

Kelt Reconditioning and Reproductive Success Evaluation Research

Project Number 2007-401-00

Report covers work performed under BPA contract # 73354 REL 45 and 73354 REL 62

Report was completed under BPA contract #73354 REL 62

Report covers work performed from: January 2021 – December 2021

Douglas R. Hatch,

Columbia River Inter-Tribal Fish Commission, Portland, OR

Report Created:

“This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the author's and do not necessarily represent the views of BPA.”

This report should be cited as follows:

Hatch, D., R. Branstetter, J. Stephenson, A. Pierce, N. Graham, J. Newell, W. Bosch, S. Everett, P. Burrows, K. Scott, L. Medeiros, L. Jenkins, C. Ray, D. Cervantes, T. Cavileer, J. Nagler, L. Caldwell, M. Fiander, C. Frederickson, J. Blodgett, and R. Johnson. Kelt Reconditioning and Reproductive Success Evaluation Research. 1/1/2021 - 12/31/2021 Bonneville Power Administration Annual Report, 2007-401-00.

Contents:

Contents:	2
LIST OF TABLES:	3
LIST OF FIGURES:	4
Abstract	6
Acknowledgments	8
Executive Introduction	9
Introduction, Problem Statement, Relevance, Goals and Objectives, and Study Area	10
SIGNIFICANCE TO FISH AND WILDLIFE PROGRAM AND OTHER REGIONAL PLANS	13
GOALS AND OBJECTIVES	14
<i>Goal 1. Develop techniques and strategies to collect, culture, care, and release wild kelt steelhead.</i>	15
<i>Goal 2. Describe the benefits to the receiving population from releasing reconditioned kelt steelhead.</i>	15
<i>Goal 3. Develop and conduct evaluations of reproductive success and homing of reconditioned kelt steelhead.</i>	15
<i>Goal 4. Carry out studies to achieve a sufficiently detailed understanding of the physiology of reconditioning in kelt steelhead to provide a scientific basis for maximizing the success of reconditioning programs.</i>	16
Methods	17
STUDY AREA	17
Steelhead Kelt Collection, Reconditioning, and Release Sites	17
Yakima River Basin	20
Snake River Basin	20
<i>Goal 1. Develop techniques and strategies to collect, culture, care, and release wild kelt steelhead.</i>	22
<i>Goal 2. Describe the benefits to the receiving population from releasing reconditioned kelt steelhead.</i>	25
<i>Goal 3. Develop and conduct evaluations of reproductive success and homing of reconditioned kelt steelhead.</i>	26
<i>Goal 4. Carry out studies to achieve a sufficiently detailed understanding of the physiology of reconditioning in kelt steelhead to provide a scientific basis for maximizing the success of reconditioning programs.</i>	28
<i>Goal 5. Project wrap up, delineation of necessary follow up monitoring and research, complete manuscripts, transition to other locations or uses of kelt reconditioning technology.</i>	30
Chapter 1: Kelt Capture, enumeration, and reconditioning in the Yakima and Snake basins. ...	31
Introduction	31
Methods	31
Results/Discussion	34
Chapter 2. Cumulative Results	49
<i>What are the benefits to the receiving population in terms of Viable Salmonid Population (VSP) parameters?</i>	50
<i>What is the reproductive success of reconditioned kelts?</i>	55
<i>What have we learned about kelt physiology and how does that inform decisions?</i>	62
<i>What is the homing fidelity of reconditioned kelts?</i>	78
<i>Where and when should we release fish?</i>	80
REPRODUCTIVE DEVELOPMENT IN WILD KELT STEELHEAD 2021	83
Introduction	83
Methods	84
Results	86
Discussion	89
Chapter 3. Building a Snake River Kelt Reconditioning Facility	92
2021 PROGRESS	92
BACKGROUND	92
Chapter 4. Project wrap up, delineation of necessary follow up monitoring and research, complete manuscripts, transition to other locations or uses of kelt reconditioning technology.	94
PRIORITIZATION OF STEELHEAD POPULATIONS FOR RECOVERY ACTIONS.	102

Adaptive Management & Lessons Learned	107
References	110
Appendices.....	130
A: MASTER KELT TRACKING TABLE.....	130
B: PUBLICATIONS.....	134
<i>Presentations 2021</i>	138
C: LIST OF METRICS AND INDICATORS	139
<i>Data Collection Methods</i>	139
<i>Data Analysis/Interpretation Methods</i>	140
<i>Metrics</i>	142
<i>Indicators</i>	143

List of Tables:

TABLE 1: KELT STEELHEAD COLLECTION, RECONDITIONING, RELEASE, AND JUVENILE COLLECTION SITES USED IN THIS STUDY. SITE NUMBER CORRESPONDS	18
TABLE 1. 1. SUMMARY OF KELT COLLECTIONS AND FISH DISPOSITION IN 2021 FROM THE SNAKE AND YAKIMA RIVERS.	35
TABLE 1. 2. SUMMARY OF FISH COLLECTIONS AND RELEASES IN THE SNAKE RIVER ASSOCIATED WITH RPA 33.	45
TABLE 2. 1. YAKIMA BASIN STEELHEAD KELTS RECONDITIONED AT PROSSER HATCHERY, NUMBER OF KELTS RELEASED, AND SURVIVAL RATE TO RELEASE, 2001 TO 2021. SOURCE: YAKAMA NATION.	51
TABLE 2. 2. SAMPLE SIZE (N), MEAN, AND GROUPING OUTPUT FOR TUKEY POST-HOC TEST FROM ANOVA OF PIT TAG DETECTIONS AT PROSSER DAM. FROM TRAMMELL ET AL. 2016.	52
TABLE 2. 3. CONTRIBUTION TO STEELHEAD SPAWNING RUN FROM RECONDITIONED KELTS RELEASED IN THE YAKIMA BASIN.	52
TABLE 2. 4. CORRESPONDING SITES BY YEAR, DETECTION, AND DETAILED LOCATION INFORMATION.	54
TABLE 2. 5. NUMBER OF JUVENILES GENOTYPED BY TRIBUTARY AND SUBSEQUENTLY ASSIGNED TO AT LEAST ONE ANADROMOUS ADULT USING PARENTAGE ANALYSIS. RESULTS ARE REPORTED FOR INDIVIDUAL YEARS AND CUMULATIVELY ACROSS ALL YEARS.	58
TABLE 2. 6. NUMBER OF GENOTYPED ADULTS WITH PIT DETECTIONS. ONLY FISH WITH BOTH A PIT TAG DETECTION AND GENOTYPES WERE INCLUDED IN THE PARENTAGE RESULTS.	60
TABLE 2. 7. NUMBER OF PROGENY ASSIGNED TO EACH TYPE OF SPAWNING EVENT WITHIN INDIVIDUAL YEARS AND ACROSS ALL YEARS. PROGENY ASSIGNED TO ADULTS WITHOUT TRIBUTARY PIT TAG DETECTIONS ARE NOT INCLUDED HERE.	60
TABLE 2. 8. AVERAGE NUMBER OF PROGENY PER PARENT SAMPLED FOR EACH CATEGORY OF FISH. RESULTS ARE REPORTED AS THE AVERAGE NUMBER OF JUVENILE DETECTIONS WITHIN EACH SPAWNING EVENT.....	60
TABLE 2. 9. RELATIVE REPRODUCTIVE SUCCESS FOR EACH CATEGORY OF FISH. RESULTS ARE NORMALIZED TO THE PRE-SPAWN MAIDEN EVEN WITHIN IN YEAR.....	61
TABLE 2. 10. LIFETIME REPRODUCTIVE SUCCESS ESTIMATE FOR MALE AND FEMALE RECONDITIONED KELTS. ADDITIVE EFFECTS OF RECONDITIONED KELT EVENTS 1 AND 2 AS NORMALIZED TO THE PRE-SPAWN MAIDEN EVENT.....	61
TABLE 3. 1. WILD STEELHEAD KELTS HELD FOR LONG-TERM RECONDITIONING FOR THE 2021 RECONDITIONING YEAR. PROSSER: PROSSER HATCHERY, DNFH: DWORSHAK NATIONAL FISH HATCHERY, WNFH: WINTHROP NATIONAL FISH HATCHERY.	84
TABLE 3. 2. WILD STEELHEAD KELTS SAMPLED DURING THE FALL OF 2021. PROSSER: PROSSER HATCHERY, DNFH: DWORSHAK NATIONAL FISH HATCHERY, WNFH: WINTHROP NATIONAL FISH HATCHERY.	85
TABLE 3. 3. WILD STEELHEAD RECONDITIONED KELTS RELEASED DURING THE FALL OF 2021. PROSSER: PROSSER HATCHERY, DNFH: DWORSHAK NATIONAL FISH HATCHERY, WNFH: WINTHROP NATIONAL FISH HATCHERY.	85
TABLE 5. 1. STEELHEAD KELT COLLECTION SITES PRIORITIZED BY HIGH QET RISK AND KELT COLLECTION POTENTIAL.	96
TABLE 5. 2. THE NUMBER OF YEARS OF AVAILABLE NATURAL ORIGIN STEELHEAD ADULT ABUNDANCE DATA REPORTED AT POPULATION LEVEL.	102

TABLE 5. 3. THE NUMBER OF YEARS OF AVAILABLE STEELHEAD JUVENILE ABUNDANCE LOCATION MONITORING DATA.	103
--	-----

List of Figures:

FIGURE 1. ANNUAL COUNTS OF UNCLIPPED STEELHEAD CROSSING BONNEVILLE DAM.	11
FIGURE 1. 1. YAKIMA RIVER STEELHEAD KELT COLLECTION FOR RECONDITIONING AND FATE FROM 2013-2021. BEGINNING IN 2019, KELTS THAT WOULD HAVE BEEN RETAINED, WERE INSTEAD TRUCKED, AND RELEASED BELOW BONNEVILLE DAM. WILL BE UPDATED IN 2022 ANNUAL REPORT.	37
FIGURE 1. 2. SNAKE RIVER STEELHEAD KELT COLLECTION FOR RECONDITIONING AND FATE FROM 2013-2021. WILL BE UPDATED IN 2002 ANNUAL REPORT.	38
FIGURE 1. 3. CONTRIBUTION TO STEELHEAD RUN FROM RECONDITIONED KELT RELEASE IN YAKIMA BASIN. * RUN IS STILL ONGOING; VALUE WILL BE UPDATED IN 2022 STEELHEAD KELT RECONDITIONING ANNUAL REPORT.	42
FIGURE 1. 4. CONTRIBUTION TO STEELHEAD RUN FROM RECONDITIONED KELT RELEASE IN SNAKE BASIN AT LGD. * INCLUDES FISH CREEK AND/OR SOUTH FORK CLEARWATER KELTS.	43
FIGURE 1. 5: CONTRIBUTION TO STEELHEAD RUN FROM RECONDITIONED KELT RELEASE IN SNAKE RIVER BASIN. X2021-2022 VALUE IS AN ESTIMATED RUN SIZE AND WILL BE CORRECTED IN THE 2021 KELT ANNUAL REPORT. B-RUN NUMBERS UTILIZE KELT MASTER PLAN DEFINITION FOR RECONDITIONED FEMALE KELT SPAWNERS.	48
FIGURE 2. 1. YAKIMA RIVER MPG STEELHEAD (PROSSER WILD ABUNDANCE) AS A PERCENTAGE OF BONNEVILLE DAM WILD GROUP A STEELHEAD ABUNDANCE, 1983 TO 2018.	53
FIGURE 2. 2. MAP OF THE SNAKE RIVER BASIN WITH YELLOW PINS INDICATING THE LAST PIT DETECTION LOCATION FOR RECONDITIONED STEELHEAD RELEASED IN 2017, 2018, AND 2019. BELOW IS THE KEY TO PIT SITES AND THE TALLY OF FISH WITH FINAL DETECTION AT EACH SITE.	54
FIGURE 2. 3. JUVENILE SAMPLING LOCATIONS IN SATUS (LOWER RIGHT) AND TOPPENISH CREEKS OVER 8 YEARS 2013-2020.	57
FIGURE 2. 4. VARIATION IN REPRODUCTIVE SUCCESS OF MAIDEN AND KELT SPAWNERS. BOTH SCALES SHOW THE NUMBER OF PROGENY ACROSS THE PORTION OF ADULTS GENOTYPED. THE SECOND PART OF THE FIGURE HAS A TRUNCATED Y-AXIS TO BETTER DEMONSTRATE THE VARIABILITY BASED ON NUMBER OF PROGENY. POSMAID= POST SPAWN MAIDEN EVENT, PREMAID= MAIDEN EVENT COLLECTED AS PRE-SPAWNER, PREKELT= FIRST SPAWNING EVENT FOR RECONDITIONED KELT, AND POSTKELT=SECOND SPAWNING EVENT FOR RECONDITIONED KELT SEE MONITORING METHODS, GENETIC ANALYSIS.	59
FIGURE 2. 5. (A) STANDARDIZED COEFFICIENTS $[(X-\text{MEAN})/SD]$ WITH 95% CONFIDENCE INTERVALS OF POTENTIAL PREDICTORS OF SURVIVAL IN A MULTIPLE LOGISTIC REGRESSION MODEL OF FEMALE STEELHEAD TROUT AT DWORSHAK NATIONAL FISH HATCHERY, AHSAHKA, IDAHO TO 70 DAYS AFTER FIRST SPAWNING IN 2015 AND 2016 (N = 291). POTENTIAL PREDICTORS WERE MEASURED AT SPAWNING. POSITIVE COEFFICIENTS INDICATE A GREATER PROBABILITY OF POST-SPAWNING SURVIVAL. COEFFICIENTS WITH CONFIDENCE INTERVALS THAT DO NOT OVERLAP ZERO WERE STATISTICALLY SIGNIFICANT ($P < 0.05$) AND ARE MARKED WITH AN ASTERISK. (B) POST-SPAWN SURVIVAL PROBABILITIES (PSURV) TO 70 DAYS, AS PREDICTED BY PLASMA TRIGLYCERIDES (MG/DL) (TG) AND PLASMA OSMOLALITY (MMOL/KG) (OS) MEASURED AT FIRST SPAWNING IN 2015 AND 2016 (N = 291). FROM JENKINS ET AL. 2020.	63
FIGURE 2. 6. SPECIFIC GROWTH RATE IN MASS (MSGR) IN FEMALE STEELHEAD TROUT SAMPLED AT 10-WEEK INTERVALS STARTING AT FIRST SPAWNING. FOOD WAS WITHHELD FROM FASTED FISH FOR THE FIRST 10 WEEKS AFTER SPAWNING. SYMBOLS INDICATE MEAN AND BARS INDICATE SEM. ABOVE EACH SAMPLING POINT, BOLD LETTERS INDICATE SIGNIFICANCE (R = REPRODUCTIVE STATUS, F = FASTING TREATMENT, R*F = INTERACTION EFFECT). ROWS OF LETTERS BELOW THE FIGURE INDICATE SIGNIFICANT DIFFERENCES OVER TIME WITHIN EACH TREATMENT GROUP. YEARS ARE SHOWN SEPARATELY DUE TO A SIGNIFICANT YEAR EFFECT AT WEEK 20-30. WITH YEARS COMBINED, FASTED FISH SHOWED SIGNIFICANTLY HIGHER MSGR THAN FED FISH OVER WEEK 10-20. FROM JENKINS ET AL. IN PREPARATION.	64

FIGURE 2. 7. MEAN (\pm SEM) LIVER INSULIN-LIKE GROWTH FACTOR BINDING PROTEIN 1 (IGFBP1) MRNA LEVELS OVER TIME IN FEMALE RAINBOW TROUT (A: TWO-YEAR-OLD; B: THREE-YEAR-OLD) FED A CONTROL-RATION OR RESTRICTED-RATION. FROM CALDWELL ET AL. 2013.	67
FIGURE 2. 8. MORTALITY OF PROSSER KELTS AFTER IVERMECTIN OR EMAMECTIN TREATMENT AT INTAKE. LOG-RANK TEST MORTALITY CURVE COMPARISON, 2011: P = 0.0019; 2012: P = 0.0340. FROM HATCH ET AL. 2013B.	69
FIGURE 2. 9. RELATIONSHIP OF PLASMA LEVELS OF ESTRADIOL (E2) AND VITELLOGENIN (VG) IN RECONDITIONED FEMALE KELTS AT PRE-RELEASE SAMPLING IN THE FALL OF 2009 (A: 10/29/2009), 2010 (B: 10/13/2010), AND 2011 (C: 10/13/2011). FISH WERE GROUPED BY CLUSTER ANALYSIS (WARD LINKAGE, 2 GROUPS SPECIFIED, DATA STANDARDIZED), AND THE RESULTING GROUPS ARE INDICATED BY CIRCLES (REMATURING FISH WITH HIGH E2 AND VG) AND TRIANGLES (NON-REMATURING FISH WITH LOW E2 AND VG). PROPOSED THRESHOLD VALUES FOR SCREENING FISH FOR MATURATION STATUS BY PLASMA E2 (500 PG ML ⁻¹) AND VG (0.1 MG ML ⁻¹) LEVEL ARE INDICATED BY LINES. FROM PIERCE ET AL. 2017.....	71
FIGURE 2. 10. (A) STANDARDIZED REPRODUCTIVE STATUS COEFFICIENTS ([X-MEAN]/SD) WITH 95% CONFIDENCE INTERVALS OF POTENTIAL PREDICTORS IN A MULTIPLE LOGISTIC REGRESSION MODEL OF REPRODUCTIVE STATUS IN CONSECUTIVE YEARS OF FEMALE STEELHEAD TROUT (N = 60) FOLLOWING MANUAL SPAWNING IN 2015 AND 2016. POSITIVE COEFFICIENTS INDICATE GREATER PROBABILITY OF CONSECUTIVE REPRODUCTIVE STATUS. COEFFICIENTS WITH CONFIDENCE INTERVALS THAT DO NOT OVERLAP ZERO WERE STATISTICALLY SIGNIFICANT (P < 0.05) AND ARE MARKED WITH AN ASTERISK. (B) THE PROBABILITY OF REPRODUCTIVE STATUS IN CONSECUTIVE YEARS IN RECONDITIONED FEMALE STEELHEAD TROUT AT DWORSHAK NATIONAL FISH HATCHERY, AHSAHKA, IDAHO, IN RELATION TO TOTAL EGG MASS AT FIRST SPAWNING (G, MASS-STANDARDIZED, N = 60). THE BARS AT THE TOP AND BOTTOM OF THE GRAPH REPRESENT THE NUMBER OF FISH THAT BECAME REPRODUCTIVELY ACTIVE OR REMAINED NON-REPRODUCTIVE FOLLOWING FIRST SPAWNING, RESPECTIVELY, FOR EACH RANGE OF TOTAL EGG MASS. FROM JENKINS ET AL. 2020.....	72
FIGURE 2. 11. SPAWN WEEK FIDELITY IN REPEAT SPAWNING FEMALE STEELHEAD TROUT FROM THE CLEARWATER RIVER, ID SAMPLED BETWEEN 2013-2015. BOXES INDICATE THE INTERQUARTILE RANGE, THE LINE INDICATES THE MEDIAN, WHISKERS SHOW THE DATA RANGE, AND ALL POINTS ARE SHOWN. REPEAT SPAWN WEEK DID NOT DIFFER SIGNIFICANTLY FROM MAIDEN SPAWN WEEK IN CONSECUTIVE SPAWNERS (ONE SAMPLE T-TEST, T ₂₂ = 0.2630, P = 0.7950), BUT WAS SIGNIFICANTLY EARLIER IN SKIP SPAWNERS (ONE-SAMPLE T-TEST, T ₁₉ = 5.4978, P<0.0001), AND REPEAT SPAWN WEEK WAS SIGNIFICANTLY EARLIER IN SKIP SPAWNERS THAN CONSECUTIVE SPAWNERS (TWO-SAMPLE T TEST, T ₄₁ =3.671, P=0.0007). FROM JENKINS ET AL. 2018.	74
FIGURE 2. 12. PLASMA ESTRADIOL-17B CONCENTRATIONS IN FEMALE STEELHEAD TROUT FROM THE CLEARWATER RIVER, IDAHO, SAMPLED IN 2015 (A) AND 2016 (B) AT 10-WEEK INTERVALS FOLLOWING MAIDEN SPAWNING. FEMALES WERE GROUPED AS REMATURING (RED: 2015, N=13; 2016, N=12) OR NON-REMATURING (BLUE: 2015, N=29; 2016, N=18). BOX HEIGHTS INDICATE INTERQUARTILE RANGE, HORIZONTAL LINES WITHIN INDICATE THE MEDIAN, AND WHISKERS SHOW THE DATA RANGE. TIME POINTS WITHIN A REMATURATION GROUP SHARING THE SAME LETTER DO NOT DIFFER SIGNIFICANTLY (ONE-WAY ANOVA FOLLOWED BY TUKEY'S MULTIPLE COMPARISON TEST, P<0.05). ASTERISKS INDICATE SIGNIFICANT DIFFERENCES BETWEEN REMATURATION GROUPS AT EACH TIME POINT (T-TEST, P<0.05). FROM JENKINS ET AL. 2019.....	76
FIGURE 2. 13. GROWTH HORMONE IN FEMALE STEELHEAD TROUT SAMPLED AT 10-WEEK INTERVALS STARTING AT FIRST SPAWNING. ANALYSIS INCLUDED FED-REPRODUCTIVE AND FED-NON-REPRODUCTIVE (CLOSED AND OPEN CIRCLES, N = 31, 35) AND FASTED-REPRODUCTIVE AND FASTED-NON-REPRODUCTIVE (CLOSED AND OPEN TRIANGLES, N = 24, 44). SYMBOLS INDICATE MEAN AND BARS INDICATE SEM. ABOVE EACH SAMPLING POINT, BOLD LETTERS INDICATE SIGNIFICANCE (R = REPRODUCTIVE STATUS, F = FASTING TREATMENT, R*F = INTERACTION EFFECT). ROWS OF LETTERS BELOW THE FIGURE INDICATE SIGNIFICANT DIFFERENCES OVER TIME WITHIN EACH TREATMENT GROUP. AT EACH SAMPLING POINT WITH A SIGNIFICANT INTERACTION EFFECT, SYMBOLS SHARING THE SAME LETTER DO NOT DIFFER SIGNIFICANTLY. FROM JENKINS ET AL. IN PREPARATION.	78
FIGURE 2. 14. SURVIVAL AND FEMALE CONSECUTIVE AND SKIP MATURATION RATES IN CRB KELT RECONDITIONING PROJECTS. SNAKE RIVER FISH INCLUDE KELTS COLLECTED AT LOWER GRANITE DAM (LGR) AND AIR SPAWNED HATCHERY ORIGIN KELTS FROM THE DNFH STOCK (DNFH HOR). MATURATION DATA FOR SKIP	

SPAWNERS IS FROM NON-MATURE FISH FROM THE PREVIOUS SEASON HELD OVER FOR AN ADDITIONAL YEAR. PROSSER 2020 CONSECUTIVE MATURATION RATES ARE NOT SHOWN, BECAUSE MATURATION WAS NOT ASSESSED BY PLASMA E2 LEVEL.	81
FIGURE 3. 1. PLASMA ESTRADIOL (E2) LEVELS IN WILD FEMALE SNAKE RIVER KELTS SAMPLED IN FALL OF 2021.....	87
FIGURE 3. 2. PLASMA ESTRADIOL (E2) LEVELS IN WILD FEMALE YAKIMA RIVER KELTS SAMPLED IN FALL OF 2021.	88
FIGURE 3. 3. PLASMA ESTRADIOL (E2) LEVELS IN WILD FEMALE UPPER COLUMBIA RIVER KELTS SAMPLED IN FALL OF 2021.	89
FIGURE 5. 1. MAP OF STEELHEAD KELT PROJECT AREA 2000-2021. NUMBERS CORRESPOND WITH SITE NUMBERS IN TABLE 1.....	19
FIGURE 5. 2. JUVENILE ABUNDANCE TREND.	106
FIGURE 5. 3. LAG 4 SPAWNERS PER SPAWNER. FIGURE SHOWS THE 4-YEAR LAG SPAWNER PER SPAWNER TREND IN LOG SPACE.....	107

Abstract

The Kelt Reconditioning and Reproductive Success Evaluation Project is a research, monitoring, and evaluation (RM&E) uncertainties category project, that was funded through the 2008 Columbia Basin Fish Accords. The objectives are to evaluate methodologies to produce viable artificially reconditioned repeat steelhead (*Oncorhynchus mykiss*) spawners and evaluate reproductive success, physiology, homing, and geographic differences. The project was recently reviewed (ISRP 2021-08) by the Independent Science Review Panel (ISRP). Our annual report contains many sections of the review document we submitted to the ISRP to provide a wider distribution of the comprehensive results. Our work occurs in both the Yakima and Snake river-basins, additionally we make some contrasts and comparisons with the Kelt Reconditioning Project in the Upper Columbia River (Project 2008-458-00). We focused on collecting steelhead kelts at juvenile bypass facilities at Prosser, Lower Granite, and Little Goose dams. These kelts were reconditioned (given prophylactic treatments and fed a specially formulated diet) at Prosser and Dworshak National Fish (DNFH) hatcheries. Survival of long-term reconditioned kelts has been 44% (22 years) at Prosser Hatchery and 37% (11 years; 44% over the last 9 years) for mixed stock collections reconditioned at both Nez Perce Tribal and Dworshak National Fish hatcheries combined. In total, we released 525 and 4,989 reconditioned kelt steelhead in the Snake and Yakima rivers since 2011 and 2000, respectively. In 2021, unmarked upstream “wild” migrant adult steelhead return counts to Bonneville Dam was the worst year since unclipped fish were recorded beginning in 1994, this places great importance on safety net programs such as the kelt reconditioning program. Years with low runs typically translate into a lower abundance of kelts, but the value of each fish increases. In 2021, we collected 530 kelts in the Snake River. From this collection of 530, 264 were PIT tagged and released back to the river, 196 were taken to DNFH for reconditioning. There were 65 reconditioned-mature (a combination of consecutive and skip spawners) fish that were released into the Snake River below Lower Granite Dam on November 10, 2021. In 2021, we collected 365 kelts in the Yakima River. From this collection of 365, 5 were PIT tagged and released back to the river, 360 were taken to Prosser Hatchery for reconditioning, and 268 reconditioned-mature consecutive spawners were released in the Yakima River below Prosser Dam on October 28, 2021. Additionally, 53 reconditioned-immature kelts were trucked and released below Bonneville Dam on October 21, 2021. Reproductive success of reconditioned steelhead was confirmed in the Yakima River tributaries of Satus and Toppenish creeks, with genetic parentage assignments

(2,700 samples from Satus and 3,087 samples from Toppenish collected from 2013 through 2019). Lifetime reproductive success for female reconditioned kelt steelhead in the Yakima River is estimated as 2.49 relative to steelhead that successfully spawn once. Studies applying tools from fish physiology and endocrinology to issues in kelt reconditioning were continued in 2021. These studies aim to achieve a sufficiently detailed understanding of the physiology of reconditioning in kelt steelhead to provide a scientific basis for maximizing the success of reconditioning programs. Screening of kelts for maturation status using plasma estradiol levels has become an essential part of the project. In 2021, we sampled blood at DNFH and Prosser hatcheries, and provided maturation status of individual fish at DNFH and Winthrop to project managers so that consecutive and skip spawners could be managed appropriately. The 2021 results were added to a comparison of the performance of the three Columbia River Basin kelt projects in terms of survival and maturation rates. In addition, we are continuing data analysis in a study where we are combining the results of a genetic analysis enabling classification of Yakima River kelts by subpopulation, with survival, physiological, and migration data. From 2008 to 2021, we have detected conclusive evidence of 146 kelts showing repeat homing and upstream movement patterns and an additional 1,414 kelts that is consistent with repeat homing in the Yakima basin. Progress continues to be made toward the realization of a dedicated kelt reconditioning facility for the Snake River in 2021. Efforts included: Review Team review of 10%, 60%, and 90% facility design documents created by R2 Resource Consultants, Inc. Development of monitoring and evaluation budget is ongoing. As the kelt program moves towards a production program we have identified some additional areas that this project can move forward on towards 3 important goals the ISRP has identified: 1. Focus on identifying and collection of at-risk populations, 2. Transferring kelt collection and rearing techniques through documentation, 3. Development of plan to provide this information to agencies that could utilize technology for at risk populations. The lessons learned and adaptive management list has continued to grow, which will provide a good outline for development of our documentation for kelt reconditioning technologies and techniques to share with agencies and groups that have interest in development of kelt reconditioning programs. Due to the ongoing COVID-19 restrictions, shutdowns of many conferences and travel have resulted in diminished opportunities for presentations, a comprehensive list of 2021 presentations are available in the appendices along with peer-reviewed articles regarding steelhead kelts produced by CRITFC and its partners along with novel additional sources.

Acknowledgments

We thank our partners at the Yakama Nation, Upper Columbia Kelt Reconditioning Program ([#2008-458-00](#)), Tom Scribner, Keely Murdoch, Matt Abrahamse, Michelle Steg-Geltner, and John Box for sharing data and sampling with our team. Many thanks to the Lower Granite Dam capture and sampling crews (Nez Perce Tribe, and Army Corps of Engineers Walla Walla District) and the US Fish and Wildlife Service and Nez Perce Tribal Hatchery staff at Dworshak National Fish Hatchery. Thank you, Lori Maxwell at Hagerman Genetic lab, for diligent and reliable laboratory processing and collection of genotypic data for this project. Also, we appreciate the help from Tim Resseguie and his team at Yakama Nation Fisheries for helping us with electroshocking and genetic collection. We thank our colleagues at CRITFC: Jeff Fryer, Denise Kelsey, David Graves, Joe Nowinski, Jayson FiveCrows, Agnes Strong, Jon Matthews, Loretta Islas, Chris Roe, Sue Wallace, Lisa Wood, Jenny Thrasher, Christine Golightly, Julie Carter, Lauren Burns, Travis Sproed, Ted Walsey, Brittney Oseth, and Amanda Kanter. Thanks to our long time collaborators at the Yakama-Nation Klickitat Fisheries Program (Zack Mays, T. Newsome, Justin Olney, JJ, OJ, and Anneliese Myers). We would like to also thank Washington Dept. of Fish and Wildlife personnel (Gabe Temple, Anthony Fritts, and Tim Webster).

Executive Introduction

Steelhead (*Oncorhynchus mykiss*) returns in the Columbia River have declined over the last two decades and counts of wild steelhead at Bonneville Dam haven't exceeded 50,000 since 2016. All steelhead stocks upstream of Bonneville are listed under the Endangered Species Act. Segregated hatcheries, mark selective fisheries, and other management strategies have failed to recover wild steelhead stocks. We need to develop new approaches aimed at restoring wild steelhead stocks. Kelt steelhead are relatively abundant in the Columbia River even in the upper most reaches, however repeat spawners represent a very small proportion of the upstream run. Developing strategies to convert the abundance of kelt steelhead into increased numbers of repeat spawners is the primary goal of the project.

This effort was initiated to explore the notion of revitalizing post-spawn wild steelhead, returning them to the river, and adding production to the population. In strong collaboration with the Yakama Nation, Nez Perce Tribe, and University of Idaho we gained knowledge of fish collection, care, and culture, measured contributions of reconditioned kelt steelhead to wild production, developed an understanding of kelt physiology, measured reproductive success, homing, and revealed life history pathways of kelt steelhead. Since 2008, this kelt reconditioning project successfully reconditioned and released 3,876 repeat spawning steelhead in the Yakima and the Snake River basins. We documented our findings in 17 published manuscripts and 84 professional presentations since 2013. The project has matured from a concept to nearly ready for production scale operations in the Yakima and Snake rivers. Future research should focus on analysis of skip spawner management strategies but other efforts such as measuring reproductive success, physiology discovery, and homing should sunset after production facilities come online.

Kelt reconditioning as a conservation tool is intended to enhance populations that have suffered decline. A near-term improvement in productivity can be a means to offset long and short-term demographic perils and minimize loss of genetic and life history diversity. Kelt reconditioning programs can work in conjunction with restoration or remediation efforts. Re-establishment or enhancement of repeat spawning in listed steelhead populations can improve productivity, diversity, and demographic stability and is particularly important during times of low steelhead abundance. The kelt project was acknowledged by National Marine Fisheries Service (NMFS 2019) as the only hatchery action in the basin that is directly improving wild steelhead abundance.

The primary goals of the project include developing and testing strategies to increase the survival and abundance of natural-origin repeat spawning steelhead in target populations while minimizing potential deleterious effects. Our proposed work is to continue addressing questions first identified by the ISRP in a 2014 review of a sister project (Steelhead Kelt Reconditioning 2008-458-00) and used in this project's last ISRP review in 2016 (ISRP 2016). These questions include: assessing how kelt reconditioning may benefit population dynamics;

evaluating the reproductive success of reconditioned kelts; achieving a sufficiently detailed understanding of the physiology of reconditioning in kelt steelhead to provide a scientific basis for maximizing success of reconditioning programs; and evaluating repeat homing of reconditioned kelt steelhead.

Introduction, Problem Statement, Relevance, Goals and Objectives, and Study Area

Columbia Basin steelhead *Oncorhynchus mykiss* have experienced a dramatic decline in abundance, resulting in listing of all distinct population segments as threatened or endangered since the late 1990's (Busby et al. 1996). A subset of the species of the family Salmonidae, including steelhead, exhibit iteroparity – the ability to spawn on two or more occasions after a period of recovery. Unfortunately, many of the factors believed to contribute to the decline of steelhead (NRC 1996; US v. Oregon 1997; ISRP 1999) have also placed significant limitations on the opportunity for successful iteroparity. Evidence suggests that kelts suffer substantial mortality during passage at mainstem hydropower facilities, in the estuary, and ocean (Evans 2002; Colotelo et al. 2012; Colotelo et al. 2013). Keefer et al. (2016) reported sea-to-sea survival of kelts from the Columbia and Snake rivers between 1% to 2%. They found kelt outmigration survival from the Snake River to below Bonneville Dam ranged from 13% to 20%. Kelt survival may also be limited by an increase in energy expenditure accompanying alteration of freshwater habitat (Love 1970; Penney and Moffitt 2014a). In the pre-dam era, repeat spawners were more prevalent, Long and Griffin (1937) reported 3.4% of scales from 1,128 steelhead collected in the lower Columbia River exhibited a spawn check. In contrast, iteroparity for upriver steelhead was estimated as 0.5% based on scales of 24,112 fish sampled at Bonneville Dam from 2004 to 2020 (Jeff Fryer personal communication). In less disturbed rivers such as the Nass and Skeena in British Columbia, contemporary repeat spawners make up an estimated 8% of the spawning run (Moore et al. 2014). In the Kalama River, an un-impounded tributary of the lower Columbia River below all mainstem hydropower facilities, it was documented that 17% of the run were repeat spawners in the 1990's (NMFS 1996).

In general, efforts to maintain or recover declining steelhead stocks within the Columbia River Basin have ranged from harvest reduction, mark-selective fisheries, habitat restoration, passage improvements (including transporting fish) at mainstem Columbia River hydro-power facilities, and hatchery propagation. These strategies have not recovered steelhead populations (Figure 1). Clearly, we need innovative approaches to obtain recovery of wild steelhead populations. Recently, the use of kelt reconditioning has been investigated as a novel method to increase total reproductive potential of steelhead populations (Hatch et al. 2013b). Kelt reconditioning was adopted as an offsite mitigation action in the 2008 Biological Opinion on the Federal Columbia River Power System as part of Reasonable and Prudent Alternative (RPA) 33 (NMFS 2008). The 2019 BiOp lists kelt reconditioning as an important strategy for wild steelhead recovery in the Columbia and Snake rivers ([NMFS 2019](#)).




Figure 1. Annual counts of unclipped steelhead crossing Bonneville Dam.

This Kelt Reconditioning proposal is intended for application towards steelhead populations that have suffered declines in abundance and are targeted for restoration. In that context, an immediate boost in freshwater productivity is a means to offset long and short-term demographic perils and minimize loss of genetic and/or life history diversity. Preferably, kelt reconditioning programs should be accompanied by restoration or remediation targeting the root cause(s) of decline (Seamons and Quinn 2010). In evaluating restoration actions for steelhead, it should be noted, that the perils faced by iteroparous species may, or may not align perfectly with those faced by semelparous species, thus remediation targeting limiting factors for semelparous species may not be equally effective for iteroparous species. For example, habitat restoration actions have not explicitly focused on attributes that might increase kelt survival. Similarly, fish passage projects at mainstem hydropower facilities generally focus on upstream adult passage improvements, with far less emphasis on guidance and passage for emigrating adults, although this trend appears to be changing (Wertheimer et al. 2008). Thus, it is possible that existing restoration projects may be sufficient to secure demographic stability of a targeted population without simultaneously reestablishing historic rates of kelt contribution. Whether “asymmetric” recovery, as described above, is detrimental, has yet to be resolved, and is directly related to uncertainty about the genetic basis of iteroparity and its value as a life-history strategy.

It is unclear whether iteroparity is a genetic trait subject to loss within a population or Distinct Population Segment (DPS) versus a ubiquitous characteristic of the species whose expression is governed solely by environmental influence (i.e., a trait that cannot be “lost”; Seamons and Quinn 2010).

Several publications posit the theoretical benefits of iteroparity (Groot and Margolis 1991, Crespi and Teo 2002; Niemälä et al. 2006) and some document a realized increase in lifetime reproductive success accompanying iteroparity (Seamons and Quinn 2010); it is generally assumed that a loss or reduction in successful iteroparity simply results in a decrease in potential productivity. However, some recent evidence suggests that the loss of successful iteroparity can have profound impacts on not only productivity, but diversity, and population stability as well. In a recent study on British Columbia summer steelhead, Moore et al. (2014) found that increasing prevalence of repeat spawners reduced the magnitude of population fluctuations. Prevalence of repeat spawners had a larger effect on population stability than freshwater and ocean residence times, suggesting that there are substantial benefits of repeat spawners to population stability. Seamons and Quinn (2010) illustrate all these potential benefits. The authors studied a small population of coastal steelhead and found that on average, female repeat spawners had 1.9 times the reproductive success of females that spawned only once, while male repeat spawners produced nearly 2.7 times the number of adult offspring as males that spawned only once. Generally, these results support other empirical and theoretical observations regarding the potential benefits of iteroparity. Perhaps more importantly, the authors found that 71% of female repeat spawners typically produced the bulk of their surviving progeny in either the first or the second spawning event while only 29% of female repeat spawners successfully reproduced during two or more spawning events. Among repeat spawning males, none of the repeat spawners produced offspring during their first reproductive attempt; those that did successfully reproduce in their second attempt, produced nearly three times as many adult offspring as males that spawned on their first attempt. Although this is a single study of one population of steelhead, if generally applicable, the observations profoundly underscore the risks associated with a reduction in iteroparity; namely:

- Successful iteroparity can increase individual lifetime reproductive success by two to three times relative to semelparous individuals.
- A non-negligible fraction of male and female spawners fails to reproduce during their first spawning attempt, suggesting that genetic diversity could be directly reduced when conditions do not support successful iteroparity.

Aside from the documented benefits of iteroparity with regard to productivity, diversity, and demographic stability; reestablishment or enhancement of iteroparity has several potential management benefits:

1. Increased iteroparity improves fishery opportunities by increasing the total number and mean size of returning adults potentially vulnerable to fisheries.

2. Kelts can be reconditioned and held for use as broodstock, thus decreasing the demographic toll of broodstock collection on escapement for natural spawning.
3. Management actions to increase successful iteroparity may be implemented quickly and could achieve a demographic boost less controversially than artificial propagation and more quickly than habitat or passage improvements.
4. Reconditioning efforts can be focused on natural origin steelhead, thus substantially reducing chance for possible deleterious hatchery effects.

Significance to Fish and Wildlife Program and other regional plans

The principle of co-management is central to the Columbia Basin Fish and Wildlife Program, and the Northwest Power and Conservation Council is committed to working collaboratively with Tribes and in accordance with federal and state fish and wildlife laws and regulations (NPCC 2014). Sovereignty, and the legal rights of Tribes founded in treaties and tribes are understood to have governmental privileges over their lands, people, and natural resources. The United States has a trust obligation to protect the aforementioned and the Northwest Power Act (NPCC 1980) aims to assist tribes in realizing their rights and responsibilities in regard to fish and wildlife.

Steelhead kelt reconditioning is part of the [Technical Recommendation 4 Supplementation](#) in Spirit of the Salmon: WY-KAN-USH-MI WA-KISH-WIT the tribes salmon recovery plan.

Steelhead kelt reconditioning programs in the Snake and Yakima rivers are included in NMFS recovery plans and in biological opinions:

- [Page 192 and Appendix C Page 110](#) of the Snake River Steelhead Recovery Plan calls to implement the Kelt Management Plan and to continue to recondition Snake River steelhead kelts to benefit B-run populations. This project directly implements kelt reconditioning on B-run steelhead in the Snake River Basin.
- [Page 2-1 and 2-6](#) in the Mid-Columbia steelhead recovery plan considers the Yakima River Kelt Reconditioning Program part of the DPS and MPG. Page 7-6 calls to develop and implement a kelt management plan. Page 7-9 recommends maintaining the existing kelt reconditioning program. Table 7-4 Recovery Strategies and Actions for the Yakima Basin MPG (page 7-35) lists the highest priority strategies that includes promoting repeat spawning of kelts, it affects all populations in the Yakima River, the key action would be using the kelt reconditioning program, it addresses all VSP parameters, and it addresses the reduced survival of kelts due to passage conditions in the Columbia and lower Yakima Rivers. Within the Yakima River specific recovery plan, Kelt Reconditioning is listed as a recovery strategy for all Yakima River populations (page 131) and continuing the program is listed as Basinwide Action #6 (page 148).

- The [NMFS 2019 CRS Biological Opinion](#) supports the Action Agencies to continue funding kelt reconditioning in the upper- and mid-Columbia as well as the Snake River on pages 43, 196 and 197.
- [Snake River Kelt Management Plans and Progress Reports 2009-2018](#) (NOAA, BPA, COE documents 2009, 2011, 2013, 2015). Annual Management Plan and progress reports related to achieving the RPA 33 objective of increasing B-run steelhead in escapement in the Snake River. These include ongoing efforts, knowledge, management and research priorities within the framework of adaptive management. Page [24](#) of the 2015 report shows long term kelt reconditioning produced significantly higher rates of repeat spawners than other management strategies.
- [The 2015 Snake River Basin Steelhead Kelt Reconditioning Facility Master Plan pages 16-24](#). Alterations to the Columbia River hydropower system have done little to improve the survival of emigrating steelhead kelts. As demonstrated, transport and short-term reconditioning appear to improve survival to the estuary but result in marginal increases in iteroparity rates. Therefore, we conclude that these strategies are unlikely to achieve the survival benefits required by RPA 33. Available evidence suggests that long-term kelt reconditioning is the most promising kelt enhancement strategy for Snake River B-run steelhead to achieve RPA 33 which is a 6% increase in escapement in the Snake River.
- [Yakima Basin Steelhead Reconditioning Project HGMP](#).

Steelhead kelt reconditioning guidance from the Council is found in many sources:

- Supporting the kelt steelhead reconditioning efforts in the Yakima and Snake river basins is stated as an accomplishment from implementation of the 2014 Fish and Wildlife Program in the [2020 Addendum, page 5](#).

The [2017 Research Plan and Critical Uncertainties](#) document (NWPCC 2017) does not call out kelt reconditioning specifically but do provide guidance for assessing propagation programs. This guidance is useful and integrated into our program to assess kelt reconditioning in terms of population growth, abundance, spatial structure, and diversity as well as potential fitness impacts to the receiving population.

Goals and Objectives

The overall goal of the Kelt Reconditioning Project is to improve wild steelhead populations by increasing the abundance of natural repeat spawners. If this research points to a positive response, we plan to move to production level implementation of kelt reconditioning. To achieve this goal, we have identified the following goals, objectives, and milestones:

Goal 1. Develop techniques and strategies to collect, culture, care, and release wild kelt steelhead.

Objective 1.1 Evaluate fish collection locations and strategies to increase the abundance of kelt steelhead for reconditioning (designed capacity ~600 kelts) at the new Snake River Facility (once it is built).

Objective 1.2 Evaluate fish culture and care strategies to improve survival and rematuration above the long-term averages for each facility (~50%), continue until new facilities are built.

Objective 1.3 Continue to search for and evaluate fish characteristics at collection that correlate with survival and rematuration of artificially reconditioned kelts to achieve repeat spawners rates that are higher than the natural repeat rate in the receiving population (~3% in the Yakima and 0.5% in the Snake).

Objective 1.4 Evaluate strategies to manage skip spawners that optimize survival and spawning contribution, available space, and cost. This is planned to continue until new facilities are constructed.

Goal 2. Describe the benefits to the receiving population from releasing reconditioned kelt steelhead.

Objective 2.1 Evaluate receiving steelhead population abundance and growth and resulting contributions from released reconditioned kelts. This is planned to continue until new facilities are constructed.

Objective 2.2 Evaluate spatial structure and diversity of reconditioned kelts by tracking PIT detections throughout the basin. This is planned to continue until new facilities are constructed.

Goal 3. Develop and conduct evaluations of reproductive success and homing of reconditioned kelt steelhead.

Objective 3.1: Collect tissue samples from representative parent and offspring fish groups in the Yakima River through 2025.

Milestone 1. Completed annual juvenile collections through 2022.

Milestone 2. Completed annual adult collections through 2025.

Objective 3.2: Calculate relative reproductive success using parentage analysis from Yakima River collections from 2022-2025.

Milestone 1. Completed annual parentage analysis of juvenile samples.

Milestone 2. Submission of manuscript for reproductive success by juvenile assignments to a scientifically accredited journal by 2022.

Milestone 3. Submission of manuscript for adult-to-adult reproductive success to an appropriate journal by 2026.

Objective 3.3: Evaluate the homing and straying rates of reconditioned kelts in the Yakima River Basin. Monitor and analyze the detection history for verifiable first and second spawnings for all reconditioned kelts to estimate quantifiable homing and straying rates.

Milestone 1: Complete and submit a manuscript describing homing of reconditioned kelt steelhead to an appropriate journal by 2022.

Goal 4. Carry out studies to achieve a sufficiently detailed understanding of the physiology of reconditioning in kelt steelhead to provide a scientific basis for maximizing the success of reconditioning programs.

Objective 4.1: Use blood sampling and plasma reproductive hormone assays to evaluate reproductive status of kelts in reconditioning programs.

Milestone 1: Every year, collection of blood samples in September, assay of plasma estradiol or 11-ketotestosterone level, and assignment of each individual as a consecutive or skip spawner before fish are released in November.

Objective 4.2: Use physiological tools in studies to improve survival and rematuration rates at kelt reconditioning facilities (with Objective 1.2).

Objective 4.3: Use physiological tools to evaluate fish characteristics at collection that correlate with survival and rematuration at kelt reconditioning facilities (with Objective 1.3). If such predictors can be identified, they could be used to select fish for reconditioning.

Objective 4.4: Evaluate physiological indices at release as predictors of post-release migration and spawning success in kelts released from reconditioning programs. If such predictors can be identified, modifications to reconditioning practices can be tested to maximize post-release fish performance.

Goal 5. Project wrap up, delineation of necessary follow up monitoring and research, complete manuscripts, transition to other locations or uses of kelt reconditioning technology.

Objective 5.1: Develop plan for necessary monitoring and research to implement after kelt reconditioning production facilities at Prosser and Nez Perce Tribal hatcheries are completed and online.

Objective 5.2: Complete and submit manuscripts and technical reports related to kelt physiology, reproductive success, and homing from 2021-2023.

Milestone 1: Complete and submit manuscript describing kelt fasting experiment conducted using DNFH hatchery kelt model to a scientifically accredited journal by 2021-2022.

Milestone 2: Complete statistical analysis and writeup of study on post-spawning ration restriction using post-spawning rainbow trout model by 2021-2022, followed by submission to a scientifically accredited journal.

Milestone 3: Complete data processing, analysis and writeup of study on predictors of post-release migration and spawning success in female kelts release from Prosser Hatchery by 2022-2023. Submission to a scientific journal or publication as a technical report to depend on results.

Objective 5.3: Survey the region for potential to transfer kelt reconditioning technology to other locations and applications such as hatchery broodstock management or reintroductions.

An additional section with rationale for each goal, objective, and milestone is provided after the description of the study area.

Methods

A list of methods is provided in the [Appendix C](#). This list provides direct hyperlinks to detailed project methods that are hosted on the [Monitoring Methods website](#).

Study Area

Steelhead Kelt Collection, Reconditioning, and Release Sites

The study area is focused in the Yakima and Snake rivers. In the Yakima River, kelts are collected at the Chandler Juvenile Collection Facility (CJCF) at rkm 76 located downstream of all known steelhead spawning areas (Table 4.1 and Figure 4.1). The CJCF is situated on an irrigation canal where approximately 50% of the outmigrating kelts pass. Kelts are transferred from the CJCF to the adjacent Prosser Hatchery for reconditioning. Juvenile collections for the relative reproductive success work are concentrated in Satus (rkm 27-62) and Toppenish (rkm 40-78) creeks.

In the Snake River, kelts are collected at the juvenile bypass screens at Lower Granite Dam (rkm 173) and more recently at Little Goose Dam (rkm 113). These fish are trucked, then reconditioned at the Dworshak National Fish Hatchery, located at the confluence of the North Fork Clearwater River (rkm 65). Additional kelt rearing space is available at Nez Perce Tribal Hatchery (rkm 38), which is also the site of the future permanent Snake River kelt reconditioning facility (Hatch et al. 2021).

This project previously collected steelhead kelts at numerous sites around the Columbia River Basin. These sites are detailed in previous annual reports (Hatch et al. 2020, Hatch et al. 2015, Hatch et al. 2013a, Hatch et al. 2012, and Branstetter et al. 2008).

Table 1: Kelt steelhead collection, reconditioning, release, and juvenile collection sites used in this study. Site number corresponds with numbers in Figure 1.

Site Number	Site	Drainage	Location	Collection site	Reconditioning site	Release Site	Juvenile Sampling Location	Dates of use
1	Chandler Juvenile Collection Facility (CJCF)	Yakima River	Rkm 76	Yes	-	-	-	1999-2021
2	Yakama Nation Prosser Fish Hatchery	Yakima River	Rkm 76	-	Yes	Yes	-	1999-2021
3	Satus Creek	Yakima River	-	-	-	-	Yes	2008-2021
4	Toppenish Creek	Yakima River	-	-	-	-	Yes	2008-2021
5	Dworshak National Fish Hatchery (DNFH)	Clearwater River	Rkm 65	Yes, (hatchery fish for experimental purposes) 2009-2018	Yes	-	-	2009-2021
6	Nez Perce Tribal Fish Hatchery (NPTH)	Clearwater River	Rkm 38	No	Yes	-	-	2016-2019
7	Lower Granite Dam Juvenile Bypass (LGDJB)	Snake River	Rkm 173	Yes	-	Yes	-	2009-2021
8	Little Goose Dam Juvenile Bypass (LGSDJB)	Snake River	Rkm 113	Yes	-	-	-	2020-2021



Figure 5. 1. Map of Steelhead kelt Project area 2000-2021. Numbers correspond with site numbers in Table 1

Yakima River Basin

The [Yakima River](#) is approximately 344 km in length and enters the Columbia River at rkm 539. The basin is 15,928 km² and average discharge is 99 m³/s. Summer steelhead populations primarily spawn upstream from Prosser Dam in Satus Creek, Toppenish Creek, Naches River, and other tributaries of the Yakima River (TRP 1995).

Chandler Juvenile Collection Facility (Yakima River)

Some post spawn steelhead (approximately 20%) migrating downriver are entrained in an irrigation canal and collected at the Chandler Juvenile Monitoring Facility (CJMF a.k.a. Chandler Juvenile Evaluation and Monitoring Facility CJEMF)) that screens migratory fishes away from the canal. The entire kelt collection for the Yakima River is conducted at the CJMF.

Yakama Nation Prosser Hatchery

Prosser Hatchery is located on the Yakima River just downstream of Prosser Dam (rkm 75.6) and adjacent to the CJMF. This facility is part of the The Yakima/Klickitat Fisheries Project, a supplementation project designated by the NPPC as the principle means of protecting, mitigating, and enhancing the anadromous fish populations in the Yakima and Klickitat Subbasins. Prosser Hatchery was constructed in 1994 with the primary function of rearing, acclimating, and releasing fall chinook salmon (*O. tshawytscha*). It is also used for rearing coho salmon (*O. kisutch*) prior to acclimation and release in the upper Yakima River Basin as well as experimental rearing of white sturgeon (*Acipenser transmontanus*) and Pacific lamprey (*Entosphenus tridentate*).

Snake River Basin

The [Snake River](#) watershed is the tenth largest among North American rivers and covers almost 280,000 km² in portions of six U.S. states: Wyoming, Idaho, Nevada, Utah, Oregon, and Washington, with the largest portion in Idaho. Most of the Snake River watershed lies between the Rocky Mountains on the east and the Columbia Plateau on the northwest. The largest tributary of the Columbia River, the Snake River watershed makes up about 41% of the entire Columbia River Basin. The Snake River enters the Columbia at rkm 523. Its average discharge at the mouth constitutes 31% of the Columbia's flow at that point. The Snake River's average flow is 1,553 m³/s. At Anatone, Washington, downstream of the confluences with the Salmon and Grand Ronde, but upstream of the Clearwater, the mean discharge is 979 m³/s. Steelhead spawn naturally throughout the lower portion of the basin with the vast amount of "B-run" steelhead produced at the Dworshak National Fish Hatchery found on the Clearwater River.

The Lower Granite Juvenile Fish Facility

The third dam on the Snake River, the Lower Granite Lock and Dam is a concrete gravity run-of-the-river dam on the Snake River, in the U.S. state of Washington. The dam is located 35 km (22 miles) south of the town of Colfax, and 35 miles (56 km) north of Pomeroy ([Wikipedia](#)).

Steelhead kelts migrating from tributaries of the Snake River above Lower Granite Dam that do not emigrate via the Removable Spillway Weir (RSW) are directed by a large bypass system to the Juvenile Fish Facility (JFF) at Lower Granite Dam (LGR) (rkm 173).

The Little Goose Juvenile Fish Facility

The Second Dam on the Snake River, the Little Goose Lock and dam is a concrete gravity run-of-the-river dam on the Snake River, in the U.S. state of Washington. The dam is located 14km (9mi) northeast of the town of Starbuck, and 40km (25 mi) north of Dayton ([Wikipedia](#)).

Steelhead kelts migrating from tributaries of the Snake River above Little Goose Dam that do not emigrate via the Removable Spillway Weir (RSW) are directed by a large bypass system to the Juvenile Fish Facility (JFF) at Little Goose Dam (LGS) (rkm 113).

Dworshak National Fish Hatchery

Kelt reconditioning facilities are located at [Dworshak National Fish Hatchery](#) (DNFH) in Ahsahka, Idaho. DNFH is located at the confluence of the North Fork of the Clearwater River (rkm 65).

The Dworshak National Fish Hatchery is a "mitigation" hatchery constructed in 1969 by the Army Corps of Engineers, which presently is co-managed by the U.S. Fish and Wildlife Service and the Nez Perce Tribe (USFWS 2009). Kelts from Lower Granite and hatchery origin fish have been reconditioned at this facility since 2009. Beginning in 2016 most of the kelts reconditioned at this location were hatchery fish that returned to the hatchery. They are then air spawned and reconditioned to learn more about kelt rematuration and how we can improve maturation and survival. Through 2019 besides the experimental groups surplus Lower Granite Dam captured "wild" kelts were held on site and trucked to NPTH in the fall. In both 2020 and 2021 all the Lower Granite Dam and Little Goose kelts collected and selected for reconditioning were trucked and reconditioned at this location due to technical issues (water sourcing) at NPTH.

Nez Perce Tribal Hatchery

Starting in mid-2016 kelt reconditioning tanks were established at the Nez Perce Tribal Fish Hatchery site situated at Nez Perce Tribal allotment site 1705, located 38 km above the mouth of the Clearwater River. This Nez Perce Tribe managed facility was constructed in 2002 and was primarily used to supplement spring and fall chinook (*O. tshawytscha*) in the Clearwater River. The majority of steelhead kelts captured at Lower Granite Dam are trucked here to be reconditioned. No fish were reconditioned here in 2020 and 2021 due to technical issues that need to be resolved before reconditioning can resume (Hatch et al 2020). This is the future site of a permanent steelhead kelt reconditioning facility which should resolve the water sourcing issues.

Goal 1. Develop techniques and strategies to collect, culture, care, and release wild kelt steelhead.

Goal 1, Objective 1.1: Evaluate fish collection locations and strategies to increase the abundance of kelt steelhead for reconditioning (designed capacity ~600 kelts) at the new Snake River Facility (once it is built and operational; scheduled 2023) and in the Yakima River.

We have operated kelt reconditioning at a research scale by collecting fish at the Chandler Juvenile Collection Facility (CJCF) and Lower Granite Dam. Production scale facilities with larger capacities may require collections at additional locations to obtain enough fish to fill the new facilities. To locate suitable collections sites, we plan to:

- Develop collection site criteria: access to kelts, abundance of kelts, ability to hold kelts prior to transport to reconditioning facility, potential conflicts with other programs, and other considerations.
- Consult co-managers and maps for potential collection sites and conduct site visits.
- Develop implementation plans for additional collection sites.

Goal 1, Objective 1.2: Evaluate fish culture and care strategies to improve survival and rematuration above the long-term averages for each facility (~50%), continue until new facilities are built.

Kelt steelhead are collected from the separator at juvenile bypass screens by dipnetting and transfer to a temporary holding tank containing oxygenated well water (13.8°C). Each fish is anesthetized in a buffered solution of tricaine methanesulfonate (MS-222) at 600 µL/L, weighed, measured for FL, and judged by experienced fish culture staff as to maturation status, sex, condition (good = 1; fair = 2; poor = 3) and coloration (bright = 1; intermediate = 2; dark = 3). Condition is based on the degree of visible external damage (e.g., abrasions, lesions, fungal infections; see Evans 2003), and coloration (an indicator of physiological state) was based on the degree of the fish's silvery, ocean-like external appearance. A uniquely coded PIT tag is injected into the pelvic girdle of each kelt (Prentice et al. 1990) for individual fish identification during reconditioning and post release tracking.

Reconditioned kelts are fed a combination of frozen krill and 6.0-mm pellet feed manufactured by Bio-Oregon. Krill is used initially to enhance the feeding response, and after 4–6 weeks the pellets are introduced. Feed is administered 3–5 times daily at a rate of 1–2% body weight or until fish seemed satiated. The tanks were covered to provide shelter from sun and relieve stress from outside movement. Any kelt mortalities were removed daily and the tanks were swept and flushed every 10–14 d as needed. The tank walls were painted white and the centers dark to discourage the kelts from rubbing the walls. We found eye damage became prevalent when fish excessively rubbed the walls (Hatch et al. 2013a). Aerators were placed in the tanks to break up the surface and introduce oxygen. The aerators appear to reduce stress by providing added security for the fish.

Before their release from the reconditioning facility, surviving steelhead are again weighed, measured (FL), sexed, and scanned for PIT tags. Fish are released from mid- October to early December, concurrent with the peak return of the natural spawning run. Releases are made downstream of both Prosser and Lower Granite dams for Yakima and Snake river origin fish, respectively. The release location and timing allow reconditioned kelts to naturally select their migration timing, spawning location, spawn timing, and mates.

Our existing fish care protocols (Hatch et al. 2013a) are achieving survival rates to the first fall around 50%. We plan to continue refining these protocols calculating and tracking survival, rematuration and making comparisons among locations described in Hatch et al. (2021).

Currently, at Dworshak National Fish Hatchery (DNFH) we are using Parasite-S (Formalin) from Syndel to control fungus while reconditioning steelhead kelts. The fungicide is delivered to the individual tanks with a MasterFlex L/S peristaltic pump and C-Flex Tubing LS/25 from Cole-Parmer. The tubing is housed in 3/4" PVC from the chemical storage room to the point of delivery in the tanks. The tanks are 15' round, water depth of 4', and an inflow of 60-65gal/min. The base treatment used is a flow-through treatment of 23.5ml/min of Parasite-S for 2hrs 46min. All incoming kelts receive daily treatments for approximately 6 months. At this point, the kelts are either released or held as skip spawners. All skip spawning kelts receive a treatment every-other day until they are released the following year.

Starting in May 2021, we will attempt to switch from Parasite-S to a 35% Perox-AID (hydrogen peroxide) product from Syndel to control fungus. Treatments will be administered at the same rate, duration, and utilize the same delivery method as the Parasite-S treatments, starting with the 2020 skip spawners. All fish in this treatment will receive every-other day treatments from May through release in October. If the treatment is successful, all steelhead kelts at this location will be transferred to this new fungicide.

Goal 1, Objective 1.3: Continue to search for and evaluate fish characteristics at collection that correlate with survival and rematuration of artificially reconditioned kelts to achieve repeat spawner rates that are higher than the natural repeat rate in the receiving population (~3% in the Yakima and 0.5% in the Snake).

All captured steelhead are scanned for existing PIT-tags, and biological data is collected as described above. To evaluate the change in fish form as a result of reconditioning, we calculate Fulton's condition factor (K) at collection and at release (Ricker 1975). We use the calculation $K = w/l$, where w = fish weight (kg) and l = fish length (cm). One-way analysis of variance (ANOVA) is used to determine differences, at collection, in kelt length, weight, condition, and color across years (Sokal and Rohlf 2000). We use ANOVA tests to evaluate the association of survival (0, 1) with several fish trait variables, including FL, weight, K, condition, and color. Additionally, we calculate correlation coefficients among fish variables along with annual abundance of kelt and prespawn steelhead, the proportion that kelt steelhead made up of the previous run, length and weight change, and standardized flow periods (Sokal and Rohlf 2000).

Goal 1 Objective 1.4: Evaluate strategies to manage skip spawners that optimize survival and spawning contribution, available space, and cost. This is planned to continue until new facilities are constructed.

Long-term reconditioning is a management strategy where emigrating kelt steelhead are collected and held in large tanks, given prophylactic treatments and fed a specially formulated diet for approximately 6 months (Hatch et al. 2013a). After 6 months, the “reconditioned” kelts are released back into the collection river as the run at large is returning from the ocean. These reconditioned fish generally mingle with the run at large and proceed to in-river, over-winter locations and spawning grounds in the spring. We have determined that both natural and artificially reconditioned kelts can pursue two alternative pathways toward rematuration and repeat spawning. One pathway is termed consecutive spawning where individuals remature and proceed to spawn in the next spawn cycle. The other pathway is termed skip spawning where individuals remature and proceed to spawn two years after their previous spawning. To illustrate, kelts collected in the spring of 2020 could spawn again in the winter/spring 2021 as consecutive spawners or wait until spring of 2022 and spawn as skip spawners. The proportion of consecutive and skip spawners in a cohort varies annually, but in general, Yakima River fish predominately follow the consecutive spawner pathway (60-70%) and will spawn the following winter/spring, while the majority of Snake River kelts follow the skip spawner life history (60-70%). Strategy choice may be controlled by a combination of genetics and environment, while also varying from year to year. We plan to develop and evaluate alternative management strategies for skip spawners.

Our skip spawner management approach in the Snake River has been to hold the fish in the hatchery for an additional 12 months. This has been very successful in terms of survival, >85% for the second year. In the Yakima River, the Prosser Hatchery kelt tanks operate on well water that in the winter is much warmer than the river. This relatively warm water disrupts the spawning schedule of skip spawners held for the additional 12 months. The Prosser Hatchery Master Plan calls for a recirculating / chilled water system to allow us to successfully hold skip spawners over the winter, but in the meantime, we are searching for an alternative skip spawner strategy for Yakima River kelts until the new facility is constructed.

Skip spawner management strategies to evaluate include:

1. Rear for an additional 12 months in the hatchery. This is the current Snake River approach and the survival and rematuration rates will be used as the baseline values to evaluate other alternatives.
2. In the fall at the time when consecutive spawners are released, transport Yakima River fish that are on skip spawner trajectory to the lower Columbia River and release. Return rates, based on PIT detections through the hydrosystem will be used as an index of repeat spawner rates and compared against strategy 1. Based on recent steelhead returns to the Yakima River, we anticipate a sample size of 75 to 100 fish in the transport group.
3. To increase the sample size of the transport group, we plan to explore other sources of reconditioned kelt steelhead for the transport test. A possible, yet currently unexplored, source of

kelts would be Dworshak Hatchery broodstock (or some other steelhead hatchery). If these fish were available, they could be air spawned in the spring, held, and reconditioned until fall, transported (for skip spawners), and evaluated by PIT detection on their return. The sample size for this group could be approximately 200 fish. This experiment is merely a possibility and will require extensive coordination with co-managers including the Fish and Wildlife Service.

Goal 2. Describe the benefits to the receiving population from releasing reconditioned kelt steelhead.

Goal 2. Objective 2.1: Evaluate receiving steelhead population abundance, growth, and resulting contributions from released reconditioned kelts. This is planned to continue until new facilities are constructed.

We plan to continue to evaluate the contribution of reconditioned kelts to receiving populations. Evaluations in the Yakima River will include ocean return steelhead counts at Prosser Dam, the number of mature reconditioned kelts released in October, and tracking the proportion of the run that is comprised of reconditioned kelts. We plan to continue tracking the proportion of the wild group A steelhead run at Bonneville Dam that is made up of Yakima River fish. The proportion will be calculated from counts of wild group A steelhead at Bonneville Dam and steelhead counts at Prosser Dam.

We will use a life cycle model to predict the effect of reconditioning and releasing a range of proportions of kelts. The predictions will account for the condition of released kelts, and the resulting probability of successfully spawning. Progeny will be assumed to contribute to the naturally spawning population. A sensitivity analysis of the rate of contribution of kelts to naturally spawning steelhead will be limited to the capacity of the reconditioning facility such that the analysis can account for a variable facility capacity.

Goal 2, Objective 2.2: Evaluate spatial structure and diversity of reconditioned kelts by tracking PIT detections throughout the basin. This is planned to continue until new facilities are constructed.

The growing network of in-stream PIT tag arrays provide an opportunity to track reconditioned kelts to many spawning locations in the Snake and Yakima rivers. We plan to use this network to characterize the spatial distribution of kelts, and to map the contribution of reconditioned kelts to abundances in streams all over the basin.

An annual record of kelt detections at each PIT tagged stream will be maintained, providing an annual record of the proportion of reconditioned kelt releases arriving at each receiving population stream. The record will also provide an indication of the proportional contribution of reconditioned kelts to each stream's spawning relative to naturally spawning maidens. The proportional distribution of kelts (spatial variability) across all receiving populations will provide an indication of the spatial diversification effect of kelt releases.

Goal 3. Develop and conduct evaluations of reproductive success and homing of reconditioned kelt steelhead.

Goal 3, Objective 3.1: Collect tissue samples from representative parent and offspring fish groups in the Yakima River through 2025.

Anadromous adult steelhead will be collected as either upstream migrants at Prosser Dam or downstream migrants at the Chandler Juvenile Collection Facility. Samples collected as upstream migrants at Prosser Dam will be treated as maidens and referred to as pre-spawn maiden collections. Post-spawn adults sampled at the Chandler facility that did not go into the reconditioning program or that did not survive the reconditioning program are referred to as post-spawn maidens. Adults collected at the Chandler facility and reconditioned prior to release in the fall are referred to as kelts.

Age-0 juveniles (juveniles collected in the same calendar year as the spawning event) will be targeted using electrofishing techniques (NMFS 2000 Electrofishing Guidelines) during the late summer and fall in the Satus and Toppenish tributaries. Additional samples from rotary screw traps may be included. Sampling will be targeted near areas where steelhead spawning has been observed or previous sampling efforts have provided progeny with anadromous parents. Technicians in the field will target age-0 juveniles with a general maximum length of 100mm in addition to the judgment of those collecting the samples based on the time of collection. Individual Fork length will be recorded.

Milestone 1. Completed annual juvenile collections through 2022.

Milestone 2. Completed annual adult collections through 2025.

Goal 3, Objective 3.2: Calculate relative reproductive success using parentage analysis from Yakima River collections from 2022-2025.

Genetic analysis will be conducted at the Hagerman Fish Culture Experiment Station in Hagerman, ID. DNA will be extracted from tissue samples using chelex beads and genotype data generated with a lab standardized panel of SNP markers using a targeted amplicon GT-seq method on a next generation sequencer. Prior to parentage analysis, a sex-determining marker (OmyY1_2SEX), three loci diagnostic for cutthroat, poor performing loci and loci with low minor allele frequency will be dropped. Unresolved duplicate samples, samples with incomplete genotypes, and non-target species samples will be omitted.

Parentage analysis will be performed using CERVUS v 3.0 (Marshall et al. 1998, Kalinowski et al. 2007). Information on fish gender is not included in the analysis. To minimize incorrect assignments, simulations will be performed annually to estimate a 99.0% confidence LOD value. Individual parentage assignments will be included if they have a minimum of 90% loci comparisons, meet the critical LOD value, and have minimal single locus mismatches. This should account for the presence of minor genotyping errors while minimizing the loss of parental assignment matches.

Relative reproductive success (RRS) will be calculated annually between classes of fish by standardizing to the pre-spawn maiden class of adults. Lifetime reproductive success (LRS) estimates for kelts will be calculated by adding the RRS of Event-1 and Event-2 estimates. This estimate of LRS does not look at individual fish that spawned across multiple years, nor does it look at the same group of fish across 2 consecutive years (e.g., Maiden in 2013, reconditioned kelts in 2014). Rather, it adds the RRS estimates of fish spawning in the same calendar year.

Milestone 1. Completed annual parentage analysis of juvenile samples.

Milestone 2. Submission of manuscript for reproductive success by juvenile assignments to a scientifically accredited journal by 2022.

Milestone 3. Submission of manuscript for adult-to-adult reproductive success to an appropriate journal by 2026.

Goal 3, Objective 3.3: Evaluate the homing and straying rates of reconditioned kelts in the Yakima River Basin. Monitor and analyze the detection history for verifiable first and second spawnings for all reconditioned kelts to estimate quantifiable homing and straying rates.

Sub-basin level homing of reconditioned kelt steelhead

In the Yakima River, kelt steelhead are reconditioned and released in the fall approximately four months prior to the next spawning season, but not all fish are appropriate for a repeat homing fidelity investigation since the released population in the past contained both mature (consecutive) and immature (skip) spawners. All long-term reconditioned kelts for this study will be released below Prosser Dam in the lower Yakima River with a PIT tag for individual identification. Final spawning destination of these individuals will be assigned through tributary, in-river, or dam PIT tag detection, recapture prior to spawning, or recollection as post spawn kelts the following emigration season. If the reconditioned kelts return to the same river system they were collected prior to or during the subsequent spawning season, then we will presume that they have rehomed to the same river basin they spawned as maidens. If they are detected in a sub-basin other than the one they were collected in, prior to, or during the subsequent spawning season, then we will presume that they strayed from the location they spawned in as maidens.

It will not be possible to determine the final destination for all fish released below Prosser Dam, therefore, we will exclude all fish that are not detected post release (likely lost tag or immediate mortalities), fish that are likely immature or skip spawners that are either lost to the Mainstem Columbia River hydrosystem during their emigration, or fish that emigrate successfully but do not successfully reascend to a sub-basin river, and have a subsequent spawning season (likely harvest or hydrosystem mortalities). Final PIT tag detections will be determined using the PTAGIS database. Once the steelhead spawning season concludes each year, the PTAGIS database will be queried to obtain the complete tag history for all PIT tag codes released from the kelt reconditioning program the previous fall ([MethodID: 4095](#)). Final detections will be assigned using the last uppermost detection location which includes in-river or dam PIT arrays, and/or recapture from sampling programs throughout the basin.

Sub-population level homing of reconditioned kelt steelhead

To investigate smaller scale sub-population level homing fidelity, we will limit our investigation to specific known sub-population maiden spawners and subsequent reconditioned kelt spawner homing events. Maiden spawner origin will be determined through detections of first-time spawners outfitted with PIT tags at tributary level or sub-population detection arrays, through parentage analysis of progeny collected in tributaries that connect confirmed maiden spawning locations to kelt steelhead parents in the reconditioning program, or kelt steelhead collected at tributary weirs.

Installation of in-stream PIT arrays provide data at both the individual and stock level. PIT arrays are installed in Satus Creek, Toppenish Creek, Ahtanum Creek, Taneum Creek, and Roza Dam in the Yakima River Basin, and in Omak Creek in the Okanogan River Basin. Wild origin steelhead are rarely PIT tagged, therefore the number of maiden spawners that will be previously PIT tagged and detected at the sub-population level during maiden spawning events is relatively low. To increase sample size at the maiden stage, we will use parentage results as a proxy for PIT tag detections of the maiden spawning event. Satus and Toppenish creeks have an ongoing reproductive success study which permits the assignment of Age-0 juveniles to kelt parents in the reconditioning program (see [Goal 3, Objective 3.1](#) for methods), thereby identifying the parents of these juveniles from the maiden spawning event. Likewise, previously PIT tagged wild origin steelhead in Omak Creek are rare, but all the wild steelhead kelts collected for the reconditioning program originated from the Omak Creek weir, therefore, confirming the origin of their first spawning.

All reconditioned kelts used in this portion of this study will be PIT tagged and released either below Prosser Dam in the Yakima River or in the mainstem Okanogan River approximately four months prior to the subsequent spawning season. Like the Sub-Basin level study, PIT tagged reconditioned kelts that could not be tracked for the reasons outlined above will be excluded from the analysis. If reconditioned kelt steelhead are detected at PIT tag arrays of their maiden tributary a second time in a subsequent spawning season, we will consider that successful homing. If reconditioned kelts return to a different tributary, then we will consider those fish to have strayed. Fish included in this study will not be included in the sub-basin level study.

Milestone 1: Complete and submit a manuscript describing homing of reconditioned kelt steelhead to an appropriate journal by 2022.

Goal 4. Carry out studies to achieve a sufficiently detailed understanding of the physiology of reconditioning in kelt steelhead to provide a scientific basis for maximizing the success of reconditioning programs.

Goal 4, Objective 4.1: Use blood sampling and plasma reproductive hormone assays to evaluate reproductive status of kelts in reconditioning programs.

Kelts will be sampled by either project leaders or the physiology team in September of each year. Non-lethal blood sampling of kelts will be conducted by caudal venipuncture according to published methods (Caldwell, et al. 2013; Houston 1990; Jenkins, et al. 2019; Pierce, et al. 2017). The volume of blood taken is a small fraction of the total, and the sampling procedure is

not particularly hard on the fish. Approximately 2-mL of blood will be drawn from anesthetized fish using heparinized syringes, centrifuged, and plasma stored at -80 °C prior to assay for estradiol (E2) in females or 11-ketotestosterone (11-KT) in males. E2 is the principal estrogen and 11-KT the principal androgen in fishes. An extraction procedure to remove interfering substances is required before steroid assays ([MethodID: 5319](#)). Standard clinical assays designed for human or mammalian plasma will not give accurate results without extraction. Extraction and assay procedures will be conducted at the CRITFC/University of Idaho physiology laboratory according to published methods (Caldwell, et al. 2014; Jenkins et al. 2019; Medeiros, et al. 2018)([MethodID: 371](#), [MethodID: 5320](#)). A histogram of log plasma E2 or 11-KT levels will be constructed and the cutoff between maturing and non-maturing individuals identified. This is typically about 1000 pg/ml E2, although it may vary depending on sampling time and the kelt population. Less information is available on males, but a cutoff of 2000-4000 pg/ml 11-KT is typical.

Goal 4, Objective 4.2: Use physiological tools in studies to improve survival and rematuration rates at kelt reconditioning facilities (with Objective 1.2).

Blood sampling, plasma hormone assays, and other plasma factor assays will be used to assess recovery from spawning, growth and metabolic status, and reproductive development in kelts during reconditioning. Blood sampling and reproductive steroid assays will be conducted as described above ([Goal 4, Objective 4.1](#)). Growth hormone (GH) and insulin-like growth factor-1 (IGF-1) assays will be conducted as described in a recent publication from our laboratory (Medeiros, et al. 2020). Plasma metabolite and osmolality assays will be conducted as previously described (Jenkins, et al. 2020; Jenkins et al. 2019).

Goal 4, Objective 4.3: Use physiological tools to evaluate fish characteristics at collection that correlate with survival and rematuration at kelt reconditioning facilities (with Objective 1.3).

Blood sampling of fish at collection for reconditioning and plasma assays will be conducted as described above ([Goal 4, Objective 4.2](#)). The survival and maturation status for each fish are tracked as part of the standard operation of reconditioning facilities. Multiple logistic regression analysis will be employed to determine predictors of survival and rematuration (e.g., Jenkins et al. 2020).

Goal 4, Objective 4: Evaluate physiological indices at release as predictors of post-release migration and spawning success in kelts released from reconditioning programs.

Blood sampling of fish during the fall survey for maturation status or at release and plasma assays will be conducted as described above ([Goal 4, Objective 4.2](#)). Migration and spawning success will be assessed by post-release PIT tag detection histories as described above ([Goal 3, Objective 3.2](#)). Multiple logistic regression analysis will be employed to evaluate predictors of migration and spawning success.

Goal 5. Project wrap up, delineation of necessary follow up monitoring and research, complete manuscripts, transition to other locations or uses of kelt reconditioning technology.

Goal 5, Objective 5.1: Develop plan for necessary monitoring and research to implement after kelt reconditioning production facilities at Prosser and Nez Perce Tribal hatcheries are completed and on line.

This project has operated at a research-scale to prove the concept of kelt reconditioning. That phase will end once dedicated kelt reconditioning facilities are constructed and on line. We will develop a plan for monitoring and evaluating the production-scale facilities. In addition, we will identify any research topics that require further study (I.e., skip spawner management strategies).

Goal 5, Objective 5.2: Complete and submit manuscripts and technical reports related to kelt physiology.

We have several studies and experiments that need to be fully analyzed, written up, and published. This work is ongoing and will extend into 2022.

Milestone 1: Complete and submit manuscript describing kelt fasting experiment conducted using DNFH hatchery kelt model to a scientifically accredited journal by 2021-2022.

Milestone 2: Complete statistical analysis and writeup of study on post-spawning ration restriction using post-spawning rainbow trout model by 2021-2022, followed by submission to a scientifically accredited journal.

Milestone 3: Complete data processing, analysis and writeup of study on predictors of post-release migration and spawning success in female kelts release from Prosser Hatchery by 2022-2023. Submission to a scientific journal or publication as a technical report to depend on results.

Goal 5, Objective 5.3: Survey the region for potential to transfer kelt reconditioning technology to other locations and applications such as hatchery broodstock management or reintroductions.

We plan to survey the region for potential applications of kelt reconditioning in other sub basins and applications of kelt reconditioning. Potential applications could be for hatchery broodstock management using air spawning techniques and then reconditioning those fish for use in future spawning years as a hedge for years with inadequate broodstock. Wild reconditioned kelts could also be used as stock for reintroductions without impacting other fish sources. This work is planned to occur from 2022-2026.

Chapter 1: Kelt Capture, enumeration, and reconditioning in the Yakima and Snake basins.

Introduction

Kelt steelhead reconditioning process evaluations involve fish culturing practices, studying alternative management strategies, and implementing research scale reconditioning programs. Adding repeat spawner steelhead to the population through reconditioning can add stability through the portfolio effect (Moore et al. 2014) and increase population abundance by increasing lifetime reproductive success (Seamons and Quinn 2010). We established “control” groups in both the Snake and Yakima rivers. These control groups were downstream migrating kelts, systematically collected, PIT tagged and released back into the river each year. These fish are monitored through PITAGIS to determine how successfully they naturally recondition in the ocean.

We define long-term reconditioning as holding and feeding post-spawn steelhead in a captive environment to increase kelt survival and create additional spawning opportunities. The long-term steelhead reconditioning diet and care treatments were established from the studies conducted in 2001 and 2002 (Hatch et al. 2002 and Hatch et al. 2003b) and summarized in Hatch et al. 2013b. These fish are typically released in the fall to over-winter and return to the spawning sites volitionally. This chapter recaps our 2021 kelt collection efforts; for a broader review of specific fish culturing practices please see (Hatch et al. 2015).

Methods

Standard Data Collection

All captured steelhead are scanned for existing PIT-tags, and biological data is collected which includes determination of kelt/maiden status, fork length, weight, condition factor (color and presence/absence of wounds/skin-body condition), coloration rating (bright, medium, dark), notation of clipped or non-clipped fins (typically adipose), and small (typically a 1 x 1 mm) tissue sample (caudal fin clip) is collected for genetic analysis. Steelhead without a PIT-tag receive a 12.5 mm PIT-tag injected into the pelvic girdle to track migration history and to determine reconditioning efficacy. All releases or mortalities are recorded including date of event, condition factor, and PIT-tag identifier. In the case of a lost PIT-tag, typically at time of release, fish are retagged, and an additional genetic sample is collected. The data is then uploaded to a central kelt database at CRITFC and any fish that is released back to the wild is entered into the PITAGIS database.

Steelhead Kelt Collection

Chandler Juvenile Monitoring Facility

Once diverted into the CJMF ([Table 1, site 1](#)), emigrating kelts are manually collected from a fish separation device (a device that allows smaller juvenile salmonids to “fall through” for processing in the juvenile facility while larger fish can be dip netted for processing and transferring to the reconditioning tanks at Prosser Hatchery (Table 1, site 2). Yakama Nation staff monitored the Chandler bypass separator during the kelt migration.

Lower Granite Dam

Steelhead kelts entering the juvenile bypass separator ([Table 1, site 3](#)) are collected by Army Corps of Engineer (COE) staff. Kelts are netted off the adult fish separator bars and moved to a fish hopper that leads into the kelt receiving tank. Staff from the Nez Perce Tribe (NPT), processed fish that were diverted into the receiving tank. Kelt steelhead, judged to be in good or better condition, with intact adipose fins, and >63cm are then trucked to DNFH for reconditioning. The transport truck had a 1.5-kiloliter tank fitted with supplemental, regulated, and compressed oxygen that was fed via air stones; also, a 12-volt powered tank aeration pump was used to circulate oxygenated water. Stress Coat® or PolyAqua® was used to replace the natural protective slime coating that may have been compromised by handling. In addition, salt was added to reduce osmo-regulatory stress. Temperature and dissolved oxygen levels were also monitored during transport. Loading densities were kept to a minimum so that no more than 20 kelts were transported at one time.

Little Goose Dam

Collection and protocols followed are the same as Lower Granite Dam ([Table 1, Site 31](#)).

Long-term Reconditioning

Long-term reconditioning is a management strategy where emigrating kelt steelhead are collected and held in large tanks, given prophylactic treatments and fed a specially formulated diet for approximately 6 months (Hatch et al. 2013b). After 6 months, the “reconditioned” kelts are released back into the collection river as the run at large is returning from the ocean. These reconditioned fish generally mingle with the run at large and proceed to in-river, over-winter locations and spawning grounds in the spring. This strategy seeks to reduce mortality in the hydro system and ocean, providing another opportunity for fish to reproduce in the wild. Techniques used in kelt reconditioning were initially developed for Atlantic salmon *Salmo salar* and Brown or Sea-trout *S. trutta*, and a review of these studies and others applicable to steelhead kelts are summarized in Evans et al. (2001). All kelt reconditioning conducted at Prosser Hatchery is primarily done in 4, 6.1m x 1.5m (20' x 5') circular tanks, 4 -smaller, 3m x 1.2m (10' x 4'), and one, 5.2m x 1.5m (17' x 5') circular tank also available. The water is spring fed, at 10-11°C. Fish are held at DNFH in one of 4, 4.6m diameter outdoor tanks, supplied with North Fork Clearwater River water at 200 liter/minute, maintained at a water height of 1.5m, with a seasonally varying temperature profile of 4.9 – 11.0°C.

Life History Strategies: Consecutive vs. Skip Spawning

The steelhead reproductive physiology research conducted by the CRITFC has determined that both natural and artificially reconditioned kelts can pursue two alternative pathways toward rematuration and repeat spawning. One pathway is termed consecutive spawning, where individuals remature and proceed to spawn in the next spawn cycle. The other pathway is termed skip spawning, where individuals remature and proceed to spawn two years after their previous spawning. To illustrate, kelts collected in the spring of 2021 could spawn again in the winter/spring of 2022 as consecutive spawners, or wait until the spring of 2023, and instead spawn as skip spawners. The proportion of consecutive and skip spawners in a cohort varies annually and is detailed in [Chapter 2](#), but in general, Yakima River fish predominately follow the consecutive spawner pathway (60-70%) and will spawn the following winter/spring, while the majority of Snake River kelts follow the skip spawner life history (60-70%), which has them held for an additional 12 months after capture, with subsequent spawning the next winter/spring. Fish strategy choice is likely controlled by a combination of genetics and environment.

Additional study is needed to evaluate reconditioning strategies for skip spawners. Our past approach was to hold the fish for an additional year in the hatchery and then release them. This scenario works well in the Snake River but is less successful in the Yakima River likely due to difficulties keeping intake water chilled adequately during the winter months. This is important, as the temperature regime of the spring water utilized at Prosser can be approximately 5-8 degrees warmer than ocean/river water during the winter months. This warmer water appears to cause maturation synchronization issues due to the prolonged holding that is needed for skip spawners. Additionally, the cost of holding fish for an additional year should be compared to results from other scenarios. We plan to continue with experimental releases at least through 2022 until we can determine if it is a worthwhile strategy to convert immature spawners, by using a combination of artificial and natural ocean reconditioning to propagate viable mature spawners. If results are promising, this may be a cost-effective method to increase kelt spawner abundance in the basin.

Summary Research-Scale Efforts to Address RPA 33 and subsequent Biological Opinion (2019)

At DNFH in 2021, we continued to conduct research toward addressing Reasonable and Prudent Alternative 33 for the Hydro-system Biological Opinion. The RPA 33 required the Action Agencies to develop, in cooperation with regional salmon managers, implementation of a Snake River steelhead kelt management plan, designed to provide at least a 6% improvement in B-run population productivity (NMFS 2008, 2010, and 2014). Toward that goal, a variety of approaches were tested and implemented including passage improvements and reconditioning kelt steelhead. The designation of B-run can be difficult to quantify in the Snake Basin. Historically, size has been used to determine B-run fish populations, which has been determined by genetic analysis to not be the sole determinant of B-run populations, since genetic assignments have provided data that B-run populations do not meet specific size requirements or overlap with A-run populations. To accurately measure how kelts contributed towards the 6% RPA 33 rule NOAA, CRITFC, Nez Perce Tribe, and the Federal Action Agencies (Bonneville Power Administration, U.S. Army Corps of Engineers, and the Bureau of Reclamation) devised a system that would recognize kelt reconditioning actions and give credit

for those fish we successfully reconditioned and released towards the 6% RPA value ([Kelt Master Plan Document](#)) .

In the spring of 2019, the NOAA published a new Hydrological Biological Opinion for the Columbia River Power System ([NMFS 2019](#)). The new rules effectively retired the 6% number in 2019 and considered all future Snake River reconditioning actions as the only hatchery actions appropriate for corrective mitigation for steelhead loss in the Snake River. We have continued to maintain this evaluation to gauge how much progress we are making reconditioning kelts in the Snake River Basin.

Results/Discussion

Steelhead Kelt Collections

Large numbers of kelt steelhead are available for collection at many sites across the Columbia River Basin. These sites generally are associated with juvenile bypass systems or weirs. From 2002-2021 a total of 23,036 downstream migrating kelts at LGD were collected. Starting in 2020, a total of 97 kelts have been collected from Little Goose Dam, and 15,384 from the CJMF, from 2000-2020, which were collected and then either released back to the river or retained for reconditioning. The Columbia River, upriver steelhead run, in 2021-22, was the worst year recorded since unclipped fish were counted ([FPC Data](#)). We collected 250, 14, and 365 kelts at LGD, LGS, and CJMF, respectively ([Table 1.1](#) and [Appendix A](#)). In 2021, the kelt collection represented 2.2 % and 23.6% of the upstream run in the Snake and Yakima rivers, respectively.

Reconditioning

Since 2011, 1,877 kelt steelhead have been retained for reconditioning from collections in the Snake River and 731 fish have survived to their first fall. Since 2000; 11,218 kelt steelhead were retained for reconditioning from collections at CJMF and 5,006 fish survived to the first fall of the annual collection period ([Appendix A](#)). Snake River collections were made at the LDG and starting in 2020 also at LGS, however it should be noted that kelt collections have also come from both the South Fork Clearwater River and Fish Creek (Lochsa River tributary) in previous years (Hatch et al. 2018).

Table 1. 1. Summary of kelt collections and fish disposition in 2021 from the Snake and Yakima Rivers.

	Lower Granite	Little Goose	Snake River total	Yakima River total
Total Collection	304	26	330	365
# Released back to river	250	14	264	5
# Taken to reconditioning facility	54	12	66	360
# Mature reconditioned fish released (consecutive spawners)	4	0	4	149
# Immature reconditioned fish retained (skip spawners)	17	1	18	NA
# Immature reconditioned fish released (skip spawners)	NA	NA	NA	53
# Skip spawners released from previous collection	58	3	61	NA

Long-term reconditioning survival (from collection to the release in the fall) is variable from year to year but has averaged 45% at the Prosser Fish Hatchery (PFH) over the last 22 years. For the last 5 years, Yakima reconditioning has been just over 57.8% survival (Figure 1.1). The staff here has several years of reconditioning experience, so we generally observe only small annual variations in survival (Figure 1.1). The reconditioning survival rate for wild Snake River kelts from 2011 through 2021 is 39.1%. Survival during the initial years at DNFH (2011-12) was compromised as a result of poor water quality which has been detailed in previous reports (Hatch et al. 2012 and Hatch et al. 2013a) and compounded by the inexperience of new hatchery staff. However, the past 5 years have seen marked improvement in survival rates, as water quality issues have been improved and staff have gained additional experience in handling and caring for wild steelhead. The survival rate for the last five-year period combined at Dworshak National Fish Hatchery and Nez Perce Tribal Fish Hatchery have averaged 47.4% (Figure 1.2). In both 2020 and 2021 the kelt collections were low (137 & 66 respectively), but long-term survival has been good (43%). Prevalence of the skip spawner life history continues to be much higher in the Snake River than in the Yakima River. On average, skip spawners compose 60% of the kelt collection in the Snake River, and approximately 40% in the Yakima River. This difference is likely the result of energy demands of longer migration distance in the Snake River (Keefer et al. 2008). Figures (1.1) and (1.2) show differences in the life history strategy by the proportion of retained fish (skip spawners) at the two facilities. Reconditioned steelhead kelt releases for the Yakima River have been at historic lows due to poor river migration conditions (high temperatures and low flows), low ocean adult returns, and high water/flooding during collection periods. The improvement in Snake River releases have trended upwards in large part to successful retention and reconditioning of skip spawners

(Figures 1.1 and 1.2) at the DNFH due to improved water quality, implementation of rearing protocols, improved life support systems, improved staff training, and experience.




Figure 1. 1. Yakima River steelhead kelt collection for reconditioning and fate from 2013-2021. Beginning in 2019, kelts that would have been retained, were instead trucked, and released below Bonneville Dam. Will be updated in 2022 Annual Report.




Figure 1. 2. Snake River steelhead kelt collection for reconditioning and fate from 2013-2021. Will be updated in 2002 Annual Report.

Skip Spawner Management Strategy in the Yakima River

Reconditioned kelt steelhead demonstrate either consecutive- or skip-spawning life histories. Skip spawners make up approximately 40% of collection in the Yakima River and 60% of the collection in the Snake River. Our primary management strategy for skip spawners was to hold the fish in the reconditioning for an additional year, thus releasing the fish in the second fall of captivity. This strategy has been successful in the Snake River where second year survival and rematuration of skip spawners has been very high (60.5% survival and 26.1% rematuration). In the Yakima River, this management strategy of holding skip spawners for a second year has been much less successful in terms of skip spawner survival. This likely is a result of warmer water temperature that is used to rear skip spawners during the winter. The water source is from a well that is ideal for summertime rearing, but that same water is much warmer than the average ambient river/ocean water temperatures where kelts would naturally recondition. This water temperature fluctuation likely had an impact on held skip spawner maturation timing.

Beginning in 2019, we experimented with transporting skip spawner kelts to the lower Columbia River and releasing them in the fall, concurrent with consecutive spawner releases in the Yakima River.

2019 Yakima River Lower Columbia River Release

We trucked and released 103 PIT-tagged, immature, reconditioned kelts below Bonneville Dam on 10/31/2019 to evaluate this management strategy. Thirty fish were detected moving upstream at the Bonneville Dam fish ladders within the first 30 days post release. Ten of these 30 fish had additional detections, four moving downstream, and six moving upstream. The four downstream moving fish were detected either in the Bonneville Dam juvenile bypass or corner collector. These migration patterns were consistent with post-spawn steelhead outmigration return to the ocean.

Final detection history for the six upstream moving fish includes three fish last heard at mainstem dams (2 at The Dalles and 1 at John Day ladders), and three fish last detected in the Yakima River tributaries. Tributary detections included one fish in Satus Creek. This fish genotyped to the Yakima River GSI reporting group and was likely a male based on genetic analysis. Another fish was last detected in Little Sheep Creek, a tributary of the Imnaha River in the Snake River basin. This fish genotyped to the MGILCS (mid-Columbia, Grande Ronde, Imnaha, lower Snake, lower Salmon, lower Clearwater) reporting group and also was likely a male based on genetic analysis (Hess et al. 2020). Finally, one fish was detected in Fifteen Mile Creek, located just upriver of the Dalles Dam, a small tributary off the mainstem Columbia River. This fish genotyped to the MGILCS reporting group and was likely male, based on genetic analysis.

We will continue to monitor PTAGIS possible detections from these lower river release groups. Skip spawning adult returns from the 2019 release group would be expected back in the summer/fall of 2021. So far, no return PTAGIS detections have been recorded for this release in 2021.

2020 Yakima River Lower Columbia River Release

On October 21, 2020, we released 52 PIT-tagged, immature, reconditioned kelts below Bonneville Dam. Due to COVID-19 concerns, blood draws were not conducted at Prosser to determine maturation status. Instead, visual maturation was determined by the hatchery manager. We initially observed 8 of these kelts moving upstream at Bonneville Dam. One fish has migrated back to Prosser already, which was a female, based on earlier visual calls when selected for reconditioning. Likely, this fish maturation status was called incorrectly and will be a consecutive spawner. We anticipate a that more thorough analysis will be ready in 2021, which will include genetic analysis to accurately determine sex and GSI origin.

We continue to monitor PTAGIS for any possible return detections which we would anticipate beginning as early as July of 2021, possibly going into 2022. At the date of this report, we have had no detections of these fish returning from the ocean.

2021 Yakima River Lower Columbia River Release

On October 28, 2021, we released 53 PIT-tagged, immature, reconditioned kelts below Bonneville Dam. Blood draws were resumed in 2022, and these fish were determined to be immature based on the blood hormone profiles established in the lab.

Long-term reconditioning mature kelt release

We evaluated the traits and survival-to-release of reconditioned kelt steelhead *Oncorhynchus mykiss* in the Yakima River (Hatch et al. 2013b). Reconditioned steelhead kelts continue to be predominantly (>92%) female. The annual survival to release average ranged from 18% at the start of the program in the early 2000's to an annual high of 76% in 2016 and averaged 45% over the course of the study (2000-21) with surviving reconditioned kelts showing increases in fork length, weight, and Fulton's K condition factor. Kelts in good condition, and those with bright coloration at the time of collection were more likely to survive. Post-release upstream migration timing of reconditioned kelts was spread out over several months and correlated well with run timing of upstream pre-spawn migrants. The empirical results we observed demonstrate the potential of kelt reconditioning to provide recovery benefits for ESA-listed, repeat spawning steelhead populations in highly developed river systems. See [Appendix A](#) for annual data.

In Figures 1.3 and 1.4, the numbers of female kelts reconditioned and released are added to the corresponding steelhead run. For example, a consecutive kelt that is collected in the Spring of 2014 (from 2013/2014 run) would be released into the following run year 2014/2015. For a skip spawner, which is typical of Snake River kelt, a fish caught in the Spring of 2014 would be released into the 2015/2016 run year. Figures 1.3 and 1.4 demonstrate that the contribution of reconditioned kelts to the overall runs in the Yakima and Snake rivers is measurable and quite

substantial in comparison to no intervention. While the total number of returned fish may appear small compared to the overall run, especially in the Snake, many of these fish are contributing towards ESA-listed populations throughout the Yakima and Snake basins. In Figure 1.4, both Snake River A and B-run populations are counted. We primarily targeted B-run fish while selecting kelts for reconditioning although there were a small proportion of A-run fish that were also collected and reconditioned. The A-run population in the Snake is much larger than the B-run population, with most of these fish coming from the Grande Ronde basin (Hatch et al. 2019).




Figure 1. 3. Contribution to steelhead run from reconditioned kelt release in Yakima Basin. * Run is still ongoing; value will be updated in 2022 steelhead kelt reconditioning annual report.




Figure 1. 4. Contribution to steelhead run from reconditioned kelt release in Snake Basin at LGD. * Includes Fish Creek and/or South Fork Clearwater Kelts.

The natural repeat spawner rate (or no intervention) measured as a return rate to Bonneville Dam, for the Snake River, is 0.27% and 2.90% for the Yakima River. The calculated benefit of reconditioning relative to leaving the fish in the river is 171.5 times for the Snake River and 15.2 for the Yakima River.

Summary Research-Scale Efforts to Address RPA 33 and subsequent Biological Opinion (2019)

Since operating at a research scale in the Snake Basin, as approved by the ISRP in the 2008 review, the capacity of our facility was much too small to meet the RPA 33 goal of increasing the LGR ladder count of B-run steelhead by 6%. However, we have demonstrated the feasibility of reaching the 6% goal. Releases of successfully reconditioned mature kelts began in 2011, but due to water quality issues in both 2011 and 2012 (Hatch et al. 2012 and 2013) these numbers were severely under representative of what we could accomplish at the initial site (Dworshak National Fish Hatchery) and with the inclusion of an additional temporary reconditioning site in 2016 (Nez Perce Tribal Hatchery). In 2020 and 2021, we only reared fish at DNFH due to low collection numbers. Table (1.1) summarizes all collections for both A and B-run, and releases associated with the RPA 33.

In 2013, we had a successful reconditioning and release at nearly 40% towards reaching the RPA 33 goal with 69 female fish released (Table 1.2). Our best year was in 2017, with releases just over 50% towards the RPA goal, at 98 mature female spawners released. In 2019, we had a lower number of mature kelts to release due to a mishap at the Nez Perce Tribal Hatchery, which resulted in a significant loss of retained skip spawner kelts from 2018. Nez Perce Tribe is taking actions to prevent such a catastrophe in the future, with protocols that should help to eliminate the chance that such an event will happen again. Since 2013, we have averaged 32% of the RPA goal, releasing an average of 56 mature female fish per year with a total of 525 mature female fish released from 2011-2021 (Table 1.2). Figure 1.5 is the schema that was devised by the Power Agencies and CRITFC to determine the credit that would be allocated towards collection and successful reconditioning of what was considered B-run kelt steelhead towards the larger B-run the year of release. The number represents the B-run steelhead released x estimated successful reproductive contribution. This is not representative of number of total kelts released and was generated under the direction of RPA 33. We will continue to estimate B-run steelhead kelt contributions in future reports for the benefit of management implications of kelt reconditioning in the Snake River Basin.

Table 1. 2. Summary of fish collections and releases in the Snake River associated with RPA 33.

Year	Collection Location	Number of Fish Collected	Number of Fish that Survived Reconditioning	% Survival	Consecutive Spawner Release	Number of Fish Retained	Mature Skip Spawners Released (Capture Year)	Total Release by Year
2011	Lower Granite Dam	111	2	1.8%	2	-	-	-
2011	S.F. Clearwater	-	-	-	-	-	-	-
2011	Fish Creek	-	-	-	-	-	-	-
2011 (subtotal)		111	2	1.8%	2	-	-	2
2012	Lower Granite Dam	124	10	8.1%	10	-	-	-
2012	S.F. Clearwater	-	-	-	-	-	-	-
2012	Fish Creek	-	-	-	-	-	-	-
								-
2012 (subtotal)		124	10	8.1%	10	-	-	10
2013	Lower Granite Dam	110	57	51.8%	57	-	-	-
2013	S.F. Clearwater	24	12	50.0%	12	-	-	-
2013	Fish Creek	-	-	-	-	-	-	-
2013 (subtotal)		134	69	51.5%	69	-	-	69
2014	Lower Granite Dam	110	34	30.9%	34	-	-	-
2014	S.F. Clearwater	-	-	-	-	-	-	-
2014	Fish Creek	12	3	25.0%	1	2	2	-

2014 (subtotal)		122	37	30.3%	35	2	2	35
2015	Lower Granite Dam	22	11	50.0%	8	3	3	8
2015	S.F. Clearwater	35	7	20.0%	4	3	0	4
2015	Fish Creek	83	25	30.1%	10	15	15	12*
2015 (subtotal)		140	43	30.7%	22	21	18	24
2016 (subtotal)	Lower Granite Dam	227	120	52.9%	19	101	77	37*^
2017 Subtotal	Lower Granite Dam	269	59	21.9%	21	58	29	98^
2018 Subtotal	Lower Granite Dam	259	177	68.3%	50	99	1	79^
2019 Subtotal	Lower Granite Dam	288	121	42.0%	39	85	58	40^
2020	Lower Granite Dam	88	53	60.2%	7	46	58	65^
2020	Little Goose Dam	49	18	36.8%	1	17	3	1
2020 Subtotal		137	71	51.8%	8	63	61	66^
2021	Lower Granite Dam	54	21	47.7%	4	17	TBD 2022	4
2021	Little Goose Dam	12	1	8.3%	0	1	TBD 2022	0

2021 Subtotal		66	22	33.3%	4	18	0	65^
Grand Total		1877	731	38.9%	279	447	246	525
				*Includes Fish Cr. kelt skip spawners.				
				^Includes previous year kelt spawners from LGD and LGO.				




Figure 1. 5: Contribution to steelhead run from reconditioned kelt release in Snake River Basin. x2021-2022 value is an estimated run size and will be corrected in the 2021 Kelt Annual Report. B-run numbers utilize [Kelt Master Plan](#) definition for reconditioned female kelt spawners.

Chapter 2. Cumulative Results

This section was part of our response to the Northwest Power and Conservation Council's recent Anadromous Habitat and Hatchery Projects Project Review ([ISRP 21-8](#)). It's included to provide a comprehensive presentation of project results over the last decade.

Steelhead kelt reconditioning programs collect steelhead kelts (post-spawned adult steelhead) to recondition them so that they can be released into the river to spawn again. Steelhead kelt reconditioning is the practice of capturing, holding, and feeding post-spawned steelhead in an artificial rearing environment for the purpose of regeneration of gonads for repeat spawning (circumventing the migration down the river, ocean rearing, and migration back to tributaries). Reconditioned kelts are released back into the river after 6 or 18 months depending on the rematuration schedule.

Starting in 2000, CRITFC and the YN collaborated to investigate steelhead kelt reconditioning. At the time, little was known or published about kelt steelhead reconditioning. Consequentially, initial work focused on very basic components: collecting kelts, how to do basic fish care, feed types and rations, treatments, evaluating maturation status, release strategies, and other basic components. We used literature published on Atlantic salmon kelt reconditioning as a starting point, but a lot was developed from trial, error, and adaptive management. Early reviews from the ISRP helped define areas for research and investigation, including evaluating the relative reproductive success of reconditioned kelt steelhead, as well as developing an understanding of kelt physiology. The Columbia Basin Fish Accords were signed in 2007 and at the same time we were seeing encouraging results from the Yakima River Kelt Project, so we expanded to include a Snake River component and a collaboration with the Nez Perce Tribe. The ISRP posed five questions to a sister Yakama Nation project ([2008-458-00](#)) in a 2014 review:

1. "The prior recommendation, by the ISRP, to establish methods to assess how kelt reconditioning may benefit population growth, abundance, spatial structure, and diversity still needs to be addressed.
2. Some modeling and a power analysis need to be conducted to clarify how many juveniles and F1 adults should be sampled to detect meaningful differences in the breeding and reproductive success of hatchery-origin, natural-origin, and reconditioned natural-origin females.
3. Methods to assess the fat levels, maturation timing, fecundity, egg size, and gamete viability of the project's reconditioned kelts need to be developed and implemented. The fate of non-maturing or skip-spawning reconditioned fish should be documented.
4. Viable plans are needed to monitor the homing and straying rates of reconditioned kelts released by the project.
5. Experiments are needed to discover the best geographic locations and times of year for release of the project's reconditioned fish."

We have used these questions as guidance for our studies since 2014 and we will organize our response on progress to date around these themes. This project has published 17 papers on these subjects since 2013.

What are the benefits to the receiving population in terms of Viable Salmonid Population (VSP) parameters?

Our project includes both Yakima and Snake rivers components and information will be provided on each of these systems. Primary focus will be on the Yakima River to build the case of benefits to population growth, abundance, spatial structure, and diversity due to differences in scale between the two systems.

From 2001 to 2021, a total of 11,218 out-migrating kelts were collected at the Chandler Juvenile Collection Facility (Table 2.1). All captured fish were PIT tagged for individual identification during reconditioning and following release. Kelts were held at Prosser Hatchery for reconditioning. The average survival rate of reconditioned kelts to release was 49%. On average, 509 kelts were collected and 234 were released. The majority (>92%) of kelts were female (Hatch et al. 2019). We evaluated the traits and survival to release of reconditioned kelt steelhead in the Yakima River program from 2001 to 2011 in an accredited peer reviewed publication (Hatch et al. 2013b).

Table 2. 1. Yakima Basin steelhead kelts reconditioned at Prosser Hatchery, number of kelts released, and survival rate to release, 2001 to 2021. Source: Yakama Nation.

Year	Number of Kelts Collected	Number of Reconditioned Kelts Released	Survival to Release
2001	508	108	21.3%
2002	420	142	33.8%
2003	482	301	62.4%
2004	694	288	41.5%
2005	427	86	20.1%
2006	279	85	30.5%
2007	422	221	52.4%
2008	472	266	56.4%
2009	510	141	27.6%
2010	1,100	426	38.7%
2011	680	223	32.8%
2012	572	333	58.2%
2013	575	308	53.6%
2014	481	316	74.4%
2015	1,098	435	43.4%
2016	525	341	73.0%
2017	117	84	71.8%
2018	227	152	67.0%
2019	371	248	66.9%
2020	463	320	69.1%
2021	360	219	60.8%
Total	11,218	5,006	44.6% (pooled)
Average	509	234	47.6%

Reconditioned kelts were released adjacently upstream of Prosser Dam from 2001 to 2007, and from 2008 to present, approximately one kilometer downstream of Prosser Dam.

Reconditioned kelts are released into the river from mid-October to early December (when water temperature of rearing water and the river were in conjunction) when the majority of the natural run is returning from the ocean. Reconditioned fish mingle with the natural run, then proceed to overwintering locations to hold, and then proceed on to spawning grounds in the spring to create redds and mate. The release location was selected to allow reconditioned kelts to migrate upstream and choose their spawning location volitionally. Spawning in the Yakima River basin generally peaks from early March to early May depending on the elevation of the stream (goes from Satus, Toppenish, Ahtanum, Naches, and upper Yakima). The timing of upstream migration by reconditioned kelts from 2008-present was determined based on PIT tag detections at Prosser Dam.

We compared return rates of repeat spawners among long-term reconditioning, short-term reconditioning, immediate transport to the lower Columbia River, and control groups in the

Yakima River. Long-term reconditioning demonstrated significantly higher return rates of repeat spawners (11-18%) than other treatments (1-3%) (Table 2.2). This result was supported despite the typical variation in river, ocean, and fish conditions among the years that were incorporated into the error term in our analysis. The data extrapolation required in our analysis does not account for variation in environmental or fish conditions between years. However, this method does provide a best and worst-case interpolation of data for earlier years in the long-term reconditioned group, thereby strengthening our ability to draw conclusions among the four treatments. For more in-depth analysis see our publication (Trammell et al. 2016).

Table 2. 2. Sample size (N), mean, and grouping output for Tukey post-hoc test from ANOVA of PIT tag detections at Prosser Dam. From Trammell et al. 2016.

Treatment	N	Mean	Grouping
Long-term min	10	11.5	A
Long-term max	10	17.6	A
Short-term	7	3.2	B
Transport	7	0.9	B
Control	7	2.7	B

Table (2.3) gives the count of upstream migrating ocean-returning steelhead, reconditioned kelts released that were subsequently detected migrating upstream, and the combined total steelhead run year count at Prosser Dam for the most recent 14 return years. The percentage of the total return that consisted of actively migrating reconditioned kelts averaged just over 5.0% over these 14 years. Since we know reconditioned kelts are predominantly female, these data demonstrate a substantial and measurable contribution to overall population demographics in comparison to no intervention. Based on average survival to release (~50%; Table 2.1) and release to spawning ground migration (~50%; Table 2.3), approximately 25% of kelts collected for reconditioning are expected to migrate to the spawning grounds.

Table 2. 3. Contribution to steelhead spawning run from reconditioned kelts released in the Yakima basin.

Run Year	Ocean Return	Reconditioned Kelts¹	Total	Reconditioned Kelts
2008-09	3,378	91	3,469	2.6%
2009-10	6,750	46	6,796	0.7%
2010-11	6,043	153	6,196	2.5%
2011-12	6,301	58	6,359	0.9%
2012-13	4,556	231	4,787	4.8%
2013-14	4,039	104	4,143	2.5%
2014-15	5,096	116	5,212	2.2%
2015-16	3,737	216	3,953	5.5%
2016-17	1,462	155	1,617	9.6%
2017-18	1,335	53	1,388	3.8%
2018-19	998	95	1,093	8.7%
2019-20	1,542	115	1,657	6.9%
2020-21	1,385	143	1,528	9.4%

2021-22	657	97	657	12.9%
---------	-----	----	-----	-------

¹ Kelts detected migrating upstream (natural and hatchery-origin).

Relative to other populations, adult productivity indices for Yakima River MPG steelhead are presently trending upward and Figure (2.1) indicates that Yakima River MPG steelhead are experiencing higher survival relative to other steelhead populations above Bonneville Dam. This could be a positive result of the kelt reconditioning program, though separating effects of habitat improvement projects and kelt reconditioning is difficult to determine.




Figure 2. 1. Yakima River MPG steelhead (Prosser wild abundance) as a percentage of Bonneville Dam wild Group A steelhead abundance, 1983 to 2018.

Impacts to spatial structure can be seen in data from the Snake River. We collect kelt steelhead at Lower Granite and (in some years) Little Goose dams, so the collection is a mixture of nearly all Snake River populations. As abundance of wild fish has decreased, our collection criteria for fish length has decreased as well, so as to spread benefits to all populations, not just to the B-run component. All reconditioned kelts are PIT tagged and released downstream of Lower Granite Dam in the fall. Figure 2.2 and Table 2.4 displays the point of last PIT detection for reconditioned kelts released in 2017, 2018, and 2019. Below the figure is a key to the PIT array locations and the number of different fish with final detections at each site. Though this is a coarse look at distribution, it does show that reconditioned fish are traveling throughout the Snake River Basin to find spawning grounds.




Figure 2. 2. Map of the Snake River basin with yellow pins indicating the last PIT detection location for reconditioned steelhead released in 2017, 2018, and 2019. Below is the key to PIT sites and the tally of fish with final detection at each site.

Table 2. 4. Corresponding sites by year, detection, and detailed location information.

SITE	2017	2018	2019	SITE DETAILS	
ACB		1		ASOTIN CREEK @ CLOVERLAND BRDG	SNAKE
AFC		1		NO/SO FORK ASOTIN CR	SNAKE
BSC	2	1	1	BIG SHEEP CR @ KM 6	IMNAHA
CCW		2	1	CATHERINE CREEK LADDER/WEIR	GRANDE RONDE
GRA	17	46	5	LOWER GRANITE DAM	SNAKE
IR1		2		LOWER IMNAHA RIVER @ KM 7	IMNAHA
IR2		4		LOWER IMNAHA RIVER @ KM 10	IMNAHA
IR3	2	5	3	UPPER IMNAHA @ KM 41	IMNAHA
IR5		2		IMNAHA WEIR UPSTREAM	IMNAHA
JOC	2	6	3	JOSEPH CREEK @ KM 3	GRANDE RONDE
KHS		1		BIG BEAR CREEK @ KENDRICK HS	POTLATCH
KRS			1	SF SALMON RIVER @ KRASSEL CR	SALMON
LAP	1			LAPWAI CREEK NEAR MOUTH	CLEARWATER
LC1	1	1	1	LOWER LOLO CREEK @ KM 21	CLEARWATER
LC2	2			UPPER LOLO CREEK @ KM 25	CLEARWATER

LOOKING		1		LOOKINGGLASS HATCHERY (PASSIVE)	GRANDE RONDE
LRL			1	LOWER LOCHSA RIVER	LOCHSA
LRU	5	2	1	LOCHSA RIVER UPPER SITE	LOCHSA
MR1			3	MINAM RIVER @ KM 0.5	WALLOWA
NFS		2		NF SALMON RIVER @ KM 0.5	SALMON
PCA		1		PANTHER CREEK	SALMON
SALR1			1	SALMON RIVER MOUTH TO KM 171	SALMON
SC1		1		LOWER SF CLEARWATER @ KM 1	CLEARWATER
SC2		1	1	LOWER SF CLEARWATER @ KM 2	CLEARWATER
SWT		1		SWEETWATER CREEK NEAR MOUTH	CLEARWATER
TAY		1		BIG CREEK @ TAYLOR RANCH	SALMON
UGR		2		UPPER GRANDE RONDE @ KM 155	GRANDE RONDE
UGS	1	1		UPPER GRANDE RONDE STARKEY	GRANDE RONDE
USE		1	1	UPPER SALMON RIVER @ KM 437	SALMON
UTR	1	1		UPPER TUCANNON RIVER @ KM 53.2	TUCANNON
VC1	1			VALLEY CREEK UPSTREAM SITE	SALMON
VC2	1			VALLEY CREEK DOWNSTREAM SITE	SALMON
WEN		3		WENAH RIVER MOUTH	GRANDE RONDE
WR1	1	1		WALLOWA RIVER @ KM 14	WALLOWA
WR2			1	WALLOWA RIVER @ KM 32	WALLOWA
ZEN		2		SECESH RIVER @ ZENA RANCH	SALMON
TOTAL	36	93	24		

Kelt reconditioning has a direct impact on population diversity by amplifying life history complexity (Jenkins 2018). Repeat spawner steelhead are very scarce in the vast majority of interior Columbia Basin streams (Keefer et al 2008 and Keefer et al. 2016). Kelt reconditioning infuses consecutive and skip spawn life history variants into the receiving populations. Moore et al. (2014) provided evidence that life-history diversity can diminish variability in population abundance and biomass via portfolio effects. Kelt reconditioning can benefit the receiving population through simply increasing spawner abundance, but also because repeat spawners are larger and have the potential for greater individual productivity (Jenkins et al. 2018). Further, simulations reported by Moore et al. (2014) showed a relationship between the proportion of repeat spawners in the population and the stability and persistence of the steelhead population. Therefore, increasing repeat spawner abundance through kelt reconditioning may stabilize population dynamics through several pathways.

What is the reproductive success of reconditioned kelts?

Reconditioned kelts are able to spawn in the wild and have been shown to have relative fitness similar to that predicted and seen in a study of natural kelts (Seamons and Quinn 2010). We have focused our primary research on relative reproductive success in both Toppenish and Satus creek drainages where lifetime reproductive success of female reconditioned kelts has averaged 2.41 that of single spawn fish as measured across 8 consecutive spawn years. Toppenish and Satus creeks (Yakima River tributaries) were selected as good study locations

after encountering difficult logistical issues in other locations. Post-spawn collections at the Chandler facility provides a reasonable number of fish to recondition. Reconditioned kelts are allowed to overwinter in the mainstem Yakima River prior to the second spawning event which allows for a volitional return and biological (physiological) synchronicity with the natural system. PIT tag arrays detect both first time spawner and reconditioned kelt migration events. Redd surveys allow us to identify areas likely to have high proportions of offspring from anadromous adults, and to minimize the collection of non-target resident fish. Availability of age-0 samples allows identification to brood year to differentiate between first and second spawning events.

Sample Collection

Juvenile collection and analysis were narrowed to Satus and Toppenish Creeks and their tributaries (Hatch et al. 2011). These systems had the best PIT-tag antenna stability, largest number of successfully collected and reconditioned kelts, access to sampling, and lowest impact to spawning habitat, which all translates into good sample sizes for analysis. The biggest downside at both locations is that during good run years, maiden fish comprised a much larger portion than the Prosser reconditioned kelts. Since 2013, sampling locations for juveniles are seen in Figure (2.3). Some sampling sites were not sampled across all years due to access constraints.



Figure 2. 3. Juvenile sampling locations in Satus (lower right) and Toppenish creeks over 8 years 2013-2020.

Results (2013-2020)

The number of juveniles successfully genotyped at the tributary level, and the corresponding number and percentage of samples assigned to at least one anadromous adult parent is shown in Table (2.5). This table does not include locations dropped because of access issues, low sampling success, or lack of assignments to an anadromous offspring. The low apparent assignment rate is due to the low number of adults and juveniles sampled compared to the relative number of unsampled anadromous steelhead and potential resident fish. In Figure (2.4), the largest number of adult fish have 0 progeny detected. This does not indicate that fish are not reproducing but that finding progeny in nature is a difficult proposition, sampling was limited, and missing assignments appears to be equally distributed amongst all groups. This holds true for fish that had a single progeny assignment as well. The second part of the Figure (2.4) has a smaller y-axis to better show the variation between spawning event types. Kelt spawning events appear to have a similar distribution of progeny when compared to maiden events (Figure 2.4).

Table 2. 5. Number of juveniles genotyped by tributary and subsequently assigned to at least one anadromous adult using parentage analysis. Results are reported for individual years and cumulatively across all years.

		Satus Cr.	Toppenish Cr.
2013	Genotyped	227	204
	Assigned	54	64
	% Assigned	24%	31%
2014	Genotyped	285	231
	Assigned	64	67
	% Assigned	22%	29%
2015	Genotyped	341	369
	Assigned	123	165
	% Assigned	36%	45%
2016	Genotyped	790	524
	Assigned	288	187
	% Assigned	36%	36%
2017	Genotyped	442	578
	Assigned	136	172
	% Assigned	31%	30%
2018	Genotyped	615	589
	Assigned	171	165
	% Assigned	28%	28%
2019	Genotyped	651	592
	Assigned	299	239
	% Assigned	46%	40%
2020	Genotyped	630	439
	Assigned	172	77
	% Assigned	27%	18%
Sum	Genotyped	3,300	3,526
	Assigned	1,008	1,136
	% Assigned	31%	32%




Figure 2. 4. Variation in reproductive success of maiden and kelt spawners. Both scales show the number of progeny across the portion of adults genotyped. The second part of the figure has a truncated y-axis to better demonstrate the variability based on number of progeny. POSmaid= Post spawn maiden event, PREmaid= Maiden event collected as pre-spawner, PreKelt= first spawning event for reconditioned kelt, and PostKelt=second spawning event for reconditioned kelt see Monitoring Methods, Genetic Analysis.

The number of genotyped parents confirmed to have entered either Satus or Toppenish creeks is shown in Table (2.6). Pre-spawn maidens have the greatest number of samples with a total of 271 males and 517 females. Post-spawn maidens have only 18 males and 90 females overall.

Reconditioned kelts have 25 male and 277 female Event-1 detections and 30 male and 309 female Event-2 detections. The number of fish increase each year but are limited by the number of kelts that can be collected, and mortality during the reconditioning process. The low number of kelt males is likely due to female skewed returns (Frederiksen et al. 2019), and male spawners staying on spawning grounds seeking additional spawning opportunities until they likely perish.

Table 2. 6. Number of genotyped adults with PIT detections. Only fish with both a PIT tag detection and genotypes were included in the parentage results.

Class	Sex	2013	2014	2015	2016	2017	2018	2019	2020	All
Pre-Spawn Maidens	Male	38	46	57	79	6	21	6	18	271
Post-Spawn Maidens	Male	4	1	7	2	1	1	1	1	18
Reconditioned Kelt Event-1	Male	3	13	7	1	0	0	1	-	25
Reconditioned Kelt Event-2	Male	5	3	13	8	1	0	0	-	30
Pre-Spawn Maidens	Female	88	70	92	141	46	37	20	23	517
Post-Spawn Maidens	Female	12	13	38	9	2	5	5	6	90
Reconditioned Kelt Event-1	Female	15	43	51	22	13	44	40	49	277
Reconditioned Kelt Event-2	Female	74	20	38	56	26	15	44	36	309

The number of progeny assigned to each class of fish is shown in Table (2.7). The majority of assignments are to pre-spawn maidens with 242 juveniles assigned to males and 441 juveniles assigned to females. All other classes of fish have lower numbers assigned as a function of the lower representation in the number of adult fish both detected and genotyped.

Table 2. 7. Number of progeny assigned to each type of spawning event within individual years and across all years. Progeny assigned to adults without tributary PIT tag detections are not included here.

Class	Sex	2013	2014	2015	2016	2017	2018	2019	2020	All
Pre-Spawn Maidens	Male	13	17	24	120	13	34	8	13	242
Post-Spawn Maidens	Male	1	0	0	4	7	1	0	0	13
Reconditioned Kelt Event-1	Male	5	6	4	1	0	0	3	NA	19
Reconditioned Kelt Event-2	Male	3	4	1	0	7	NA	NA	NA	15
Pre-Spawn Maidens	Female	43	41	26	165	76	42	35	13	441
Post-Spawn Maidens	Female	8	1	27	2	0	9	9	1	57
Reconditioned Kelt Event-1	Female	5	17	32	17	63	67	112	30	343
Reconditioned Kelt Event-2	Female	18	12	16	40	39	38	58	9	230

The average number of progeny per parent is shown in Table (2.8). Numbers varied greatly between both years and between classes in within years. Males had ranges between 0.00 and 7.00 progeny per parent.

Table 2. 8. Average number of progeny per parent sampled for each category of fish. Results are reported as the average number of juvenile detections within each spawning event.

Class	Sex	2013	2014	2015	2016	2017	2018	2019	2020	All
Pre-Spawn Maidens	Male	0.34	0.37	0.42	1.52	2.17	1.62	1.33	0.72	1.06

Post-Spawn Maidens	Male	0.25	0.00	0.00	2.00	7.00	1.00	0.00	0.00	1.28
Reconditioned Kelt Event-1	Male	1.67	0.46	0.57	1.00	NA	NA	3.00	NA	1.34
Reconditioned Kelt Event-2	Male	0.60	1.33	0.08	0.00	7.00	NA	NA	NA	1.80
Pre-Spawn Maidens	Female	0.49	0.59	0.28	1.17	1.65	1.14	1.75	0.57	0.95
Post-Spawn Maidens	Female	0.67	0.08	0.71	0.22	0.00	1.80	1.80	0.17	0.68
Reconditioned Kelt Event-1	Female	0.33	0.40	0.63	0.77	4.85	1.52	2.38	0.83	1.46
Reconditioned Kelt Event-2	Female	0.24	0.60	0.42	0.71	1.50	2.53	1.32	0.18	0.94

Relative reproductive success for each category of fish is seen in Table (2.9). Variance in the number's parallels that of the average number of progeny per parent with higher variance seen in males.

Table 2. 9. Relative reproductive success for each category of fish. Results are normalized to the Pre-Spawn Maiden even within in year.

Class	Sex	2013	2014	2015	2016	2017	2018	2019	2020	AVG
Pre-Spawn Maidens	Male	1	1	1	1	1	1	1	1	1
Pos-Spawn Maidens	Male	0.73	0	0	1.32	3.23	0.62	0	0	0.84
Reconditioned Kelt Event-1	Male	4.87	1.25	1.36	0.66	NA	NA	2.25	NA	1.76
Reconditioned Kelt Event-2	Male	1.75	3.61	0.18	0.00	3.23	NA	NA	NA	1.60
Pre-Spawn Maidens	Female	1	1	1	1	1	1	1	1	1
Pos-Spawn Maidens	Female	1.36	0.13	2.51	0.19	0	1.59	1.03	0.29	0.89
Reconditioned Kelt Event-1	Female	0.68	0.67	2.22	0.66	2.93	1.34	1.39	1.47	1.42
Reconditioned Kelt Event-2	Female	0.50	1.02	1.49	0.61	0.91	2.23	0.75	0.32	0.99

Lifetime reproductive success (LRS) of reconditioned kelts is shown in Table (2.10). Male kelt LRS varied between 0.0 and 6.83 times that of fish sampled as pre-spawn maidens within the same year. Across years 2013-2016 male kelts have an LRS of 3.42. Female kelt LRS had annual variation between 1.18 and 3.84 with an average of 2.41.

Table 2. 10. Lifetime reproductive success estimate for male and female Reconditioned kelts. Additive effects of Reconditioned Kelt Events 1 and 2 as normalized to the Pre-Spawn Maiden event.

Class	Sex	2013	2014	2015	2016	2017	2018	2019	2020	AVG
Reconditioned Kelt Lifetime	Male	6.83	4.86	1.54	0.66	NA	NA	NA	NA	3.42
Reconditioned Kelt Lifetime	Female	1.18	1.70	3.71	1.27	3.84	3.57	2.11	1.80	2.41

The presence of kelt offspring demonstrates that reconditioned kelts successfully spawn in the wild. Lifetime reproductive success of female reconditioned kelts was calculated to be 2.41 times that of the pre-spawn maidens. This is similar to findings by Seamons and Quinn (2010) who theorized and found that lifetime reproductive success of natural repeat spawners should scale with the number of breeding spawners. We specifically found that the relative reproductive success of the second spawning event (post reconditioning) for female reconditioned kelts (0.99) is similar to that of putative first-time spawners and demonstrates the potential to boost numbers additively over their first spawning event.

The 2020 spawning event was the eighth consecutive year that we successfully assigned multiple progeny to reconditioned kelts. The methodology of focusing sampling efforts on age-0 fish in areas that anadromous spawning was expected to have occurred, and an increased sampling rate of juvenile has resulted in an increase in the number of successful assignments to both maiden and kelt fish. Future sampling will continue to focus on age-0 fish in areas that spawning was expected to have occurred. We will also start to incorporate analysis of adult offspring using upstream migrants sampled at Prosser Dam or the Chandler facility.

Reconditioned kelt steelhead have demonstrated that they are capable of spawning in the wild. With additional sampling in future years, including adult to adult estimates, we hope to have more accurate numbers and modeling potential. Current data shows that reconditioned kelt steelhead contribute to the productivity of the natural population on a scale similar to that of natural kelts, which will help to preserve this important life history in the Columbia River Basin. We anticipate that we will submit a manuscript on reproductive success in an accredited scientific journal by 2022.

What have we learned about kelt physiology and how does that inform decisions?

Previously, little was known about post-spawning physiology in iteroparous salmonids. There is no evidence for anything like the programmed post-spawning mortality that occurs in the semelparous members of *Oncorhynchus*. Iteroparity is the ancestral trait of *Oncorhynchus* (Crespi and Teo 2002), and the capacity for iteroparity is viewed as a trait of the species *O. mykiss* (Seamons and Quinn 2010). However, individuals may or may not be able to successfully express this trait. Post-spawning kelts experience high mortality due to injuries, susceptibility to pathogens, and bioenergetic exhaustion, as well as anthropogenic changes in the river corridor (Colotelo et al. 2014; Keefer et al. 2018; Keefer et al. 2008). For kelts that survive, little is known about the physiology of gonadal recrudescence, and questions regarding the reproductive status of reconditioned kelts were identified as critical uncertainties during earlier ISRP reviews of the project (ISRP 2011). To fill these knowledge gaps, beginning in 2009, we initiated a research program to investigate the physiology of post-spawning steelhead, with the goal of providing a scientific basis for maximizing the success of steelhead kelt reconditioning programs. Research covered the broad areas of energetics and survival; feeding, digestion and growth; stress; smoltification; disease; and reproduction. We conducted studies using wild kelts, hatchery-origin kelts from Dworshak National Fish Hatchery (DNFH), and post-spawning rainbow trout. Depending on the goals and the experimental system, we used both observational and manipulative experimental designs. Our main methods were non-lethal blood sampling and measurement of plasma hormones and other factors related to physiological status.

Energetics and Survival

Studies conducted by the Department of Fish and Wildlife Sciences at the University of Idaho have described aspects of energy depletion during spawning migration, sexual maturation and kelt emigration in Snake River steelhead (Penney and Moffitt 2014b, 2015). As expected, energy was progressively depleted from early migration to the kelt stage. Muscle lipids were

used more rapidly than muscle protein. Muscle lipid use appeared to slow during fall and winter. Liver lipid stores were depleted more rapidly than muscle stores, and saturated and monounsaturated fatty acids were depleted more rapidly than polyunsaturated fatty acids, which may reflect conservation of polyunsaturated fatty acids due to their role in cell membrane structure and function. At the kelt stage, fish that were rated in “better condition” based on visual appearance had greater energy reserves, consistent with findings on survival in captive reconditioning and natural repeat spawning rates (Hatch, et al. 2013b; Keefer et al. 2008). Kelt length was positively correlated with energy density and muscle protein, suggesting greater energy reserves in larger fish, consistent with a decrease in resting metabolic rate with increasing body size (Clarke and Johnston 1999).

Most mortality in kelt reconditioning occurs over the initial interval after fish are captured after spawning or spawned in captivity (Hatch et al. 2013b; Jenkins, et al. 2020). In the DNFH hatchery, based off kelts undergoing reconditioning, a regression model analyzing factors associated with survival over the first 70 days after spawning found that plasma osmolality and triglyceride levels were positively associated with survival (Fig. 2.5) (Jenkins et al. 2020). In contrast, static measures of energy reserves such as condition factor and muscle lipid level were not significantly associated with survival, and neither were the reproductive measures size-adjusted total egg mass (TEM) or plasma estradiol (E2) level. The osmolality result indicates that osmoregulatory failure in freshwater is a major cause of mortality in kelts, similar to findings on predictors of pre-spawn mortality in Fraser River sockeye salmon (Jeffries, et al. 2011). The plasma triglyceride result indicates that the functioning of physiological systems enabling access to lipid stores is essential for survival. Measures that reflect functioning of physiological systems performed better as indicators of fish condition than static measures such as condition factor. Kelt use of muscle protein stores were not assessed in this study, and would be a potential future direction, as use of protein stores for energy can be significant in salmonid physiology (Mommensen 2004).




Figure 2. 5. (A) Standardized coefficients $[(x-\text{mean})/SD]$ with 95% confidence intervals of potential predictors of survival in a multiple logistic regression model of female steelhead trout at Dworshak National Fish Hatchery, Ahsahka, Idaho to 70 days after first spawning in 2015 and 2016 ($n = 291$). Potential predictors were measured at spawning. Positive coefficients indicate a greater probability of post-spawning survival. Coefficients with confidence intervals that do not overlap zero were statistically

significant ($P < 0.05$) and are marked with an asterisk. (B) Post-spawn survival probabilities (PSurv) to 70 days, as predicted by plasma triglycerides (mg/dL) (TG) and plasma osmolality (mmol/kg) (OS) measured at first spawning in 2015 and 2016 ($n = 291$). From Jenkins et al. 2020.

Feeding, Digestion and Growth

Growth is minimal over the period after spawning in both hatchery and wild kelts (Jenkins, et al. 2019; Pierce, et al. 2017). Many fish continue to lose weight over the first 10 weeks after spawning, whereas even among fish with positive growth, growth rates are 5 to 10-fold lower than later during reconditioning (Jenkins et al. 2019). Suppressed growth during this period is due to minimal feed consumption, although fish are offered highly palatable feeds such as krill, multiple times throughout the day. However, in kelts that were fasted for the first 10 weeks after spawning, growth rates immediately increased after food was made available, and exceeded growth rates of fish that had been fed throughout, a compensatory growth response to the fast (Fig. 2.6) (Hatch et al. 2020; Jenkins, et al. In preparation). This implies that there is a process of recovery from spawning that must occur before feeding and growth can begin in earnest, and that this process does not necessarily depend on feeding. Feeding motivation and feed consumption are strongly reduced during the recovery period. The physiology underlying the recovery period is not well understood. However, it is reasonable to hypothesize that it may relate to restoration of digestive system function after a prolonged fast, and reversal of the catabolic state associated with elevated levels of the stress hormone cortisol that occurs during the spawning period in salmonids (Barry, et al. 2010; Dickhoff 1989; Schreck, et al. 2001).




Figure 2. 6. Specific growth rate in mass (MSGR) in female steelhead trout sampled at 10-week intervals starting at first spawning. Food was withheld from Fasted fish for the first 10 weeks after spawning. Symbols indicate mean and bars indicate SEM. Above each sampling point, bolded letters indicate significance (R = Reproductive status, F = Fasting Treatment, R*F =

Interaction effect). Rows of letters below the figure indicate significant differences over time within each treatment group. Years are shown separately due to a significant year effect at week 20-30. With years combined, Fasted fish showed significantly higher MSGR than Fed fish over week 10-20. From Jenkins et al. In Preparation.

Maintenance of a functioning digestive system is energetically costly, and animals that routinely fast for prolonged periods often reduce this cost by atrophy of the digestive epithelia (Secor, et al. 2002; Zaldua and Naya 2014). The gut-somatic index decreases linearly over time in fasted juvenile rainbow trout with a loss of ~40% of the relative mass of the gut over 147 days of fasting (Simpkins, et al. 2003). In Atlantic salmon that fasted for 50 days, restoration of the gut upon refeeding required at least one week, during which feed intake was reduced (Krogdahl and Bakke-McKellep 2005). As would be expected after their prolonged fasting spawning migration, the stomach and intestinal epithelia are atrophied in post-spawning summer run steelhead trout, and degenerative changes are found in the liver (Penney and Moffitt 2014a). Some kelts showed evidence of regeneration of the gut epithelia, although it was not clear whether this was a consequence of or preparation for feeding activity. In some species, food in the lumen of the gut stimulates restoration of function (Secor et al. 2002), and force feeding has been successfully employed in Atlantic salmon kelt reconditioning projects (Eales, et al. 1991) and with precocious Chinook salmon parr (Bernier, et al. 1993). To determine if this might be effective with steelhead kelts, we force fed DNFH hatchery origin kelts after spawning (Hatch, et al. 2012). However, this resulted in decreased survival, and the stomachs of mortalities contained partially digested food 35 days after force feeding. This suggests that there was a delay in digestion associated with the restoration of digestive function. The metabolic cost of restoration of gut function may have outweighed benefits obtained from the food. Based on these results, force feeding does not appear to be a promising method for steelhead kelt reconditioning.

The decreased feeding motivation of kelts during the period after spawning may be due to changes in the levels of orexigenic and anorexigenic endocrine and neuroendocrine factors. One such factor is ghrelin, a strongly orexigenic (appetite stimulating) hormone produced by the stomach (Kojima and Kangawa 2005). Unlike other appetite stimulating factors, ghrelin crosses the blood-brain barrier. In humans, ghrelin is reduced in stomach atrophy (Osawa, et al. 2005). Thus, it is possible that steelhead kelts may have reduced appetite due to decreased plasma ghrelin caused by stomach atrophy. Ghrelin is highly conserved, and commercially available mammalian ghrelins are effective in fish species. In most trials in fish, ghrelin administration increased food intake (Riley, et al. 2005; Shepherd, et al. 2007; Unniappan and Peter 2004, 2005). However, in several studies, appetite has not been stimulated (Jonsson, et al. 2007), or has even been suppressed (Jonsson, et al. 2010). To determine if ghrelin has potential as an appetite stimulant in kelt steelhead, we tested ghrelin administration in juvenile rainbow trout (Branstetter, et al. 2010). Fish were intraperitoneally implanted with Alzet micro-osmotic pumps that administered rat ghrelin or vehicle at a constant rate over time and tracked for 20 days. Feed consumption and growth were quantified. Fish in the ghrelin treated tanks exhibited unusual behavior. During feeding, fish were at the top of the tank, with their backs out of the water, and were highly active. Ghrelin treated fish pursued pellets, but often did not ingest them. Fish in the control treatment tanks did not show these behaviors. Feed intake in

ghrelin treated tanks was lower than controls, and growth rates were lower than controls. Thus, ghrelin appeared to inhibit appetite and growth under our experimental conditions. As well as stimulating appetite, ghrelin stimulates corticotrophin-releasing factor (CRF) neural circuits in the brain (Jonsson et al. 2010). Stimulation of the CRF system produces increases in activity level and suppresses feed intake (Lowry and Moore 2006), consistent with our observations on the behavior of ghrelin treated fish. These findings suggest that the effect of ghrelin administration may depend on the relative magnitude of the effects mediated by these different mechanisms. Due to the complexities involved, we decided not to pursue ghrelin treatment any further.

Stress

The stress axis or hypothalamic-pituitary-interrenal (HPI) axis is the central endocrine pathway mediating the stress response in fish (Barton 2002; Wendelaar Bonga 1997). Activation of the HPI axis results in an increase in circulating corticosteroids, such as cortisol (Medeiros, et al. 2010; Wendelaar Bonga 1997). High circulating cortisol levels are found during the final stages of gonadal maturation and during spawning in salmonids (Laidley and Leatherland 1988). Cortisol initially plays an adaptive function during stress, but chronic elevations of cortisol contribute to the deleterious effect of chronic stress (Barton and Iwama 1991). Chronic cortisol elevation induces a catabolic state in which protein synthesis is inhibited and energy is mobilized from carbohydrate, protein and lipid stores and immune function is suppressed (Mommsen, et al. 1999; Wendelaar Bonga 1997) (Schreck 1981, 1990). Elevated cortisol levels and the resulting catabolism, anorexia, and suppression of immune function are proposed to cause post-spawning death in the semelparous *Oncorhynchus* species (Dickhoff 1989). In addition, activation of the stress axis inhibits the reproductive endocrine axis (Barton 2002; Donaldson 1990). Therefore, we hypothesize that survival, recovery from spawning, and gonadal recrudescence in steelhead kelts may depend on a post-spawning reduction in circulating cortisol levels. Although basal (unstressed) circulating cortisol levels are difficult to measure due to the rapid response to capture stress, we have indirect evidence that it takes approximately 6-10 weeks for cortisol levels decrease after spawning in rainbow trout. Plasma levels of the growth stimulatory hormone insulin-like growth factor-1 (IGF1), for which we have recently established a laboratory assay (Medeiros, et al. 2020), increased after spawning for 6-8 weeks in rainbow trout, and did not respond to rations until 8 weeks (Hatch et al. 2020). Consistent with its catabolic function, cortisol suppresses IGF1 (Pierce, et al. 2005a). Liver gene expression levels of IGF binding protein 1b (IGFBP1b) decreased for 8-10 weeks after spawning in rainbow trout (Fig. 2.7) (Caldwell, et al. 2013). IGFBP1b is an indicator of stress and is directly induced by corticosteroids (Kajimura and Duan 2007; Pierce, et al. 2006). A fuller characterization of post-spawning changes in the HPI axis would be very useful.




Figure 2. 7. Mean (\pm SEM) liver insulin-like growth factor binding protein 1 (*igfbp1*) mRNA levels over time in female rainbow trout (A: two-year-old; B: three-year-old) fed a control-ration or restricted-ration. From Caldwell et al. 2013.

Along with studies describing the recovery from spawning, we conducted one manipulative study. In teleosts, most of the long-term changes in response to cortisol are mediated by cytosolic glucocorticoid receptors (GRs) located in target tissues, although other receptors exist. The GR antagonist mifepristone (RU486) blocks the physiological response to cortisol in salmonids and other teleosts (McDonald and Wood 2004; McDonald, et al. 2004; Medeiros, et al. 2014; Medeiros and McDonald 2013; Rodela, et al. 2009), interrupting the cortisol signaling cascade that leads to depressed appetite and the other undesirable effects of chronic stress. To determine if long term RU486 administration might speed up recovery from spawning, we tested if this treatment was effective in increasing growth in post-spawning female rainbow (Hatch, et al. 2017). Post-spawning rainbow trout were implanted with a time release formulation of RU486 in a mixture of vegetable shortening and vegetable oil, or vehicle alone. However, high mortality (50%) occurred over the 42-day course of the experiment, possibly associated with the poor condition of the fish available for the experiment and repeated handling. Mortality did not differ between treatment groups, and no differences were detected in post-spawning growth. This experiment needs to be repeated with a higher number of surviving fish before any conclusions regarding the efficacy of the treatment can be made.

Smoltification

The ability to osmoregulate in saltwater increases with body size in salmonids, and consequently kelts would not be expected to need to undergo smoltification as a preadaptation to adjust to saltwater (Clarke and Hirano 1995). However, numerous other physiological changes occur during smoltification, which may be necessary for fish to successfully make the transition to the ocean (McCormick 2012). Gill sodium-potassium ATPase activity increased in kelts as compared with spawning steelhead, however, plasma thyroxine levels were lower in kelts (Buelow and Moffitt 2015). Thus, it is unclear whether kelts undergo a preadaptation process similar to smoltification, since both factors typically increase during smoltification (McCormick 2012). Further study is needed to characterize smoltification related changes in steelhead kelts. Archival tagging studies have generally not found extensive mortality associated with ocean entry in steelhead kelts, although numbers are low (Hayes, et al. 2012;

Miller, et al. 2021; Nielsen, et al. 2011; Null, et al. 2013). Plasma ion concentrations were higher in better condition steelhead kelts in the Clearwater River, suggesting that loss of the ability to osmoregulate in freshwater may be an early indicator of impending mortality (Buelow and Moffitt 2015). Our results in hatchery kelts reconditioned in freshwater confirm this supposition (Fig. 2.5) (Jenkins et al. 2020). Thus, based on the limited information currently available, it appears that freshwater osmoregulation is more critical to the survival of kelts than saltwater osmoregulation.

Disease

Immunosuppression and susceptibility to opportunistic pathogens are observed during spawning in multiple fish species. In steelhead kelts, these problems are exacerbated due to bioenergetic exhaustion and injuries sustained during spawning, downstream migration, and collection at dams and weirs. The crowded conditions the fish experience in culture also greatly facilitate disease transmission. The principal opportunistic pathogens that impact kelts are external *Saprolegnia* infestation, bacterial infections, and gill copepods. Consistent daily formalin treatment to control *Saprolegnia* outbreaks has been found to be essential to minimize mortality in kelt reconditioning. The installation of automated formalin treatment systems has resulted in reduced mortality at both the Prosser and Dworshak reconditioning sites. Prophylactic treatment of kelts at intake by injection with oxytetracycline, a broad-spectrum antibiotic, has been found to decrease mortality due to bacterial infection. In 2009, we did not treat fish at intake with oxytetracycline, due to concerns from a fish disease specialist employed by the US Fish and Wildlife Service about excessive antibiotic use, and experienced significantly higher mortality (Branstetter, et al. 2009).

Infestation of fish gills with freshwater copepod ectoparasites (*Salmincola* spp.) is a chronic problem in steelhead and Atlantic salmon kelt reconditioning (Hatch, et al. 2013a; McGladdery and Johnston 1988), which can cause mortality and suppress rematuration (Branstetter, et al. 2007; Duston and Cusack 2002). Treatment of kelts for parasitic copepods is necessary for successful reconditioning. The standard treatment for copepods in the Columbia River Basin had been ivermectin gavage (Johnson and Heindl 2001; Roberts, et al. 2004). However, neurotoxic effects of ivermectin have also been reported (Johnson, et al. 1993; O'Halloran, et al. 1992; Palmer, et al. 1987), and were also observed after ivermectin administration to kelts. We evaluated and implemented an alternative treatment using emamectin benzoate, which was more effective at controlling copepods and resulted in substantially reduced mortality (Fig. 2.8) (Branstetter, et al. 2011; Hatch, et al. 2014; Hatch et al. 2012).




Figure 2. 8. Mortality of Prosser kelts after ivermectin or emamectin treatment at intake. Log-rank test mortality curve comparison, 2011: $p = 0.0019$; 2012: $p = 0.0340$. From Hatch et al. 2013b.

Reproduction

The bulk of the progress to date in research on kelt steelhead physiology has been in the area of female reproductive development. We have documented that the consecutive and skip spawning life histories are found in reconditioned kelts, established methods for screening fish for maturation status before release, and estimated the expected productivity benefit from released consecutive and skip spawners spawning in the wild. We have evaluated a hypothesized trade-off between reproductive investment in maiden spawning and post-spawning survival and post-spawning reproductive investment. As part of this work, we found interesting relationships between growth and nutritional status early during oogenesis and subsequent reproductive parameters at repeat spawning, shedding light on the physiological mechanisms regulating egg size and fecundity. Finally, we have investigated the timing of the rematuration decision and factors associated with it, establishing that initiation of rematuration occurs early during reconditioning, that it is associated with faster growth and better nutritional status, and that both physiological status at spawning and post-spawning feeding conditions influence the rematuration decision. This implies that, although optimal conditions in reconditioning can maximize consecutive spawning rates, pre-capture environmental conditions likely set a limit to consecutive spawning rates. We have found no evidence for functionally significant regulation of reproductive trajectory in steelhead at a second “permissive” decision window proposed for Atlantic salmon. Much of the progress in this area has been made possible by access to hatchery-origin kelts at DNFH.

The consecutive and skip spawning life histories are found throughout iteroparous seasonally spawning teleost species, and substantially impact fisheries management and conservation (Rideout, et al. 2005; Rideout and Tomkiewicz 2011). Consecutive spawners initiate gonadal rematuration on schedule to spawn at a one-year interval after initial (aka “maiden”) spawning, whereas skip spawners rest for a year, typically spawning at a two-year interval, although longer resting intervals are possible. In salmonids, post-spawning life histories are plastic: multiple repeat spawners can spawn at different intervals (Chaput and Jones 2006; Halttunen

2011; Hubley and Gibson 2011; Moore, et al. 2014; Riva-Rossi, et al. 2007). Larger, older fish, with a longer migration distance and/or later spawn timing are more likely to skip spawn (Chudyk 1976; Jonsson, et al. 1991; Keefer et al. 2018; Keefer et al. 2008; Moore et al. 2014). In general, both the presence of repeat spawners, and variation in repeat spawning interval increase population stability via the portfolio effect (Moore et al. 2014; Schindler, et al. 2010). Repeat spawners are also more fecund than maidens due to their larger size, multiplying their productivity benefit (Chaput and Jones 2006; Halttunen 2011). Early studies on wild steelhead kelts undergoing reconditioning confirmed that the consecutive and skip spawning life histories are found in reconditioned kelts (Fig. 2.9) (Pierce et al. 2017). Plasma levels of estradiol (E2), the principal female steroid, and vitellogenin (VG), a phospholipoprotein produced by the liver that functions to shuttle energy into the ovary during oogenesis, diverged into two groups by early fall: consecutive spawners had high levels of both factors, and potential skip spawners had lower levels. The migration patterns of released fish from the two categories concurred with their reproductive status, with consecutive spawners migrating upriver at higher rates, and skip spawners occasionally detected migrating upriver the following season. The time course of the increase in plasma E2 and VG was such that complete separation occurred by late July to mid-August. This enabled blood sampling and identification of consecutive and skip spawning individuals before release time, and appropriate management of the two life history types, which has now become a routine part of the project. Visual assessment of maturation status of Prosser kelts at release time was approximately 70% correct, which is not surprising given that this is five or more months before spawning.




Figure 2. 9. Relationship of plasma levels of estradiol (E2) and vitellogenin (VG) in reconditioned female kelts at pre-release sampling in the fall of 2009 (A: 10/29/2009), 2010 (B: 10/13/2010), and 2011 (C: 10/13/2011). Fish were grouped by cluster analysis (Ward linkage, 2 groups specified, data standardized), and the resulting groups are indicated by circles (rematuring fish with high E2 and VG) and triangles (non-rematuring fish with low E2 and VG). Proposed threshold values for screening fish for maturation status by plasma E2 (500 pg ml⁻¹) and VG (0.1 mg ml⁻¹) level are indicated by lines. From Pierce et al. 2017.

The most significant potential criticism of kelt reconditioning projects, in our view, is a hypothesized reproductive life-history tradeoff between reproductive investment in maiden spawning and energy retained for subsequent survival and investment in repeat spawning. We believe that this is the issue underlying ISRP statements from more than decade ago suggesting that kelt reconditioning might select for reduced fitness (ISRP 2008). A recent adult-to-adult reproductive success study on Hood River steelhead found that iteroparous individuals had reduced success on their maiden spawning, which was interpreted as being due to reduced maiden spawning investment (Christie, et al. 2018). However, there are alternative explanations for these results, and the existence and magnitude of any potential trade-off have not been directly assessed. Using the DNFH hatchery kelt model, we were able to directly test these predictions, both in terms of survival and investment in repeat spawning. Size-adjusted maiden spawning total egg mass (TEM), a reasonable measure of reproductive investment, was not significantly related to survival after spawning (Fig. 2.5) (Jenkins et al. 2020). However, maiden TEM was the only factor that predicted consecutive spawning, and the relationship was positive: fish with greater maiden TEM were more likely to become consecutive spawners (Fig. 2.10) (Jenkins et al. 2020). Consecutive spawning requires greater investment of energy during the period after spawning than skip spawning, so this result shows that individual females with greater maiden reproductive investment also invest more in repeat spawning. As described in Stearns' analysis of life-history tradeoffs, this result implies that variation in condition between individuals is of greater importance than a tradeoff between current and future reproduction in determining reproductive investment, as found in other studies (Hendry, et al. 2000; Stearns 1992). Thus, if anything, kelt reconditioning is selecting for individuals that are best adapted to current conditions.




Figure 2. 10. (A) Standardized reproductive status coefficients ($[x\text{-mean}]/SD$) with 95% confidence intervals of potential predictors in a multiple logistic regression model of reproductive status in consecutive years of female steelhead trout ($n = 60$) following manual spawning in 2015 and 2016. Positive coefficients indicate greater probability of consecutive reproductive status. Coefficients with confidence intervals that do not overlap zero were statistically significant ($P < 0.05$) and are marked with an asterisk. (B) The probability of reproductive status in consecutive years in reconditioned female steelhead trout at Dworshak National Fish Hatchery, Ahsahka, Idaho, in relation to total egg mass at first spawning (g, mass-standardized, $n = 60$). The bars at the top and bottom of the graph represent the number of fish that became reproductively active or remained non-reproductive following first spawning, respectively, for each range of total egg mass. From Jenkins et al. 2020.

As well as evaluating potential relationships of maiden spawning reproductive performance to post-spawning outcomes, the DNFH hatchery kelt model enabled us to directly assess repeat spawning reproductive performance in reconditioned kelts. Based on issues with spawn timing in captive broodstock programs, we wanted to determine whether ovulation timing was shifted in reconditioned kelts. Consecutive spawning fish ovulated the same week on average as their maiden spawning, whereas skip spawners ovulated 3 weeks earlier (Fig. 2.11) (Jenkins, et al. 2018). Earlier migration timing for natural skip spawners versus maiden or consecutive spawners has been reported in Atlantic salmon and steelhead (Keefer et al. 2018; Niemela, et al. 2006). Ovulation timing for both consecutive and skip spawners was within the range for maiden spawners in this population. Consecutive spawners had 1.28-fold, and skip spawners 1.52-fold, the expected productivity of maiden spawners, based on total egg mass. When standardized for fish size, consecutive spawners had relatively smaller and skip spawners relatively larger eggs, than maiden spawners. These results are similar to findings in repeat-spawning Atlantic salmon (Reid and Chaput 2012). Interestingly, measures of energy balance such as growth rates and plasma triglyceride level correlated positively with size-standardized egg size at 10 weeks after spawning and with size-standardized total egg mass and fecundity at 20 weeks after spawning (Jenkins et al. 2020). This suggests that egg size and fecundity are set based on physiological condition during early- to mid- oogenesis, and that egg size is set first, in repeat spawning steelhead and potentially in other salmonids. If generally true, this might inform evolutionary models on optimal egg size and the egg size/fecundity tradeoff (Einum, et al. 2002; Hendry and Day 2003; Kinnison, et al. 2001), and provide a mechanism linking environmental conditions with reproductive performance.




Figure 2. 11. Spawn week fidelity in repeat spawning female steelhead trout from the Clearwater River, ID sampled between 2013-2015. Boxes indicate the interquartile range, the line indicates the median, whiskers show the data range, and all points are shown. Repeat spawn week did not differ significantly from maiden spawn week in consecutive spawners (one sample t-test, $t_{22} = 0.2630$, $p = 0.7950$), but was significantly earlier in skip spawners (one-sample t-test, $t_{19} = 5.4978$, $p < 0.0001$), and repeat spawn week was significantly earlier in skip spawners than consecutive spawners (two-sample t test, $t_{41} = 3.671$, $p = 0.0007$). From Jenkins et al. 2018.

Because of the importance of consecutive and skip spawning in steelhead kelt reconditioning programs, we have sought to better understand the physiological decision underlying the development of these two life histories. Work in this area has been guided by the critical period model of salmonid maturation decisions. This was developed for Atlantic salmon (*Salmo salar*) and proposes that the initial maturation decision takes place during a seasonally defined critical period approximately one year before spawning and is gated by stores of metabolic fuels (Thorpe 2007; Thorpe, et al. 1998). The aspect of metabolic fuel storage that regulates entry into a reproductive cycle is not known but is functionally defined as either the absolute level or the rate of change of body size (growth) and/or lipid reserves (Taranger, et al. 2010; Thorpe 2007). Endocrine signaling factors associated with lipid metabolism and/or growth are presumed to provide the mechanistic link between energetic status and the reproductive endocrine axis (Taranger et al. 2010). A second “permissive” critical window is proposed to occur approximately 6 months later, during which maturation can be arrested if energy stores are not sufficient. Thus, consecutive spawners must initiate rematuration during the first window and continue at the second, whereas skip spawning results from either not initiating at the first window or arresting at the second. However, the timing of the critical periods and the relevant aspect of metabolic fuel storage and network of associated signaling factors are not known in detail in any species. In steelhead kelts, the extreme energy depletion from maiden spawning migration and the maiden spawning event itself would be expected to influence this decision process.

Evidence from wild kelts, our DNFH hatchery kelt model, and the post-spawning rainbow trout model shows that the functionally important critical period occurs early during the post-spawning period and may extend to before spawning. Plasma levels of estradiol (E2), the principal female reproductive steroid in salmonids, reliably diverged between consecutive and skip spawners at 20 weeks post-spawning in DNFH kelts (Fig. 2.12) (Jenkins et al. 2019). The timing of the increase was somewhat sooner at approximately 13 weeks in wild kelts (Pierce et al. 2017), whereas the difference occurred at 8-10 weeks in post-spawning rainbow trout (Caldwell, et al. 2014). The difference in timing likely relates to metabolic rate, which would be slower in the DNFH fish due to their larger size and the colder water temperatures at this site due to earlier spawn timing. Elevations in plasma E2 only occur after upstream events in the brain-pituitary-ovary-liver endocrine axis (Lubzens, et al. 2010; Taranger et al. 2010; Wootton and Smith 2015). The amount of time required for these events is not known with any certainty; however, in maturing coho and Atlantic salmon maiden females, increases in plasma E2 were first detected 10 months before ovulation (Andersson, et al. 2013; Campbell, et al. 2006). Thus, the increase in plasma E2 sets a later bound to the initiation critical period. Examination of individual profiles of E2 over time do not show more than a few individuals where E2 initially increases, and then decreases after the time of the purported second permissive critical period (Jenkins et al. 2019; Pierce et al. 2017). Thus, we do not find support for functionally significant regulation of rematuration during this window in steelhead kelts. This part of Thorpe's model makes sense for fall spawners like Atlantic salmon, which may need to reassess energy reserves after the winter. However, summer steelhead would need to begin migration from ocean feeding areas toward river mouths in mid-summer, and thus the second critical period makes little sense in the context of the summer steelhead life history.




Figure 2. 12. Plasma estradiol-17 β concentrations in female steelhead trout from the Clearwater River, Idaho, sampled in 2015 (A) and 2016 (B) at 10-week intervals following maiden spawning. Females were grouped as rematuring (red: 2015, N=13; 2016, N=12) or non-rematuring (blue: 2015, N=29; 2016, N=18). Box heights indicate interquartile range, horizontal lines within indicate the median, and whiskers show the data range. Time points within a rematuration group sharing the same letter do not differ significantly (One-Way ANOVA followed by Tukey's Multiple Comparison Test, $P < 0.05$). Asterisks indicate significant differences between rematuration groups at each time point (T-test, $P < 0.05$). From Jenkins et al. 2019.

Initiation of maturation as a consecutive spawner during the post-spawning period is associated with greater growth during this period, although growth is minimal during recovery from spawning in kelts and rainbow trout (Fig. 2.6) (Caldwell et al. 2013; Jenkins et al. 2019; Pierce et al. 2017). Plasma triglyceride levels are elevated in consecutive spawners at 10 weeks post-spawning, consistent with greater feeding during this period versus skip spawners (Jenkins et al.

2019). To clarify the relative roles of physiological condition at spawning and post-spawning feeding conditions on the consecutive maturation decision, we conducted an experiment in which DNFH hatchery kelts were fasted during the first 10 weeks after spawning. To explore the role of the growth hormone/insulin-like growth factor (GH/IGF) endocrine axis in reproductive decisions in steelhead kelts, we used recently developed assays in our laboratory (Medeiros et al. 2020) to measure plasma GH and IGF1 levels in this experiment. The GH/IGF axis is the principal physiological system that regulates growth in salmonids, as in other vertebrates (Norbeck, et al. 2007; Perez-Sanchez, et al. 2018; Wood, et al. 2005). During fasting and under other catabolic conditions, GH increases, and functions to mobilize stored energy (Bergan-Roller and Sheridan 2018; Bjornsson, et al. 2018; Norbeck et al. 2007; Pierce, et al. 2005b). Thus, elevated plasma GH can be used as an indicator of catabolic status. Some DNFH hatchery females in both the fed and fasted groups initiated rematuration as consecutive spawners. However, fish in the fasted group that were able to do this had significantly lower plasma GH levels at spawning than other groups, indicating that they were less catabolic at this time point (Fig. 2.13) (Jenkins et al. 2020). Plasma GH also increased during the fast as in subadult salmonids (Bjornsson et al. 2018; Pierce et al. 2005b). From this, we can conclude that 1) both metabolic state at spawning and post-spawning feeding conditions regulate consecutive spawning, 2) post-spawning feeding conditions were more important under our experimental conditions, 3) GH is a good indicator of metabolic state, and 4) the GH/IGF axis responds to nutritional status even after the prolonged fasting spawning migration of summer steelhead. These results imply that both pre-capture conditions and conditions in culture will affect consecutive spawning rate and suggest that steelhead show physiological adaptation to a period of continued fasting after spawning.




Figure 2. 13. Growth Hormone in female steelhead trout sampled at 10-week intervals starting at first spawning. Analysis included Fed-Reproductive and Fed-Non-reproductive (closed and open circles, $n = 31, 35$) and Fasted-Reproductive and Fasted-Non-reproductive (closed and open triangles, $n = 24, 44$). Symbols indicate mean and bars indicate SEM. Above each sampling point, bolded letters indicate significance (R = Reproductive status, F = Fasting Treatment, $R*F$ = Interaction effect). Rows of letters below the figure indicate significant differences over time within each treatment group. At each sampling point with a significant interaction effect, symbols sharing the same letter do not differ significantly. From Jenkins et al. In Preparation.

What is the homing fidelity of reconditioned kelts?

To investigate repeat homing fidelity in artificially reconditioned steelhead kelts, we split our investigations into a larger sub-basin level homing fidelity and a much smaller sub-population level homing fidelity. For sub-basin level homing fidelity, we are using the Yakima River as our test population to investigate if post spawn steelhead kelts that are captured emigrating out of the Yakima River, reconditioned, released, then return to the Yakima River basin to spawn a second time. The Yakima River steelhead population consists of four genetically unique major population groups (MPG): the Upper Yakima River, encompassing the mainstem and all tributaries above the confluence with the Naches River; the Naches River system, including Ahtanum Creek and Yakima River mainstem from the confluence of the Naches River down to Toppenish Creek; Toppenish Creek; and Satus Creek (Loxterman and Young 2003). To examine homing fidelity on the sub-population level we investigated the same parameters on a smaller scale using Satus Creek, Toppenish Creek, Ahtanum Creek, Taneum Creek, above Roza Dam (Upper Yakima population) in the Yakima River Basin, and Omak Creek in the Okanogan River

Basin as our test populations to determine if kelts return to the same locations to spawn a second time.

Sub-basin level homing of reconditioned kelt steelhead

After excluding reconditioned kelts that were unsuitable for measuring homing fidelity (see methods), we ended up with a total of 1,103 fish from 2012-2021 that were used for the sub-basin homing study (Hatch et al. 2020). Out of the reconditioned kelts that met the qualifications, a total of 1,103 individual fish were last detected moving upstream at a Yakima Basin mainstem dam PIT tag array, detected at tributary PIT tag arrays, recaptured at Roza Dam prior to spawning, or were recaptured as post spawned kelts at the Chandler Juvenile Collection Facility the following emigration season. Only 2 reconditioned kelts had final PIT detections in another sub-basin outside of the Yakima River during this nine-year period. One fish had a final detection on 4/16/2014 in the Walla Walla River and the other was detected in Asotin Creek (tributary of the Snake River) on 4/9/2020. At the sub-basin level this equates to an extremely low stray rate of 0.2%. As noted in Table (2.10) below, we did have an additional 3 fish with final detections at Ice Harbor, Little Goose, and Lower Granite Dams on the lower Snake River during March or April. Although these reconditioned kelts could not be assigned to a specific spawning tributary to be classified as strays at the basin scale, their inclusion in our analysis would increase the stray rate of reconditioned kelts to only 0.5%.

Sub-population level homing of reconditioned kelt steelhead

To investigate homing fidelity at the sub-population level, a total of 146 reconditioned kelts met the qualifications for this analysis (see methods). Each of these fish had sub-population maiden spawner detections and subsequent detections post reconditioning. Maiden spawner origin was determined through detections of first-time spawners outfitted with PIT tags at tributary level or sub-population detection arrays, kelt steelhead collected at a tributary weir, or through parentage analysis of progeny collected in tributaries that confirmed maiden spawning locations. In the Yakima River, all 65 previously PIT tagged fish that were detected as a maiden spawner and reconditioned kelts returned to spawn in the same tributary as their original detection or were detected again at Roza Dam (Upper Yakima population). We have found no evidence of straying in this group of fish.

We also used an ongoing reproductive success study in Satus and Toppenish Creeks to assign Age-0 juveniles to fish later taken into the reconditioning program (Hatch et al. 2020). These parentage results were used as proxy locations for comparison to PIT detections following reconditioning and release from the kelt reconditioning program. A total of 70 fish had both a proxy maiden location (parentage results) and a post reconditioning tributary PIT detection. All fish demonstrated fidelity to their maiden location, with 34 returning to Satus Creek and 36 to Toppenish Creek. No fish were detected at other PIT tag arrays that would have indicated straying behavior.

In Omak Creek (Okanogan River tributary), 11 kelt steelhead were collected at the weir migrating out of the tributary and following reconditioning were released near the mouth of the Okanogan River. All 11 fish were later detected back at the Omak Creek weir on their repeat

spawning run. At the sub-population level, we had 100% homing fidelity post reconditioning, as all 146 fish were detected at the same PIT tag arrays or recaptured at the same location as their maiden spawning event (Table 2.10).

Table 2.10. Steelhead homing in the Yakima River and Omak Creek based on PIT detections.

Location	Sub-Basin Level Homing		Sub-Population Level Homing			
	Unknown Maiden Tributary		PIT Tag First - Second		Progeny - PIT Second	
	Rehome	Stray	Rehome	Stray	Rehome	Stray
*Yakima R.	1,103	2 (3)	65	0	70	0
Omak Cr.	—	—	11	0	—	—
Totals	1,103	2 (3)	76	0	70	0

*We did find three reconditioned kelts with final detections at one of the four lower Snake River dams in March/April. These fish were not detected at a sub-basin or sub-population level, but could be considered strays at the basin scale.

Where and when should we release fish?

Discussion / data comparing different locations and consecutive / skip spawner life histories *Geographic Comparison of Reconditioning Programs*

Survival and maturation data from Prosser, Winthrop, and Dworshak hatcheries are shown in Figure (2.14). Survivals in the Prosser project increased beginning in 2012 and have been in the 70-80% range from 2016 to 2020. In 2012, the Prosser project began treating all kelts with emamectin benzoate by intraperitoneal injection for copepod infestation. Previous treatment had been with ivermectin by gavage. We attribute the increased survival to implementation of a less toxic treatment. The performance of the Prosser project over the past 5 years has been exceptional and is a tribute to the quality of the fish care in this project. The Prosser project also has the advantage of collecting fish at the reconditioning location, whereas fish must be transported from the collection location to the reconditioning location in other projects (Hatch et al. 2020). Survival of kelts collected at Lower Granite Dam increased to levels similar to the Prosser project in 2018 and 2019 (not including the mortality event at NPTH in 2019), suggesting that survival rates similar to that found in the Prosser project may be achievable with the Snake River fish (Hatch et al. 2020). Survival of fish at both Winthrop and in the DNFH decreased in 2020, which can be attributed to the poorer overall fish condition in 2020, which was noted at both sites. The reasons for this can vary annually, but is likely a result that 2020 was a low run year, and collection criteria were adjusted to maximize collection, which included fish in poorer condition. Survivals of DNFH hatchery fish were somewhat lower than found for wild origin fish, which may be due to the effects of anesthesia and manual spawning at the hatchery. Further, hatchery returning steelhead have been lethally spawned at DNFH since the hatchery was established in the 1970s, which could have resulted in selection against iteroparity. Overall, results suggest that survivals above 50% are attainable in CRB kelt reconditioning, even in inland populations with a long migration (Hatch et al. 2021).




Figure 2. 14. Survival and female consecutive and skip maturation rates in CRB kelt reconditioning projects. Snake River fish include kelts collected at Lower Granite Dam (LGR) and air spawned hatchery origin kelts from the DNFH stock (DNFH HOR). Maturation data for skip spawners is from non-mature fish from the previous season held over for an additional year. Prosser 2020 consecutive maturation rates are not shown, because maturation was not assessed by plasma E2 level.

With a few exceptions, consecutive rematuration rates in the Prosser project have consistently been near 60%. Maturation rates for Snake River and Upper Columbia fish have generally been lower, and overall, the ranking over the past few years has been Prosser > Upper Columbia > Snake River, which is consistent with the hypothesis that fish with a longer and energetically more demanding migration tend to repeat spawn as skip spawners (Keefer, et al. 2008). The consecutive maturation rate for 2020 at Prosser is not shown because fish were not blood sampled due to the COVID-19 pandemic (Hatch et al. 2021). Based on visual appearance, the 2020 Prosser maturation rate was 83.8%, but this should not be compared to other maturation rates as it is not based on plasma estradiol level. Some of the variation in maturation rates is attributable to conditions during reconditioning. For example, 2010 was a substantially higher collection year at Prosser than typical, resulting in high densities during reconditioning (Hatch et al. 2013b), Winthrop fish were not given effective treatment for copepods in 2012, and there

were issues with fish care on the Snake River project in 2014, 2016, and 2017 (Hatch et al. 2015, 2017, and 2018). The pattern of low consecutive maturation rates in these years indicates that consecutive maturation rate is sensitive to husbandry conditions. However, there were no issues with fish care at Winthrop or in the Snake River in 2020, and yet consecutive maturation rates decreased at both sites. This can be attributed to the effects of pre-capture environmental conditions, and the collection of fish in poorer condition may have also played a role. The variation in consecutive maturation rate from year to year means that projects either need to be able to accommodate varying numbers of skip spawners, or alternative strategies such as transporting and releasing skip spawners need to be explored.

Skip maturation rates in most CRB kelt reconditioning projects have been uniformly high, ranging from 80 to 100%. Surprisingly, however, skip maturation rates at Prosser, the most well established and longest running of the reconditioning projects, have generally been lower than in the other projects. The reasons for this difference are not known, but it may relate to water quality during the winter at the different locations. In other projects, maturation rates of skip spawners have been high even in years with low maturation rates of consecutive spawners, such as 2014 and 2016 in the Snake River project or 2017 in the Winthrop project. These results indicate that most kelts that are not rematuring after one summer of reconditioning will mature as skip spawners the next year. In addition, skip spawners have larger eggs and are more fecund than maiden or consecutive spawners, and have greater energy reserves at release (Jenkins, et al. 2018). Thus, skip spawners provide a source of steelhead spawners to seed habitat in years when the numbers of maiden spawners or survival and consecutive maturation rates of reconditioned kelts are low. Given the critically low number of steelhead spawners throughout the CRB in recent years and the species conservation status, this is a resource worth supporting.

Reproductive development in wild kelt steelhead 2021

Introduction

An understanding of the reproductive status of female kelt steelhead during reconditioning and at release is required to maximize the success of Columbia River Basin kelt reconditioning projects. Natural steelhead production is limited by the number of female spawners. In order to contribute to ESA-listed steelhead populations, female kelts must not only survive reconditioning but also remature and produce viable eggs. Questions regarding reproductive performance of reconditioned fish underlie issues raised regarding kelt reconditioning projects during ISRP review (ISRP 2011). We believe these issues can be best addressed by research aimed at an improved understanding the life history and physiology of post-spawning steelhead.

Iteroparous female salmonids have two major post-spawning life history trajectories (Chaput and Jones 2006; Keefer et al. 2008; Rideout et al. 2005; Rideout and Tomkiewicz 2011). After a spawning event, some fish are able to restore energy lost during migration and spawning, redevelop a mature ovary, and spawn the next year. These fish are termed consecutive spawners. Other fish do not initiate redevelopment of the ovary for the next spawning season, but instead skip a year. These fish are termed skip spawners. We hypothesize that these life history trajectories are the result of the effect of energy balance on maturation decisions made during seasonally defined critical periods. The influential critical period model of the first reproductive maturation (puberty) in salmonids posits that maturation is initiated during a decision window approximately one year prior to spawning (Campbell et al. 2006; Satterthwaite et al. 2009; Shearer and Swanson 2000; Thorpe 2007). This decision is made based on energy reserves. We hypothesize that a similar decision mechanism regulates rematuration in post-spawning steelhead. Consistent with this idea, we found that energy restriction affected reproductive development within 10 weeks after spawning in female rainbow trout (Caldwell et al. 2013; Caldwell et al. 2014). In post-spawning fish, energy driven decisions take place in the context of the extreme energy deficit incurred by migration and spawning (Penney and Moffitt 2014a; Penney and Moffitt 2014b; Penney and Moffitt 2015). Threshold energy levels for maturation or rematuration are determined by the genetic makeup of the fish and subject to selection (Carlson and Seamons 2008; Hutchings 2011).

Studies conducted in 2009-2011 established that blood levels of estradiol and vitellogenin diverge between rematuring and non-rematuring fish during reconditioning. Estradiol is the principal female gonadal steroid in fishes, which regulates many aspects of reproductive development, and vitellogenin is a phospholipoprotein produced by the liver under regulation by estradiol which provides most of the material for ovarian development. Estradiol indicates maturation earlier than vitellogenin, and the cost of the estradiol assay is about 1/4th of the cost of the vitellogenin assay.

During 2021, we measured estradiol level in a large number of blood samples. We collected blood from fish in the reconditioning programs at Dworshak National Fish Hatchery (DNFH, Asahka, ID) and Prosser Hatchery (Prosser, WA), ran plasma estradiol assays, and provided maturation status to project managers so that rematuring fish could be released and non-rematuring fish retained for further reconditioning (at DNFH) or below Bonneville Dam (for immature fish that were held at the Prosser hatchery). Additionally, we collaborated with colleagues in the Upper Columbia reconditioning project at Winthrop National Fish Hatchery (WNFH, Winthrop, WA) to measure estradiol levels in samples they collected from their reconditioned kelts.

Methods

Fish Collection and Husbandry

Steelhead kelts were collected and reconditioned at Prosser Hatchery, Washington, Dworshak National Fish Hatchery (DNFH), Idaho, and Winthrop National Fish Hatchery (WNFH), Washington as described elsewhere (Abrahamse and Murdoch 2013; Abrahamse and Murdoch 2014; Trammell et al. 2016).

Sampling

Fish were blood sampled on the indicated dates (Table 3.1, 3.2, and 3.3). During blood sampling, blood (2 mL) was drawn from the caudal vein using heparinized syringes (ammonium heparin, 10 mg/mL) and centrifuged (5 min, 5000 g). Plasma was collected and frozen on dry ice in the field prior to storage at -80°C. In addition to blood sampling, the length, weight and sex of fish were recorded, and a reading of muscle lipid levels was taken with a Distell Fish Fatmeter (Distell Inc., West Lothian, Scotland), using the rainbow trout muscle lipid setting (Trout-1) at the two most anterior measurement sites recommended by the manufacturer (Colt and Shearer 2001; Crossin and Hinch 2005).

Table 3. 1. Wild steelhead kelts held for long-term reconditioning for the 2021 reconditioning year. Prosser: Prosser Hatchery, DNFH: Dworshak National Fish Hatchery, WNFH: Winthrop National Fish Hatchery.

Location	Fish type	Total # Fish	Collected in 2021	Collected in 2020	Notes
Prosser	Wild kelts	360	360	N/A	
DNFH	Wild kelts	122	56	66	Very low collection rate for 2021
WNFH	Wild kelts	81	64	17	

Table 3. 2. Wild steelhead kelts sampled during the fall of 2021. Prosser: Prosser Hatchery, DNFH: Dworshak National Fish Hatchery, WNFH: Winthrop National Fish Hatchery.

Location	Sample date	Fish type	# Fish	Notes
Prosser	9/9/21	Wild kelts	242	All fish were collected in 2021
DNFH	9/23/21	Wild kelts	84	Fish were collected in 2020 (n = 62) and 2021 (n = 22)
WNFH	9/21/21	Wild kelts	22	Fish were collected in 2020 (n = 10) and 2021 (n = 12)

Table 3. 3. Wild steelhead reconditioned kelts released during the fall of 2021. Prosser: Prosser Hatchery, DNFH: Dworshak National Fish Hatchery, WNFH: Winthrop National Fish Hatchery.

Location	Release date	Fish type	# Fish	Notes
Prosser	10/28/21	Wild kelts	202	All surviving fish collected in 2021
DNFH	11/10/21	Wild kelts	65	Fish were collected in 2020 (n = 61) and 2021 (n = 4)
WNFH	11/9 and 11/12/2021	Wild kelts	14	Fish were collected in 2020 (n = 8) and 2021 (n = 6), 9 released in the Methow River and 5 in the Columbia River

Estradiol Assay

Fish plasma level of estradiol-17 β (E2) is an indicator of reproductive development. Fish plasma samples must be solvent extracted prior to E2 assay to remove interfering substances. Plasma samples (250 μ L) were extracted twice consecutively in 10 mL glass tubes with anhydrous diethyl ether (JT Baker, Avantor Performance Materials, Inc.; Center Valley, PA, USA). 2.0 mL diethyl ether was added to each tube and samples were vortexed for 1 m, the phases allowed to separate for 7 minutes, and then frozen on dry ice. After 5 minutes, the aqueous phase was inspected to ensure that it was frozen solid, and the solvent fraction was then poured off into a 5 mL glass tube. Diethyl ether extracts were then placed in a 54°C water bath (OA-SYS™ Heating System; Organomation Associates, Inc; Berlin, MA) and dried down under a gentle stream of N₂ directed *via* a nitrogen evaporator manifold (N-EVAP™ 112; Organomation Associates, Inc; Berlin, MA). A second extraction of the remaining aqueous fraction from each plasma sample was then performed, again using 2.0 mL diethyl ether, as described above; this second extract was pooled with the first extract. Dried extracts of fish plasma were resuspended in 250 μ L assay buffer from the estradiol assay kit.

Plasma E2 concentrations were assayed by an enzyme immunoassay using an acetylcholinesterase linked estradiol tracer (Cayman Chemical, Ann Arbor, MI). Extracted

plasma samples were appropriately diluted and duplicate technical replicates assayed in the EIA according to the manufacturer's instruction manual provided with the kit.

Results

Plasma E2 levels were bimodally distributed in blood samples taken from female kelts in all projects at a pre-release sampling in the fall (Figs 3.1, 3.2, 3.3). As found in previous years, the division between the lower and higher modes was approximately 1000 pg/mL E2 at all locations. A few fish with E2 levels close to 1000 pg/mL could group with either mode. These fish could be maturing more slowly than the rest of the upper mode. Consequently, the division between modes was adjusted to include these fish as rematuring so that the fish could be released. As is typical, the rematuration rate of female kelts as consecutive spawners in 2021 was highest at Prosser - females rematured at a rate of 72.9%. Consecutive spawners from other programs on the Snake River and Upper Columbia River had lower rates of rematuration for 2021, with only 18.2% of the Snake River fish rematuring and 50% of the Upper Columbia River fish rematuring. As with previous years, the rematuration rate of female kelts held for a second year of reconditioning was higher than consecutive spawners for the Snake River fish (98.4%) as well as the Upper Columbia River fish (88.9%).

Snake River 20/21 Kelts

Overall Maturing: 77.1%

2020 Maturing: 98.4%

2021 Maturing: 18.2%




Figure 3. 1. Plasma estradiol (E2) levels in wild female Snake River kelts sampled in fall of 2021.

Prosser 2021 Kelts

Maturing: 72.5%

Borderline: 0.4%

Non-maturing: 27.1%




Figure 3. 2. Plasma estradiol (E2) levels in wild female Yakima River kelts sampled in fall of 2021.

Winthrop 20/21 Kelts

Overall maturing: 66.7%
2020 Kelts Maturing: 88.9%
2021 Kelts Maturing: 50%




Figure 3. 3. Plasma estradiol (E2) levels in wild female Upper Columbia River kelts sampled in fall of 2021.

Due to the relatively low survival rate for hold over fish at Prosser hatchery (averaging approximately 13.7%), all fish are now being released regardless of maturation status. Thus, there are no skip spawners at Prosser. Fish believed to be maturing were released just downstream of Prosser Dam while immature fish were trucked downstream to below Bonneville Dam. Skip spawner survival for fish collected in 2020 and held for further reconditioning was on par or better compared with previous years: 92.4% at DNFH and 41.2% at WNFH. Survival of fish collected in 2021 to release was 56.1% at Prosser, 39.3% at DNFH, and 18.8% at WNFH. Of note, there was significantly higher than average mortality at DNFH and WNFH during the summer of 2021.

Discussion

It is now well established that some female steelhead kelts remature after a summer of reconditioning, whereas other fish do not, and that plasma estradiol level from mid-June onward indicates maturation status (Jenkins et al. 2018). Evidence in both steelhead kelts and

post-spawning rainbow trout suggests that the initial decision to remature is made early, before mid-July for kelts and during the 10 weeks after spawning in rainbow trout (Bromage et al. 1992; Caldwell et al. 2013; Caldwell et al. 2014; Hatch et al. 2013; Jenkins et al. 2018). Plasma estradiol levels in rematuring and non-rematuring kelts for 2021 at all sites were similar to previous years and were similar to those seen in other projects.

Female consecutive maturation rates were variable among the projects this season. It is possible that this relates to pre-capture environmental conditions. In previous years, the relatively low consecutive maturation rates found in Snake River kelts has been in line with what has been observed previously in Snake River steelhead, and steelhead from the Skeena and Nass systems in British Columbia, which have a life history similar to Snake River B-run steelhead. These cohorts have been found to repeat spawn predominantly as skip spawners (Chudyk 1976; Keefer et al. 2008; Moore et al. 1995). This has been hypothesized to be due to the longer migration and later spawn timing of these fish. With only 30% of fish rematuring after a year of reconditioning (on average), consecutive rematuration is observed in less than half of this population, implying that pre-capture environmental conditions may dictate the reproductive strategy employed. This could be the result of the warmer water temperatures the Columbia River Basin has been experiencing the past couple of summers, requiring a longer recovery period before the kelts are able to mature again (even with reconditioning). This is supported by the consistently high rates of maturation in the fish held for a second year of reconditioning. This year's higher mortality at WNFH and DNFH, coupled with the lower-than-average rate of consecutive rematuration, provides support for the theory that pre-capture environmental conditions are an important factor when evaluating what contributes to rematuration. Should we observe a high rate of maturation in the skip spawners next year, it will provide further evidence that the reconditioning program is providing an (artificial) refuge for life histories that would otherwise be in decline, which is especially important during years with low adult returns.

Non-rematuring fish collected from both the Snake and Upper Columbia rivers held for a second year rematured at very high rates (88.9% or higher) in 2021. This adds to a growing body of data showing that non-rematuring females will remature as skip spawners if held for a second year. Skip spawning is a natural life history in Columbia Basin steelhead. Increased size, fecundity, and energy reserves in skip spawners would be expected to result in greater relative reproductive success versus maidens or consecutive repeat spawners (Jenkins et al. 2018). The presence of skip spawners increases life history diversity, which would be expected to increase population stability in steelhead populations (Moore et al. 2014; Schindler et al. 2010). Moreover, whether and how much culture conditions can influence the proportion of consecutive and skip spawning kelts in captive reconditioning is not well understood. These considerations suggest that Columbia Basin kelt reconditioning programs should find ways to accommodate the skip spawner life history, a fact that's becoming increasingly important as run numbers show an overall decrease and the site's survival rate increases. Together, these data point out the important role the program is playing in keeping the local population healthy and diverse.

Hold over survival in Snake River fish is relatively high (typically over 60%), which contrasts with the low survival experienced by fish held for an additional year at Prosser. The difference could be due to population-specific life history differences but could also be due to over winter water quality. The Prosser facility is located in an area that experiences heavy runoff. Considering that the low hold over survival has occurred since the site began holding non-maturing fish in 2016, the suitability of Prosser as a hold over reconditioning facility was re-evaluated. Considering the continued low returns, it was decided that all fish would once again be returned to the river, regardless of maturation status.

Typically, there is high mortality in recently collected fish; however, both DNFH and WNFH experienced higher mortality than is typically observed. Survival of recently collected fish at both locations tends to average around 60%. The trend for decreased survival is most likely the result of the poor ocean conditions and elevated river water temperatures (i.e., climate change) during their return migration. Compounding the issue is that females appear to be more affected by the worsening conditions (Hatch et al. 2013; Keefer et al. 2008), making the reconditioning program more likely to be negatively impacted by this effect. To counteract the effects of climate change and declining return numbers, it seems prudent to increase the number of collection sites, and to investigate ways in which collection efforts could be improved (e.g., better holding conditions, improved sorting and transportation that decrease stress levels, discussing how changes to spill patterns could help collection numbers).

Chapter 3. Building a Snake River Kelt Reconditioning Facility

2021 Progress

Significant progress was made toward the realization of a dedicated kelt reconditioning facility for the Snake River in 2021. Efforts included: the Facility Review team reviewing 10%, 60%, and 90% design documents. Currently, the team is reviewing the 90% design documents generated by R2 Resource Consultants, Inc. Work developing a monitoring and evaluation budget for the new facility has continued.

We anticipate that Final Design Documents will be completed in 2022 and we will begin steps toward construction of the Snake River Kelt Reconditioning Facility.

Background

In the Columbia Basin Fish Accord Agreement that CRITFC is party to, \$2M was included for capital construction of a Snake River Kelt Reconditioning Facility. The Northwest Power and Conservation Council (NWPCC) three-step review process is triggered for any artificial production initiative that involves the construction of new production facilities. In 2016, we drafted a [Master Plan](#), reviewed the plan with co-managers and action agencies and submitted it the NWPCC for review by the Independent Science Review Panel (ISRP). In December 2016, the NWPCC accepted our Master Plan and recommended that we proceed to final design of the facility. This Master Plan would result in the fabrication of new facilities at an existing propagation facility.

The initial review of the Master Plan by the ISRP was completed in May 2016. The ISRP response is summary was:

“The Master Plan is well written and contains an excellent summary of the extensive steelhead reconditioning work that has occurred in the Basin. Moreover, we compliment the proponents for investigating and addressing the many difficulties associated with steelhead reconditioning. Numerous challenges associated with fish culture had to be addressed, including establishing appropriate holding and rearing environments, formulating diets, and developing disease control protocols. The effects of long-term reconditioning on gamete viability, fidelity to natal streams, and ability to reproduce in nature were investigated. Comparisons that evaluated the potential benefits of various kelt treatments that ranged from simple direct transportation past downstream dams to long-term reconditioning lasting from 6 to 20 months were also conducted. In general, the results of these assessments indicated that long-term reconditioning of kelts appears to be a promising approach that might lead to a viable conservation strategy for steelhead.

The proponents acknowledge that the submitted Master Plan does not yet have all the necessary components for a Step 1 review. It currently lacks a Hatchery Genetic Management Plan (HGMP), and work is needed on the program's Research, Monitoring and Evaluation Plan and Comprehensive Environmental Assessment. Before producing these elements of the Master Plan, the proponents requested that the ISRP determine if the program's preferred location for a long-term reconditioning facility, for Snake River B-run steelhead, is appropriate.

More information is needed before a decision about the location of the proposed long-term reconditioning facility can be reached. Specifically, information on the following issues is requested in the updated Step 1 Master Plan. Additional comments provided in the ISRP's full report should also be considered in the revision.

The biological and ecological rationale for annually increasing B-run steelhead escapement by 180 reconditioned female kelts needs to be explained in the Master Plan. Clarification on why male kelts are not included in the proposed reconditioning program is needed.

The biological escapement goals for B-run steelhead populations in the Snake River subbasin should be in the Master Plan along with a description of what project "success" entails. To what extent, for example, are reconditioned kelts expected to contribute to the rebuilding of natural steelhead populations and eventually to fisheries?

If available, information on the abundance and status and trends of B-run steelhead populations in the Clearwater and Salmon River subbasins should be provided in the Master Plan. Current spawning levels of B-run steelhead in the Snake River Basin should also be described with reference to numerical objectives for natural spawning steelhead. Additionally, a brief overview of the factors limiting each of these populations should be added to the Plan. Substantial hatchery and habitat restoration actions affecting B-run steelhead are occurring in the Snake River subbasin. The Master Plan should briefly describe these programs and indicate how the proponent's goal of annually releasing 180 reconditioned kelts will be coordinated with ongoing habitat restoration and existing hatchery programs.

As it is currently designed, the kelt reconditioning program will recondition female B-run steelhead kelts without targeting specific populations. It would seem that capturing, reconditioning, and releasing kelts from populations that have the potential to accommodate additional spawners would be a more efficient and productive way of directing this strategy. The Master Plan should explain why a more focused program was not considered. The Master Plan should discuss the infrastructural needs of a more focused and integrated reconditioning program. If the project, for instance, were to narrow its focus on B-run populations that could benefit from the addition of reconditioned kelts, would facilities at Dworshak National Fish Hatchery be adequate to meet these new escapement objectives? The Master Plan should compare the benefits and drawbacks of increasing B-run steelhead escapements by modifying harvest regulations, by long-term reconditioning for adult release, and long-term reconditioning for captive breeding and smolt release.

Some discussion of the genetic risks that may accompany reconditioning (e.g., heritable epigenetic effects and domestication selection) needs to be added to the Master Plan or incorporated into the Plan's HGMP."

We revised the [Master Plan](#) and submitted the document to the ISRP in July 2016 and received "meets scientific review criteria (qualified)" recommendation on September 27, 2016.

At the November 2016 NWPCC meeting in Coeur d' Alene, we presented our [Master Plan to the Council's Fish Committee](#). The Fish Committee received the plan favorably and recommended that it be presented to the full Council in December. At the December Council meeting we again presented the [Master Plan](#) and received a recommendation from the Council to proceed to the Final Design stage of the 3-step process.

In 2017, advancements were made in drafting a Monitoring and Evaluation (M&E) Plan and environmental compliance documents. Also, in 2017, we met with BPA and determined that BPA would solicit through a Request for Proposals (RFP) for a firm to design and build the kelt facility. The pace of this action has been slow and there are several components that must be completed. These include a Memorandum of Understanding (MOU) for construction, Operation and Maintenance (O&M) funding plans, Facility Designs, completion of the Northwest Power and Planning Council's Step 3, and construction of the facility.

In 2019, discussions between CRITFC and BPA continued regarding soliciting for a design and construction firm and development of a M&E and O&M plan. Bonneville Power Administration solicited proposals for the design phase in November 2019. A review group was formed to select a design firm and received completed designs at the end of 2020. If the current schedule holds, construction will be completed in 2022 and facility use will begin in 2023.

Chapter 4. Project wrap up, delineation of necessary follow up monitoring and research, complete manuscripts, transition to other locations or uses of kelt reconditioning technology.

The ISRP has identified 3 major points that this project should focus on before moving towards full-scale basin wide production:

1. Focus on identifying at-risk populations for collection utilizing genetic tools to focus reconditioning efforts on these populations specifically.
2. Transferring kelt reconditioning knowledge and techniques to other at-risk populations in the Columbia River Basin to further assist in recovery efforts.
3. Development of a plan to provide the documented kelt technological knowledge to other specific agencies for implementation. We intend to have this component completed by the 2022 Annual Report.

Due to declining returns, staff from The Nez Perce Tribe analyzed Snake River Basin spring/summer Chinook and steelhead population extinction risk and presented their results to the Fish and Wildlife Committee at the [May Committee meeting](#). The NPT staff finds that the results of their analysis are troubling, and the status of these populations suggest that approximately 50% of the Snake River steelhead populations will reach quasi-extinction levels by 2025. A similar call to action occurred when the fish were first listed under the ESA which spurred captive broodstock programs, gene-banking, and other required emergency actions. Steelhead kelt reconditioning is a management tool that could reduce the chances of extirpation for these identified steelhead populations and should be considered basinwide where appropriate.

Due to these dire straits facing steelhead populations, prioritizing certain populations for recovery actions may be somewhat misleading, because all steelhead populations are in demographic trouble, so any recovery action should be beneficial to any population that these actions are applied to. The NPT and CRITFC have set a prioritization table based on populations that are the most imperiled, while also recognizing which populations we can most successfully collect for reconditioning in the Snake River basin, please see Table 5.1. This table contributes towards addressing point 1 from the ISRP.

Table 5. 1. Steelhead kelt collection sites prioritized by high QET risk and kelt collection potential.																
MPG	Location	Priority Ranking 1 or 2	Population	Endemic Hatchery Stock Yes or No	High QET Risk Yes or No	Kelt Personnel Required for Collection	Holding Type	Holding Location	Holding Capacity	Holding Duration (Hours)	Time to DNFH (Hours)	Estimated Cost for 2022 (x\$1,000)	Estimated Number of Kelts	Agencies Responsible	Comments	Location
Salmon River	Pahsimeroi River Weir	1	Pahsimeroi	No	Yes	Yes	on site transport tank	direct transport	20	6	6.5	12-16	20	IDFG		Pahsimeroi River Weir
Clearwater River	Lolo Creek Weir	1	Lolo Pop	No	Yes	Yes	on site transport tank	direct transport	20	6	1	>100	5	NPT	very few adults captured	Lolo Creek Weir
Salmon River	Rapid River Weir	1	Little Salmon	No	Yes	Yes	on site transport tank	direct transport	20	6	3	7-10	5	IDFG	very few adults observed	Rapid River Weir
Salmon River	Lemhi River Weir	1	Lemhi	No	Yes	Yes	on site transport tank	direct transport	20	6	6.5	>100	5	IDFG	very few adults captured	Lemhi River Weir
Salmon River	Yankee Fork (Flat Pole) Weir	1	Salmon Upper Mainstem	No	Yes	Yes	on site transport tank	direct transport	20	6	8	>100	5	IDFG		Yankee Fork (Flat Pole) Weir
Salmon River	Upper Salmon River (Sawtooth) Weir	1	Salmon Upper Mainstem	No	Yes	Yes	on site transport tank	direct transport	20	6	8	12-16	5	IDFG		Upper Salmon River (Sawtooth) Weir
Salmon River	Morgan Creek Weir	2	Pahsimeroi	No	Yes	Yes	on site transport tank	direct transport	20	6	7	>100	5	IDFG	very few adults captured	Morgan Creek Weir
Salmon River	Iron Creek Weir	2	Pahsimeroi	No	Yes	Yes	on site transport tank	direct transport	20	6	6.5	>100	5	IDFG	very few adults captured	Iron Creek Weir
Salmon River	Poison Creek Weir	2	Pahsimeroi	No	Yes	Yes	on site transport tank	direct transport	20	6	6.5	>100	5	IDFG	very few adults captured	Poison Creek Weir

Salmon River	Twelve Mile Creek Weir	2	Pahsimeroi	No	Yes	Yes	on site transport tank	direct transport	20	6	6	>100	5	IDFG	very few adults captured	Twelve Mile Creek Weir
Salmon River	Tower Creek Weir	2	Lemhi	No	Yes	Yes	on site transport tank	direct transport	20	6	5.5	>100	5	IDFG	very few adults captured	Tower Creek Weir
Salmon River	Carmen Creek Weir	2	Lemhi	No	Yes	Yes	on site transport tank	direct transport	20	6	5.5	>100	5	IDFG	very few adults captured	Carmen Creek Weir
Salmon River	Fourth of July Creek Weir	2	Salmon Upper Mainstem	No	Yes	Yes	on site transport tank	direct transport	20	6	5.5	>100	5	IDFG	very few adults captured	Fourth of July Creek Weir
Grande Ronde River	Cottonwood Creek Weir	2	Grande Ronde Lower Mainstem	No	Unk	Yes	on site transport tank	direct transport	20	6	2	5-7	5	WDFW	very few adults captured	Cottonwood Creek Weir
Salmon River	Challis Creek Weir	2	East Fork Salmon	No	Unk	Yes	on site transport tank	direct transport	20	6	7	>100	5	IDFG	very few adults captured	Challis Creek Weir
Lower Snake River	Tucannon River Weir	1	Tucannon	Yes	No	Yes	on site transport tank	direct transport	20	6	2	5-7	10	WDFW		Tucannon River Weir
Lower Snake River	Alpowa Creek Weir	2	Asotin	No	No	Yes	on site transport tank	direct transport	20	6	1.5	5-7	5	WDFW	very few kelts captured	Alpowa Creek Weir
Lower Snake River	Asotin Creek Weir	1	Asotin	No	No	Yes	on site transport tank	direct transport	20	6	1.5	5-7	40	WDFW		Asotin Creek Weir
Lower Snake River	George Creek (Asotin Creek) Weir	2	Asotin	No	No	Yes	on site transport tank	direct transport	20	6	1.5	5-7	20	WDFW		George Creek (Asotin Creek) Weir
Lower Snake River	Tenmile Creek Weir	2	Asotin	No	No	Yes	on site transport tank	direct transport	20	6	1.5	5-7	10	WDFW		Tenmile Creek Weir
Clearwater River	Big Bear Creek Weir	1	Lower Clearwater	No	No	Yes	on site transport tank	direct transport	20	6	1	>100	10	IDFG		Big Bear Creek Weir
Clearwater River	East Fork Potlatch River Weir	1	Lower Clearwater	No	No	Yes	on site transport tank	direct transport	20	6	1.5	>100	5	IDFG	very few kelts observed	East Fork Potlatch River Weir

Clearwater River	Jacks Creek Weir	2	Lower Clearwater	No	No	Yes	on site transport tank	direct transport	20	6	0.5	<5	5	NPT		Jacks Creek Weir
Clearwater River	Clear Creek Weir (Kooskia NFH)	1	Lower Clearwater	No	No	Yes	adult trap holding pond	Kooskia NFH	100	48	1	5-7	5	NPT	very few kelts observed	Clear Creek Weir (Kooskia NFH)
Clearwater River	Fish Creek Weir	1	Lochsa	No	No	Yes	on site transport tank	direct transport	20	6	2	5-7	10	IDFG	Current IDFG policy limitations	Fish Creek Weir
Clearwater River	Crooked River Weir	2	South Fork Clearwater	No	No	Yes	on site transport tank	direct transport	20	6	2	5-7	5	IDFG	very few adults captured	Crooked River Weir
Clearwater River	Mill Creek (SF Clearwater River)	2	South Fork Clearwater	No	No	Yes	on site transport tank	direct transport	20	6	1	>100	5	NPT		Mill Creek (SF Clearwater River)
Clearwater River	Newsome Creek Weir	2	South Fork Clearwater	No	No	Yes	on site transport tank	direct transport	20	6	1	>100	0	NPT	location above SF Clearwater River velocity barrier	Newsome Creek Weir
Grande Ronde River	Joseph Creek Weir	1	Joseph	No	No	Yes	on site transport tank	direct transport	20	6	2	>100	40	NPT		Joseph Creek Weir
Grande Ronde River	Lostine River Weir	1	Wallowa	No	No	Yes	on site transport tank	direct transport	20	6	3	7-10	20	NPT		Lostine River Weir
Grande Ronde River	Lookingglass Hatchery Ladder	1	Grande Ronde Upper Mainstem	No	No	Yes	on site transport tank	direct transport	20	6	4	7-10	5	ODFW	potential for multi-day holding	Lookingglass Hatchery Ladder
Grande Ronde River	Wallowa Hatchery Ladder	2	Wallowa	No	No	Yes	on site transport tank	direct transport	20	6	3	7-10	0	ODFW	No fish passed above weir	Wallowa Hatchery Ladder

Grande Ronde River	Upper Grande Ronde River Weir	2	Grande Ronde Upper Mainstem	No	No	Yes	on site transport tank	direct transport	20	6	4.5	7-10	5	ODFW		Upper Grande Ronde River Weir
Grande Ronde River	Catherine Creek Weir	2	Grande Ronde Upper Mainstem	No	No	Yes	on site transport tank	direct transport	20	6	4.5	7-10	5	ODFW		Catherine Creek Weir
Grande Ronde River	Big Canyon - Deer Creek Weir	1	Wallowa	No	No	Yes	on site transport tank	direct transport	20	6	3.5	7-10	10	ODFW		Big Canyon - Deer Creek Weir
Imnaha River	Screw trap at Cow Creek	1	Imnaha	Yes	No	Yes	large aluminum tank	Thorn Creek Facility	20	48	5	20-30	5	NPT	5-hour direct transport without Thorn Creek holding	Screw trap at Cow Creek
Imnaha River	Imnaha River Weir at Thorn Creek	2	Imnaha	Yes	No	Yes	large aluminum tank	Thorn Creek Facility	20	48	5	20-30	10	NPT	5-hour direct transport without Thorn Creek holding	Imnaha River Weir at Thorn Creek
Imnaha River	Horse Creek Weir	2	Imnaha	Yes	No	Yes	large aluminum tank	Thorn Creek Facility	20	48	5	>100	10	NPT	5-hour direct transport without Thorn Creek holding	Horse Creek Weir
Imnaha River	Little Sheep Creek Weir	2	Imnaha	Yes	No	Yes	on site transport tank	direct transport	20	48	4	7-10	10	ODFW	1.5-hours to transfer to Gumboot Facility	Little Sheep Creek Weir
Imnaha River	Camp Creek Weir	2	Imnaha	Yes	No	Yes	on site transport tank	direct transport	20	6	4	7-10	10	NPT	1.5-hours to transfer to Gumboot Facility	Camp Creek Weir
Imnaha River	Grouse Creek Weir	2	Imnaha	Yes	No	Yes	on site transport tank	direct transport	20	6	5	>100	10	NPT	1-hour to transfer to Gumboot Facility	Grouse Creek Weir

Imnaha River	Crazy Man Creek Weir	2	Imnaha	Yes	No	Yes	acclimation raceway	Gumboot Facility	100	48	4.5	15-20	5	NPT	5-hour direct transport without Gumboot Facility holding	Crazy Man Creek Weir
Imnaha River	Mahogeny Creek Weir	2	Imnaha	Yes	No	Yes	acclimation raceway	Gumboot Facility	100	48	4.5	>100	5	NPT	5-hour direct transport without Gumboot Facility holding	Mahogeny Creek Weir
Imnaha River	Imnaha River Weir at Gumboot Creek	1	Imnaha	Yes	No	Yes	acclimation raceway	Gumboot Facility	100	48	4.5	15-20	10	ODFW	5-hour direct transport without Gumboot Facility holding	Imnaha River Weir at Gumboot Creek
Imnaha River	Gumboot Creek Weir	1	Imnaha	Yes	No	Yes	acclimation raceway	Gumboot Facility	100	48	4.5	>100	10	NPT	0.5-hour to transfer to Gumboot Facility	Gumboot Creek Weir
Salmon River	Johnson Creek Weir	1	South Fork Salmon	No	No	Yes	on site transport tank	direct transport	20	6	6	>100	5	NPT	very few adults captured	Johnson Creek Weir
Salmon River	East Fork Salmon River Weir	1	East Fork Salmon	No	No	Yes	on site transport tank	direct transport	20	6	8	12-16	20	IDFG		East Fork Salmon River Weir
Salmon River	S. Fork Salmon River Weir at Knox Creek	1	South Fork Salmon	No	No	Yes	on site transport tank	direct transport	20	6	4	>100	5	IDFG		S. Fork Salmon River Weir at Knox Creek
Lower Snake River	Lower Granite Dam	1	Mixed Populations	NA	NA	No	large steel tank	Lower Granite Dam JFF	200	48	2	0	50	COE		Lower Granite Dam
Lower Snake River	Little Goose Dam	1	Mixed Populations	NA	NA	No	large aluminum tank	Little Goose Dam JFF	100	48	2	0	20	COE		Little Goose Dam

Lower Snake River	Lower Monumental Dam	1	Mixed Populations	NA	NA	No	large aluminum tank	Lower Monumental Dam JFF	100	48	2.5	<5	20	COE	Alternative release location for reconditioned kelts collected here (Tucannon River stock)	Lower Monumental Dam
-------------------	----------------------	---	-------------------	----	----	----	---------------------	--------------------------	-----	----	-----	----	----	-----	--	----------------------

Prioritization of steelhead populations for recovery actions.

Kelt reconditioning has proven to be effective at recapturing Steelhead production through iteroparity that otherwise would have been lost, but reconditioning efforts have thus far been applied at a coarse spatial scale, i.e., at MPG and arguably at ESU levels, without a concerted effort to target specific populations. This approach minimized collection effort and cost during the proof-of-concept phase but focusing on the most at-risk populations would maximize the conservation value of the technique. Although it would be challenging to collect, recondition and release at the population level, there do exist population level productivity measurements that can be monitored for sensitivity to reconditioning efforts, and kelt collection could target populations deemed most at risk or at highest potential gain. Based on juvenile productivity and adult return data available in the [Coordinated Assessments](#), we propose to refine a potential collection strategy that is sensitive to the benefits of specific collection sites impacting populations that stand to gain the most from reconditioning.

Table 5.2 shows the number of years of natural origin spawning abundance data monitored for specific MPGs. Table 5.3 shows the juvenile data monitored at specific sites for each of the corresponding adult populations. Adult monitoring encompasses a broader scale of measurement than the juvenile monitoring. The reported juveniles account for a fraction of the total juveniles that would be associated with the adult population monitored, but the trend in the juvenile responses should provide a relative sense of the benefits associated with the trend in adults.

An in-depth analysis of spawner and juvenile abundance trends has the potential to reveal where there may be population-level tactical advantage to collecting and reconditioning kelts. We propose to look at the potential collection sites identified in Table (5.1), and reference these to the juvenile production patterns, looking for patterns in juvenile production, and potentially identifying production in relation to relative spawning abundances (depending on ability to associate with adults with juveniles).

Table 5. 2. The number of years of available natural origin steelhead adult abundance data reported at population level.

MPG	POPULATION NAME	LOCATION NAME	N
Clearwater	Clearwater River Lower Mainstem - summer Steelhead	Clearwater River Lower Mainstem	36
Clearwater River	Lochsa and Selway River summer Steelhead	Lochsa and Selway River (Superpopulation)	36
Clearwater River	South Fork Clearwater River and Lolo Creek summer Steelhead	South Fork Clearwater River and Lolo Creek (Superpopulation)	36
Grande Ronde	Grande Ronde River Upper Mainstem - summer Steelhead	Grande Ronde River Upper Mainstem	52

Grande Ronde	Joseph Creek - summer Steelhead	Joseph Creek	48
Salmon	Little Salmon River - summer Steelhead	Little Salmon River	36
Salmon River	Lower and upper Middle Fork Salmon River summer Steelhead	Lower and Upper Middle Fork Salmon River (Superpopulation)	36
Salmon River	South Fork Salmon River and Secesh River summer Steelhead	South Fork Salmon River and Secesh River (Superpopulation)	36
Salmon River	Upper Salmon River Panther Creek to Headwaters summer Steelhead	Upper Salmon River Panther Creek to Headwaters (Superpopulation)	36

Table 5. 3. The number of years of available steelhead juvenile abundance location monitoring data.

LOCATIONNAME	SMOLTEQLOCATION	N
Clearwater River Lower Mainstem	Big Bear Creek	7
Clearwater River Lower Mainstem	East Fork Potlatch River, rkm = 1	7
Lochsa River	Colt Killed Creek, rkm =1	16
Lochsa River	Crooked Fork Lochsa River, rkm = 3	16
Lochsa River	Fish Creek	4
Lochsa River	Fish Creek, rkm = 2	23
Lochsa River	Lochsa River	1
Lolo Creek	Lower Granite Dam	6
South Fork Clearwater River	American River, rkm = 2	9
South Fork Clearwater River	Crooked River, rkm = 1	14
South Fork Clearwater River	Lower Granite Dam	2
South Fork Clearwater River	Red River, rkm = 6	9
Imnaha River	Lower Granite Dam	24
Asotin Creek	Asotin Creek	31
Tucannon River	Tucannon River	24
Lemhi River	Hayden Creek, rkm = 1	15
Lemhi River	Lemhi River, rkm = 49	19
Lemhi River	Lemhi River, rkm=7	11
Little Salmon River	Rapid River, rkm = 6	13
Middle Fork Salmon River Lower Mainstem	Big Creek	14
Middle Fork Salmon River Upper Mainstem	Marsh Creek, rkm = 11	17
Middle Fork Salmon River Upper Mainstem	Marsh Creek, rkm = 8	11
North Fork Salmon River	North Fork Salmon River, rkm =1	6

Pahsimeroi River	Pahsimeroi River, rkm = 2	20
Salmon River Upper Mainstem	Salmon River, rkm= 617	20
Secesh River	Lower Granite Dam	5
South Fork Salmon River	South Fork Salmon River, rkm = 118	24
South Fork Salmon River	South Fork Salmon River, rkm = 63	6

Figure 5.1 shows the juvenile production trend using the logarithm of juveniles in year t divided by juveniles in year $t+1$, i.e., $\log(J(t+1)/J(t))$. There does not appear to be any persistent year over year trend (zero implies no change in log scale). Figure 5.2 shows the trend in absolute juvenile abundance. We will examine patterns in juvenile and adult abundance trends and identify if there are populations that stand out. These will be associated with the collection sites and ordered such that kelt collection and reconditioning selects for the populations that might benefit most. Figure 3 shows the lag-4 spawner per spawner trend ($\log(S(t+4)/S(t))$), which implies that spawners four years from a brood year represent a large portion of returns, and as such $S(t+4)/S(t)$ will be an approximation for the population growth (λ). Compared to the juvenile pattern, there seems to be a recent persistent decline, but overall, on a 5–10-year scale, there seems to be a cyclic pattern. Our goal is to review all these patterns and look for signals at the population level that might indicate preferential treatment. Ultimately, we envision these data along with other factors including but limited to: accessibility, distance from reconditioning facility, kelt collection infrastructure, cost / fish collected to inform a kelt collection plan.




Figure 5. 1. Juvenile year over year production. Yearly values show $\log(J(t+1)/J(t))$.




Figure 5. 2. Juvenile abundance trend.



Figure 5.3. Lag 4 spawners per spawner. Figure shows the 4-year lag spawner per spawner trend in log space.

Adaptive Management & Lessons Learned

1. Columbia River steelhead populations upstream of Bonneville Dam are listed under ESA and need novel recovery strategies.
2. There is a relatively large abundance of kelt steelhead in the Columbia River Basin even in the upper most areas.
3. In general, repeat spawning steelhead make up a very small proportion of the spawning run.
4. Increasing repeat spawners in steelhead populations can have many positive effects on populations including increasing; genetic diversity, lifetime fecundity, and fitness since genes are distributed across generations.
5. Long-term reconditioning kelt steelhead provides 5 to over 100 times more repeat spawners than leaving the fish in the river.
6. Physiology studies have provided us with a much better understanding of energetic and physiological status of kelts, improved our understanding of alternative life histories in post-spawning fish, and improved survival and health of reconditioned fish.

7. Blood hormone assays are useful to classify consecutive and skip spawner steelhead. Future work needs to focus on optimizing strategies for skip spawner contributions.
8. There appears to be a reduction in the B-run steelhead composition between the maiden and kelt stage, but the B-run composition of repeat spawners is similar to the kelt composition. Underlying biological and behavioral factors contributing to such discrepancies are not well understood but likely warrant further investigation of potential causes. With more data including escapement comparisons, it may be possible to refine the confidence in estimated rates of iteroparity among Reporting Group's (RG's).
9. Age appears to be less of a factor in rates of iteroparity than size. While the A-run life history was observed to be present among all reporting groups, so too were the B-run life history.
10. Despite the understanding in recent years that the B-run life history is relatively uncommon outside the middle and south forks of both the Clearwater River and Salmon River, our results suggest otherwise. In fact, age 2-ocean fish were dominant among all 10 reporting groups. This finding has implications for management of steelhead populations in the basin and provides evidence that regionally based classifications of life history types or their distributions warrants reconsideration.
11. The upper Salmon River region produces a disproportionate number of Snake River kelt steelhead and is presumably an important factor in spawner abundance for that region. This result is mirrored among hatchery-origin fish.
12. Adding a production level kelt reconditioning facility at Nez Perce Tribal Hatchery will make achieving the goal RPA 33 possible, i.e., increase the abundance on adult B-run steelhead by 6%.
13. The Snake River Kelt Reconditioning Facility Master Plan was submitted and favorably review by the ISRP and recommended to proceed to final design by the NWPCC in December of 2016.
14. Reproductive success studies are underway at a variety of scales: hatchery analog, spawning channel, and natural river. Results are positive.
15. Artificially reconditioned kelt steelhead appear to repeat home with high fidelity. Data indicates that natural repeat spawners in the Snake River exhibited a 15% stray rate.
16. Concluded with the Cle Elum spawning channel.
17. Kelt biophysiological decision to remature is made soon after spawning.
18. As a result of this project 1,700 kelt steelhead were collected in the Snake River since 2012 and 697 of those fish were reconditioned and released back into the Snake River.
19. As a result of this project 7,868 kelt steelhead were collected in the Yakima River since 2008 and 3,416 of those fish were reconditioned and released back into the Yakima River.
20. Kelt Reconditioning, during years of low steelhead returns, effectively acts as a stop gap or safety net measure that should allow for a larger production of the juvenile population than normal under poor return years. This increase in juvenile production, should rearing and migration conditions improve, would translate into additional adult returns later, thus decreasing the time period for recovery after poor run years.
21. GSI analysis revealed that >50% of the reconditioned kelts released in the Snake are from B-run MPGs (Clearwater and Salmon rivers). These fish are important for meeting the goal of RPA 33. The National Marine Fisheries Service has issued a new Biological Opinion for operation of the Columbia River Hydrosystem (NMFS 2019) and plan to issue supplanting BiOp for the FCRPS in 2020.
22. Conducted and produced valuable kelt research on rematuration of steelhead kelts and how environmental factors play into rematuration and how we may be able to better identify sequential/skip spawners to address management of steelhead kelts in the Columbia River Basin See Jenkins et al 2020 and Medeiros et al 2020.

23. New and improved redundancy systems and protocols are being put in place at Nez Perce Tribal Hatchery to prevent catastrophic loss at the facility until a dedicated facility is constructed, which will have better fail-safe systems in place.

References

- Abrahamse MS & Murdoch KG 2013 Upper-Columbia River Steelhead Kelt Reconditioning Project: 2012 Annual Report to the U.S. Dept. of Energy, Bonneville Power Administration, Project No. 2008-485-000. Portland, OR: Prepared by Yakama Nation Fisheries Resource Management.
- Abrahamse MS & Murdoch KG 2014 Upper-Columbia River Kelt Reconditioning Program Update. Portland, OR: Prepared by Yakama Nation Fisheries Resource Management.
- Ackerman, M. W., C. Habicht, and L. W. Seeb. 2011. Single-nucleotide polymorphisms (SNPs) under diversifying selection provide increased accuracy and precision in mixed-stock analyses of sockeye salmon from Copper River, Alaska. *Transactions of the American Fisheries Society* 140:865–881.
- Ackerman, M. W., J. McCane, C. A. Steele, M. R. Campbell, A. P. Matala, J. E. Hess, and S. R. Narum. 2012. Chinook and steelhead genotyping for genetic stock identification at Lower Granite Dam. 2011 Annual Report. Submitted to: U.S. Department of Energy, Bonneville Power Administration, Division of Fish and Wildlife, P.O. Box 3621 Portland, OR Contract #53239; Project #2010-026-00. Available at: <https://pisces.bpa.gov/release/documents/documentviewer.aspx?doc=P128035>
- Ackerman, M. W., J. McCane, C. A. Steele, N. V. Vu, M. R. Campbell, A. P. Matala, J. E. Hess, and S. R. Narum. 2014. Chinook and steelhead genotyping for genetic stock identification at Lower Granite Dam. 2013 Annual Report. Submitted to: U.S. Department of Energy, Bonneville Power Administration, Division of Fish and Wildlife, P.O. Box 3621 Portland, OR Contract #57343; Project #2010-026-00. Available at: <https://pisces.bpa.gov/release/documents/documentviewer.aspx?doc=P128035>
- Ali M, Nicieza A & Wootton RJ 2003 Compensatory growth in fishes: a response to growth depression. *Fish and Fisheries* 4 147-190.
- Anderson, E. C. 2010. Computational algorithms and user-friendly software for parentage-based tagging of Pacific salmonids. Final report submitted to the Pacific Salmon Commission's Chinook Technical Committee (US Section). 46 p.
- Baker DM, Davies B, Dickhoff WW & Swanson P 2000 Insulin-like growth factor I increases follicle-stimulating hormone (FSH) content and gonadotropin-releasing hormone-stimulated FSH release from coho salmon pituitary cells in vitro. *Biology of Reproduction* 63 865-871.
- Bar N 2014 Physiological and hormonal changes during prolonged starvation in fish. *Canadian Journal of Fisheries and Aquatic Sciences* 71 1447-1458.
- Barry TP, Marwah A & Nunez S 2010 Inhibition of cortisol metabolism by 17 alpha,20 beta-P: Mechanism mediating semelparity in salmon? *General and Comparative Endocrinology* 165 53-59.
- Barry TP, Unwin MJ, Malison JA & Quinn TP 2001 Free and total cortisol levels in semelparous and iteroparous chinook salmon. *Journal of Fish Biology* 59 1673-1676.

- Baumann G 2002 Growth hormone binding protein. The soluble growth hormone receptor. *Minerva Endocrinol* 27 265-276.
- Baxter, Colden, F. Richard Hauer, and William W. Woessner. 2003. "Measuring Groundwater–Stream Water Exchange: New Techniques for Installing Minipiezometers and Estimating Hydraulic Conductivity." *Transactions of the American Fisheries Society* 132 (3): 493–502. doi: 10.1577/1548-8659(2003)132<0493: MGWENT>2.0.CO; 2.
- Baxter, C. V., and R. F. Hauer. 2000. "Geomorphology, Hyporheic Exchange, and Selection of Spawning Habitat by Bull Trout (*Salvelinus confluentus*).” *Canadian Journal of Fisheries and Aquatic Sciences* 57: 1470–81.
- Beckman BR 2011 Perspectives on concordant and discordant relations between insulin-like growth factor 1 (IGF1) and growth in fishes. *General and Comparative Endocrinology* 170 233-252.
- Beckman BR, Larsen DA, Moriyama S, Lee-Pawlak B & Dickhoff WW 1998 Insulin-like growth factor-I and environmental modulation of growth during smoltification of spring chinook salmon (*Oncorhynchus tshawytscha*). *General and Comparative Endocrinology* 109 325-335.
- Bell, G. 1980. The costs of reproduction and their consequences. *The American Naturalist* 116(1):45-76.
- Benedet S, Andersson E, Mittelholzer C, Taranger GL & Bjornsson BT 2010 Pituitary and plasma growth hormone dynamics during sexual maturation of female Atlantic salmon. *General and Comparative Endocrinology* 167 77-85.
- Bergan-Roller HE & Sheridan MA. 2018. The growth hormone signaling system: Insights into coordinating the anabolic and catabolic actions of growth hormone. *General and Comparative Endocrinology* 258 119-133.
- Bhatta S, Iwai T, Miura C, Higuchi M, Shimizu-Yamaguchi S, Fukada H & Miura T 2012 Gonads directly regulate growth in teleosts. *Proceedings of the National Academy of Sciences of the United States of America* 109 11408-11412.
- Boe J, Montfort J, Nguyen T & Fostier A 2006 Identification of new participants in the rainbow trout (*Oncorhynchus mykiss*) oocyte maturation and ovulation processes using cDNA microarrays. In *Reprod Biol Endocrin*.
- Biga PR, Peterson BC, Schelling GT, Hardy RW, Cain KD, Overturf K & Ott TL 2005 Bovine growth hormone treatment increased IGF-I in circulation and induced the production of a specific immune response in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 246 437-445.
- Bjornsson BT 1997 The biology of salmon growth hormone: from daylight to dominance. *Fish Physiology and Biochemistry* 17 9-24.
- Bjornsson BT, Einarsdottir IE, Johansson M & Gong NP 2018 The Impact of Initial Energy Reserves on Growth Hormone Resistance and Plasma Growth Hormone-Binding Protein Levels in Rainbow Trout Under Feeding and Fasting Conditions. *Frontiers in Endocrinology* 9.
- Bjornsson BT, Johansson V, Benedet S, Einarsdottir IE, Hildahl J, Agustsson T & Jonsson E 2002 Growth hormone endocrinology of salmonids: regulatory mechanisms and mode of action. *Fish Physiology and Biochemistry* 27 227-242.

- Bjornsson BT, Taranger GL, Hansen T, Stefansson SO & Haux C 1994 The Interrelation between Photoperiod, Growth-Hormone, and Sexual-Maturation of Adult Atlantic Salmon (*Salmo-Salar*). *General and Comparative Endocrinology* 93 70-81.
- Blaise O, Weil C & Le Bail PY 1995 Role of IGF-I in the control of GH secretion in rainbow trout (*Oncorhynchus mykiss*). *Growth Regulation* 5 142-150.
- Bobé J, Montfort J, Nguyen T & Fostier A 2006 Identification of new participants in the rainbow trout (*Oncorhynchus mykiss*) oocyte maturation and ovulation processes using cDNA microarrays. In *Reprod Biol Endocrin.*
- Bon E, Corraze G, Kaushik S & LeMenn F 1997 Effects of accelerated photoperiod regimes on the reproductive cycle of the female rainbow trout: I—Seasonal variations of plasma lipids correlated with vitellogenesis. *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 118 183-190.
- Bonneville Power Administration and U.S. Army Corps of Engineers 2014. 2013 Snake River Kelt Management Plan. www.salmonrecovery.org. (April 2015).
- Bowersox, B. J., R. Banks and N. Davids. 2011. Potlatch River steelhead monitoring and evaluation project, Annual Report 2010. IDFG #12-103, April 2012.
- Branstetter R, Stephenson J, Pierce AL, Hatch DR, Bosch B, Fast D, Blodgett J, Everett SR, Paddlety J, Dasher R, et al. 2011 Steelhead Kelt Reconditioning and Reproductive Success. 2011 Annual Report to the U.S. Dept. of Energy, Bonneville Power Administration, Project No. 2007-401-000. Portland, OR: Prepared by the Columbia River Inter-Tribal Fish Commission.
- Branstetter R, Stephenson J, Pierce AL, Hatch DR, Bosch B, Fast D, Blodgett J, Everett SR, Paddlety J, Dasher R, et al. 2010 Steelhead Kelt Reconditioning and Reproductive Success. 2010 Annual Report to the U.S. Dept. of Energy, Bonneville Power Administration, Project No. 2010-401. Portland, OR: Prepared by the Columbia River Inter-Tribal Fish Commission.
- Breton B, Govoroun M & Mikolajczyk T 1998 GTH I and GTH II secretion profiles during the reproductive cycle in female rainbow trout: relationship with pituitary responsiveness to GnRH-A stimulation. *General and Comparative Endocrinology* 111 38-50.
- Brodersen J, Nilsson PA, Hansson LA, Skov C & Bronmark C 2008 Condition-dependent individual decision-making determines cyprinid partial migration. *Ecology* 89 1195-1200.
- Bromage N, Jones J, Randall C, Thrush M, Davies B, Springate J, Duston J & Barker G 1992 Broodstock management, fecundity, egg quality and the timing of egg-production in the rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 100 141-166.
- Buelow, J. L. 2011. Physiological Characteristics of Steelhead Kelt steelhead (*Oncorhynchus mykiss*) in the Snake River, Idaho. Master's thesis, University of Idaho, Moscow.
- Bull JJ & Shine R 1979 Iteroparous animals that skip opportunities for reproduction. *American Naturalist* 114 296-303.
- Burgner RL, Light JT, Margolis L, Okazaki T, Tautz A & Ito S 1992 Distribution and origins of steelhead trout (*Oncorhynchus mykiss*) in offshore waters of the North Pacific ocean. Vancouver, B.C.: International North Pacific Fisheries Commission.
- Busby, P. J., T. C. Wainwright, G. J. Bryant, L. J. Liehr, R. S. Waples, F. W. Waknitz and F. W. Lagomarsino. 1996. Status review of west coast steelhead from Washington, Idaho, Oregon, and California. U.S. Department of Commerce, NOAA Technical Memorandum

- NMFS-NWFSC-27. Available at:
<http://www.nwfsc.noaa.gov/publications/techmemos/tm27/tm27.htm>
- Burton MPM 1994 A critical period for nutritional control of early gametogenesis in female winter flounder, *Pleuronectes americanus* (Pisces, Teleostei). *Journal of Zoology* 233 405-415.
- Caisman JM 2015 Partial migration in steelhead (*Oncorhynchus mykiss*): Identifying factors that influence migratory behavior and population connectivity across a watershed. In *College of Graduate Studies*, p 80. Moscow, Idaho: University of Idaho.
- Caldwell LK, Pierce AL & Nagler JJ 2013a Metabolic endocrine factors involved in spawning recovery and rematuration of iteroparous rainbow trout (*Oncorhynchus mykiss*). *General and Comparative Endocrinology* 194 124-132.
- Caldwell LK, Pierce AL & Nagler JJ 2013b Metabolic endocrine factors involved in spawning recovery and rematuration of iteroparous female rainbow trout (*Oncorhynchus mykiss*). *Gen Comp Endocrinol* 194 124-132.
- Caldwell LK, Pierce AL, Riley LG, Duncan CA & Nagler JJ 2014 Plasma nesfatin-1 is not affected by long-term food restriction and does not predict rematuration among iteroparous female rainbow trout (*Oncorhynchus mykiss*). *PLoS One* 9 e85700.
- Campbell B, Beckman BR, Fairgrieve WT, Dickey JT & Swanson P 2006a Reproductive investment and growth history in female Coho salmon. *Transactions of the American Fisheries Society* 135 164-173.
- Campbell B, Dickey J, Beckman B, Young G, Pierce A, Fukada H & Swanson P 2006b Previtellogenic oocyte growth in salmon: relationships among body growth, plasma insulin-like growth factor-1, estradiol-17beta, follicle-stimulating hormone and expression of ovarian genes for insulin-like growth factors, steroidogenic-acute regulatory protein and receptors for gonadotropins, growth hormone, and somatolactin. *Biology of Reproduction* 75 34-44.
- Campbell, M. R., C. C. Kozfkay, T. Copeland, W. C. Schrader, M. W. Ackerman, and S. R. Narum. 2012. Estimating abundance and life history characteristics of threatened wild Snake River steelhead stocks by using genetic stock identification. *Transactions of the American Fisheries Society* 141:1310–1327.
- Campbell, NR, Harmon SA, Narum SR. 2015. Genotyping-in-Thousands by sequencing (GT-seq): a cost effective SNP genotyping method based on custom amplicon sequencing. *Molecular Ecology Resources* 15(4): 855–867.
- Carlson SM & Seamons TR. 2008. A review of quantitative genetic components of fitness in salmonids: implications for adaptation to future change. *Evolutionary Applications* 1 222-238.
- Casquet J., C. Thebaud, R.G. Gillespie. 2012. Chelex without boiling, a rapid and easy technique to obtain stable amplifiable DNA from small amounts of ethanol-stored spiders. *Molecular Ecology Resources*. Vol. 12, Iss. 1, 136-141.
- Chapman, D. W. Critical review of variables used to define effects of fines in redds of large salmonids. *Trans. Am. Fish. Soc.*, 117: 1–21 (1988).
- Chaput G & Benoit HP 2012 Evidence for bottom-up trophic effects on return rates to a second spawning for Atlantic salmon (*Salmo salar*) from the Miramichi River, Canada. *ICES Journal of Marine Science* 69 1656-1667.

- Chaput G & Jones R 2006 Reproductive rates and rebuilding potential for two multi-sea-winter Atlantic salmon (*Salmo salar* L.) stocks of the Maritime provinces. Ed FaO Canada: Canadian Science Advisory Secretariat.
- Christie MR, McNickle GG, French RA & Blouin MS 2018 Life history variation is maintained by fitness trade-offs and negative frequency-dependent selection. *Proceedings of the National Academy of Sciences of the United States of America* 115 4441-4446.
- Chudyk WE 1976 The life history of adult steelhead sampled in the Tyee test fishery in the Skeena river estuary and comparisons with other steelhead stocks in British Columbia. Ed BCFaW Branch.
- Chyb J, Mikolajczyk T & Breton B 1999 Post-ovulatory secretion of pituitary gonadotropins GtH I and GtH II in the rainbow trout (*Oncorhynchus mykiss*): regulation by steroids and possible role of non-steroidal gonadal factors. *Journal of Endocrinology* 163 87-97.
- Cohick WS & Clemmons DR 1994 Enhanced Expression of Dihydrofolate-Reductase by Bovine Kidney Epithelial-Cells Results in Altered Cell Morphology, Igf-I Responsiveness, and Igf Binding Protein-3 Expression. *Journal of Cellular Physiology* 161 178-186.
- Colotelo, A. H., R. A. Harnish, and B. W. Jones, and 10 other authors. 2014. Passage Distribution and Federal Columbia River Power System Survival for Steelhead Kelt steelhead Tagged Above and at Lower Granite Dam, Year 2. PNWL-23051, prepared for the U.S. Army Corp of Engineers, Walla Walla District, Walla Walla Washington, by Pacific Northwest National Laboratory, Richland, Washington.
[http://www.salmonrecovery.gov/Files/Comprehensive%20Evaluation/Colotelo-tal_2013_-%20Kelt steelhead-Passage-Distribution-Survival_PNWD-22101.pdf](http://www.salmonrecovery.gov/Files/Comprehensive%20Evaluation/Colotelo-tal_2013_-%20Kelt%20steelhead-Passage-Distribution-Survival_PNWD-22101.pdf).
- Colt J & Shearer KD 2001 Evaluation of migrational delays on the reproductive success of adult hatchery spring Chinook salmon in the Columbia and Snake Rivers (objective 2 only). 2001 Report to the U.S. Army Corps of Engineers, Contract W66QKZ00805700. Seattle: Northwest Fisheries Science Center, National Marine Fisheries Service.
- Copeland, T., R.V. Roberts, B.N. Oldemeyer, and K.A. Apperson. 2013. Idaho steelhead monitoring and evaluation studies, Annual Report 2012. IDFG 13-07.
- Courter II, Child DB, Hobbs JA, Garrison TM, Glessner JJG & Duery S 2013 Resident rainbow trout produce anadromous offspring in a large interior watershed. *Canadian Journal of Fisheries and Aquatic Sciences* 70 701-710.
- Crespi BJ & Teo R 2002 Comparative phylogenetic analysis of the evolution of semelparity and life history in salmonid fishes. *Evolution* 56 1008-1020.
- Crim LW, Wilson CE, So YP, Idler DR & Johnston CE 1992 Feeding, reconditioning, and rematuration responses of captive Atlantic salmon (*Salmo salar*) kelt. *Canadian Journal of Fisheries and Aquatic Sciences* 49 1835-1842.
- Crossin GT & Hinch SG 2005 A nonlethal, rapid method for assessing the somatic energy content of migrating adult pacific salmon. *Transactions of the American Fisheries Society* 134 184-191.
- Crossin GT, Phillips RA, Trathan PN, Fox DS, Dawson A, Wynne-Edwards KE & Williams TD 2012 Migratory carryover effects and endocrinological correlates of reproductive decisions and reproductive success in female albatrosses. *General and Comparative Endocrinology* 176 151-157.

- Daughaday WH, Mariz IK & Blethen SL 1980 Inhibition of access of bound somatomedin to membrane receptor and immunobinding sites: a comparison of radioreceptor and radioimmunoassay of somatomedin in native and acid-ethanol-extracted serum. *J Clin Endocrinol Metab* 51 781-788.
- Daughaday WH & Rotwein P 1989 Insulin-Like Growth Factor-I and Factor-II - Peptide, Messenger Ribonucleic-Acid and Gene Structures, Serum, and Tissue Concentrations. *Endocrine Reviews* 10 68-91.
- De Mones A, Fostier A, Cauty C & Jalabert B 1989 Ovarian early postovulatory development and estrogen production in rainbow trout (*Salmo gairdneri* R.) from a spring-spawning strain. *General and Comparative Endocrinology* 74 431-441.
- Dickhoff WW 1989 Salmonids and annual fishes: Death after sex. In *Development, maturation, and senescence of neuroendocrine systems: A comparative approach*, pp 253-266. Eds MP Schreibman & CG Scanes. New York: Academic Press.
- Duan C & Plisetskaya EM 1993 Nutritional Regulation of Insulin-Like Growth Factor-I Messenger-Rna Expression in Salmon Tissues. *Journal of Endocrinology* 139 243-252.
- Duan CM 1998 Nutritional and developmental regulation of insulin-like growth factors in fish. *Journal of Nutrition* 128 306s-314s.
- Duan CM & Xu QJ 2005 Roles of insulin-like growth factor (IGF) binding proteins in regulating IGF actions. *General and Comparative Endocrinology* 142 44-52.
- Dumas J, Barriere L, Blanc D, Godard J & Kaushik SJ 1991 Reconditioning of Atlantic salmon (*Salmo salar*) kelts with silage-based diets - growth and reproductive performance. *Aquaculture* 96 43-56.
- Dyer AR, Upton Z, Stone D, Thomas PM, Soole KL, Higgs N, Quinn K & Carragher JF 2004 Development and validation of a radioimmunoassay for fish insulin-like growth factor I (IGF-I) and the effect of aquaculture related stressors on circulating IGF-I levels. *Gen Comp Endocrinol* 135 268-275.
- Ensing, D., Crozier, W.W., Boylan, P., O'Maoil_eidigh, N. & McGinnity, P. (2013) An analysis of genetic stock identification on a small geographical scale using microsatellite markers, and its application in the management of a mixed-stock fishery for Atlantic salmon *Salmo salar* in Ireland. *Journal of Fish Biology* 82:2080–2094.
- Evans, A.F., R.E. Beaty, D.R. Hatch, J. Blodgett, & D. Fast. 2001. Kelt reconditioning: A research project to enhance iteroparity in Columbia Basin steelhead (*Oncorhynchus mykiss*). 2000 Annual Report to U.S. Dept. of Energy, Bonneville Power Administration, Project No. 2000-017. Prepared by the Columbia River Inter-Tribal Fish Commission, Portland, OR.
- Everest, F. L., R. L. Beschta, J. C. Scrivener, K. V. Koski, J. R. Sedell, and C. J. Cederholm. Fine sediment and salmonid production: A paradox. In: *Streamside Management: Forestry and Fishery Interactions* (E. O. Salo and T. W. Cundy, Eds., pp. 98–142). Seattle: College of Forest Resources, University of Washington (1987).
- Farrell AP, Gallagher PE, Fraser J, Pike D, Bowering P, Hadwin AKM, Parkhouse W & Routledge R 2001 Successful recovery of the physiological status of coho salmon on board a commercial gillnet vessel by means of a newly designed revival box. *Canadian Journal of Fisheries and Aquatic Sciences* 58 1932-1946.

- Ferriss BE, Trudel M & Beckman BR 2014 Regional and inter-annual trends in marine growth of juvenile salmon in coastal pelagic ecosystems of British Columbia, Canada. *Marine Ecology Progress Series* 503 247-261.
- Fisker S 2006 Physiology and pathophysiology of growth hormone-binding protein: methodological and clinical aspects. *Growth Horm IGF Res* 16 1-28.
- Fleming, I. A. 1998. Pattern and variability in the breeding system of Atlantic salmon (*Salmo salar*), with comparisons to other salmonids. *Canadian Journal of Fisheries and Aquatic Sciences* 55 (Suppl. 1), 59–76.
- Ford, M. J. (ed.). 2011. Status review update for Pacific salmon and steelhead listed under the Endangered Species Act: Pacific Northwest. U.S. Dept. Commer., NOAA Tech. Memo. NMFS-NWFSC-113, 281 p.
- Fostier A, Weil C, Terqui M, Breton B & Jalabert B 1978 Plasma estradiol-17-beta and gonadotropin during ovulation in rainbow trout (*Salmo gairdneri* R). *Annales De Biologie Animale Biochimie Biophysique* 18 929-936.
- Frederiksen CR, Fast DE, Bosch WJ & Temple GM 2015 Yakima Steelhead VSP Project: Yakima River Steelhead Population Status and Trends Monitoring, 10/15/2013-10/14/2014 Annual Report, 2010-030-00. Bonneville Power Administration.
- Frederiksen, C.R., D.E. Fast, W.J. Bosch, and G.M. Temple. Yakima Steelhead VSP Project: Yakima River Steelhead Population Status & Trends Monitoring, 10/15/2011 - 10/14/2013 Annual Report, 2010-030-00
- Frederiksen DR, Fast D & Temple G 2012 Yakima Steelhead Viable Salmonid Population (VSP) Status & Trends Monitoring. Yakima Steelhead VSP Project Annual Report 2011, Project No. 201003000. Toppenish, WA: Prepared by Yakama Nation Fisheries and Washington Department of Fish and Wildlife.
- Fruchtmann S, Jackson L & Borski R 2000 Insulin-like growth factor I disparately regulates prolactin and growth hormone synthesis and secretion: studies using the teleost pituitary model. *Endocrinology* 141 2886-2894.
- Gabillard JC, Kamangar BB & Montserrat N 2006 Coordinated regulation of the GH/IGF system genes during refeeding in rainbow trout (*Oncorhynchus mykiss*). *Journal of Endocrinology* 191 15-24.
- Gauthey Z, Freychet M, Manicki A, Herman A, Lepais O, Panzerat S, Elozeigi A, Tentelie C & Labonne J 2015 The concentration of plasma metabolites varies throughout reproduction and affects offspring number in wild brown trout (*Salmo trutta*). *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 184 90-96.
- Geist, David R. 2000. "Hyporheic Discharge of River Water into Fall Chinook Salmon (*Oncorhynchus tshawytscha*) Spawning Areas in the Hanford Reach, Columbia River." *Canadian Journal of Fisheries and Aquatic Sciences* 57: 1647–56.
- Gomez JM, Boujard T, Fostier A & LeBail PY 1996 Characterization of growth hormone nycthemeral plasma profiles in catheterized rainbow trout (*Oncorhynchus mykiss*). *Journal of Experimental Zoology* 274 171-180.
- Gray ES, Young G & Bern HA 1990 Radioreceptor assay for growth hormone in coho salmon (*Oncorhynchus kisutch*) and its application to the study of stunting. *J Exp Zool* 256 290-296.

- Gross MR 1996 Alternative reproductive strategies and tactics: Diversity within sexes. *Trends in Ecology & Evolution* 11 92-98.
- Guidelines for Electrofishing Waters Containing Salmonids Listed under the Endangered Species Act (Backpack Electrofishing Guidelines, NMFS, June 2000)
- Halttunen E 2011 Staying alive - the survival and importance of Atlantic salmon post-spawners. In Department of Arctic and Marine Biology: University of Tromso, Norway.
- Halver JE & Hardy RW 2002 Fish Nutrition. San Diego: Academic Press.
- Ham BR, Myrick CA, Barrows FT, Yeoman CJ, Duff GC, Maskill MG & Sealey WM 2015 Feed Characteristics Alter Growth Efficiency of Cutthroat Trout. *Journal of Fish and Wildlife Management* 6 83-91.
- Hanson KC, Ostrand KG, Gannam AL & Ostrand SL 2010 Comparison and validation of nonlethal techniques for estimating condition in juvenile salmonids. *Transactions of the American Fisheries Society* 139 1733-1741.
- Hardy RW & Barrows FT 2002 Diet Formulation and Manufacture. In Fish Nutrition, pp 506-600. Eds JE Halver & RW Hardy. San Diego: Academic Press.
- Hatch D.R., R. Branstetter, J. Stephenson, A. Pierce, S. Willis, J. Newell, W. Bosch, S. Everett, N. Graham, L. Medeiros, L. Jenkins, B. Hoffman, N. Hoffmann, T. Cavileer, J. Nagler, M. Fiander, C. Frederickson, J. Blodgett, D. Fast, and R. Johnson. 2020 Kelt Reconditioning and Reproductive Success Evaluation Research. 1/1/2019 - 12/31/2019 Bonneville Power Administration Annual Report, 2007-401-00. Portland, OR: Prepared by the Columbia River Inter-Tribal Fish Commission.
- Hatch D.R., R. Branstetter R, Stephenson J, Pierce A, Lessard R, Newell J, Matala A, Bosch W, Everett S, Graham N, et al. 2019. Kelt Reconditioning and Reproductive Success Evaluation Research. 1/1/2018 - 12/31/2018 Bonneville Power Administration Annual Report, 2007-401-00. Portland, OR: Prepared by the Columbia River Inter-Tribal Fish Commission.
- Hatch, D. R., P. J. Anders, A. F. Evans, J. Blodgett, B. Bosch, D. Fast, & T. Newsome. 2002. Kelt reconditioning: A research project to enhance iteroparity in Columbia Basin steelhead (*Oncorhynchus mykiss*). Project 2000-017-00, Annual Report the Bonneville Power Administration, Portland, OR.
- Hatch, D.R., R. Branstetter, J. Blodgett, B. Bosch, D. Fast, & T. Newsome. 2003. Kelt reconditioning: A research project to enhance iteroparity in Columbia Basin steelhead (*Oncorhynchus mykiss*). 2002 Annual Report to U.S. Dept. of Energy, Bonneville Power Administration, Project No. 2000-017. Prepared by the Columbia River Inter-Tribal Fish Commission, Portland, OR.
- Hatch, D.R., R. Branstetter, J. Stephenson, A.L. Pierce, A. Matala, R. Lessard, W. Bosch, L.K. Caldwell, S.R. Everett, J. Newell, N. Graham, L. Jenkins, M. Elliot, T. Cavileer, J. Nagler, M. Fiander, J. Blodgett, C. Frederiksen, D. Fast, K. J.M. Whiteaker, R. Johnson. 2015. Kelt Reconditioning and Reproductive Success Evaluation Research. 1/1/2014-12/31/2014 Annual Report to the U.S. Dept. of Energy, Bonneville Power Administration, Project No. 2007-401-000. Portland, OR: Prepared by the Columbia River Inter-Tribal Fish Commission.
- Hatch D.R., Branstetter R, Stephenson J, Pierce AL, Matala A & Newell J. 2013a. Steelhead Kelt Reconditioning and Reproductive Success, 2012 Annual Report to the U.S. Dept. of

- Energy, Bonneville Power Administration, Project No. 2007-401-000. Portland, OR: Prepared by the Columbia River Inter-Tribal Fish Commission.
- Hatch D.R., Branstetter R, Stephenson J, Pierce AL, Whiteaker JM & Bosch B. 2012. Steelhead Kelt Reconditioning and Reproductive Success, 2011 Annual Report to the U.S. Dept. of Energy, Bonneville Power Administration, Project No. 2007-401-000. Portland, OR: Prepared by the Columbia River Inter-Tribal Fish Commission.
- Hatch D.R., Branstetter R, Stephenson J, Pierce AL, Matala A, Whiteaker JM, Lessard R, Caldwell LK, Everett SR, Bosch B, et al. 2014. Kelt Reconditioning and Reproductive Success Evaluation Research, 2013 Annual Report to the U.S. Dept. of Energy, Bonneville Power Administration, Project No. 2007-401-000. Portland, OR: Prepared by the Columbia River Inter-Tribal Fish Commission.
- Hatch, D.R., D.E. Fast, W.J. Bosch, R. Branstetter, J.W. Blodgett, J.M. Whiteaker, & A.L. Pierce. 2013b. Survival and traits of reconditioned kelt steelhead (*Oncorhynchus mykiss*) in the Yakima River, Washington. *North American Journal of Fisheries Management* 33: 615–625.
- Hatch, D., R. Branstetter, J. Stephenson, A. Pierce, A. Matala, R. Lessard, W. Bosch, S. Everett, J. Newell, N. Graham, L. Medeiros, L. Jenkins, T. Tall Bull, M. Elliott, K. Huggler, T. Cavileer, J. Nagler, M. Fiander, C. Frederickson, J. Blodgett, D. Fast, J. Whiteaker, and R. Johnson. 2016a. Kelt Reconditioning and Reproductive Success Evaluation Research: 2015 Annual Technical Report. Bonneville Power Administration Annual Report, 2007-401-00, 156 p.
- Hatch DR, Branstetter R, Stephenson J, Pierce AL, Newell J, Bosch B, Everett S, Graham N, Medeiros L, Jenkins L et al. 2017. Kelt reconditioning and reproductive success evaluation research, 1/1/2016 - 12/31/2016 Bonneville power administration 2016 annual technical report. Portland, OR: Columbia River Inter-Tribal Fish Commission, 2007-401-00.
- Hatch DR, Branstetter R, Stephenson J, Pierce AL, Newell J, Bosch WJ, Graham N, Medeiros LR, Jenkins LE, Hoffman B, et al. 2018 Kelt Reconditioning and Reproductive Success Evaluation Research. Portland, OR: Bonneville Power Administration Annual Report, 2007-401-00 (Prepared by the Columbia River Inter-Tribal Fish Commission).
- Henderson BA & Wong JL 1998 Control of lake trout reproduction: role of lipids. *Journal of Fish Biology* 52 1078-1082.
- Hess, J. E., N. Campbell, A. P. Matala and S. R. Narum. 2012. Genetic assessment of Columbia River Stocks: 2011 annual report. Submitted to Bonneville Power Administration. Contract #41224; Project # 2008-907-00
<https://pisces.bpa.gov/release/documents/documentviewer.aspx?doc=P120015>
- Hess, J. E., J. M. Whiteaker, J. K. Fryer and S. R. Narum. 2014. Monitoring Stock-Specific Abundance, Run Timing, and Straying of Chinook salmon in the Columbia River Using Genetic Stock Identification (GSI). *Transactions of the American Fisheries Society* 34:184–201.
- Hess J.E.. 2020, GENETIC ASSESSMENT OF COLUMBIA RIVER STOCKS, 1/1/2019 - 12/31/2019 Annual Report, 2008-907-00Hevroy EM, Hunskar C, de Gelder S, Shimizu M, Waagbo R, Breck O, Takle H, Sussort S & Hansen T 2013 GH-IGF system regulation of attenuated muscle growth and lipolysis in Atlantic salmon reared at elevated sea temperatures.

- Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology 183 243-259.
- Hill GE 2011 Condition-dependent traits as signals of the functionality of vital cellular processes. Ecology Letters 14 625-634.
- Hockersmith, E., J. Vella, L. Stuehrenberg, R. Iwamoto, & G. Swan. 1995. Yakima River radio telemetry study: steelhead, 1989-93. Prepared for Bonneville Power Administration, P.O. Box 3621, Portland, OR. Project Number 89-089. Seattle, WA: Coastal Zone and Estuarine Studies Division, Northwest Fisheries Science Center, National Marine Fisheries Service, NOAA.
- Howell, P., K. Jones, D. Scarnecchia, L. LaVoy, W. Kendra. & D. Ortmann. 1985. Stock assessment of Columbia River anadromous salmonids Volume II: steelhead stock summaries stock transfer guidelines – information needs. Final Report to Bonneville Power Administration, Contract DE-AI79–84BP12737, Project 83-335.
- Huang YS, Rousseau K, Le Belle N, Vidal B, Burzawa-Gerard E, Marchelidon J & Dufour S 1998 Insulin-like growth factor-I stimulates gonadotrophin production from eel pituitary cells: a possible metabolic signal for induction of puberty. Journal of Endocrinology 159 43-52.
- Hutchings JA 2011 Old wine in new bottles: reaction norms in salmonid fishes. Heredity 106 421-437.
- ISRP 2011 Retrospective Report 2011. Independent Scientific Review Panel.
- ISRP. 2014. Review of 2014 Progress Report for the Yakama Nation's Upper Columbia Kelt Reconditioning Program, #2008-458-00, an Accord Proposal. Independent Scientific Review Panel, 2014-9.
- Iwama G.K., McGeer J.C. & Pawluk M.P. (1989). The effects of five fish anesthetics on acid-base balance, hematocrit, cortisol and adrenaline in rainbow trout. Canadian Journal of Zoology 67:2065-2073.
- Jeffries KM, Hinch SG, Donaldson MR, Gale MK, Burt JM, Thompson LA, Farrell AP, Patterson DA & Miller KM 2011 Temporal changes in blood variables during final maturation and senescence in male sockeye salmon *Oncorhynchus nerka*: reduced osmoregulatory ability can predict mortality. Journal of Fish Biology 79 449-465.
- Jenkins LE, Pierce AL, Graham N, Branstetter R, Hatch DR & Nagler JJ. 2018. Reproductive performance and energy balance in consecutive and repeat spawning female steelhead reconditioned in captivity. Transactions of the American Fisheries Society 147 959-971.
- Jenkins LE, Pierce AL, Graham ND, Medeiros LR, Hatch DR & Nagler JJ 2019 Elevated plasma triglycerides and growth rate are early indicators of reproductive status in post-spawning female steelhead trout (*Oncorhynchus mykiss*). Conservation Physiology 7.
- Jenkins, L.E., A.L. Pierce, Christopher C. Caudill, N.D. Graham, L.R. Medeiros, D.R. Hatch, J.J. Nagler. 2020. Effects of physiological condition on aspects of repeat spawning in female Steelhead Trout reconditioned in captivity. Transactions of the American Fisheries Society.
- Jenkins RC & Ross RJM 1996 Acquired growth hormone resistance in catabolic states. Baillieres Clinical Endocrinology and Metabolism 10 411-419.
- Johnston CE, Gray RW, McLennan A & Paterson A 1987 Effects of photoperiod, temperature, and diet on the reconditioning response, blood chemistry, and gonad maturation of

- Atlantic salmon kelts (*Salmo salar*) held in fresh water. *Canadian Journal of Fisheries and Aquatic Sciences* 44 702-711.
- Jorgensen C, Ernande B, Fiksen O & Dieckmann U 2006 The logic of skipped spawning in fish. *Canadian Journal of Fisheries and Aquatic Sciences* 63 200-211.
- Jensen, D.W., E. Ashley Steel, A.H. Fullerton, and G.R. Pess. 2009. Impact of Fine Sediment on Egg-To-Fry Survival of Pacific Salmon: A Meta-Analysis of Published Studies. *Reviews in Fisheries Science*, 17(3):348-359.
- Justice, C., S. White, and D. McCullough. 2012. Spawning Gravel Composition Survey Methods. A Component of Monitoring Recovery Trends in Key Spring Chinook Habitat Variables and Validation of Population Viability Indicators BPA Annual Report 2012.
- Kadri S, Mitchell DF, Metcalfe NB, Huntingford FA & Thorpe JE 1996 Differential patterns of feeding and resource accumulation in maturing and immature Atlantic salmon, *Salmo salar*. *Aquaculture* 142 245-257.
- Kalinowski S.T., M.L. Taper, and T.C. Marshall. 2007. Revising how the computer program CERVUS accommodates genotyping error increases confidence in paternity. *Molecular Ecology* 16:1099-1106.
- Kato Y, Murakami Y, Sohmiya M & Nishiki M 2002 Regulation of human growth hormone secretion and its disorders. *Internal Medicine* 41 7-13.
- Keefer ML, Wertheimer RH, Evans AF, Boggs CT & Peery CA. 2008. Iteroparity in Columbia river summer-run steelhead (*Oncorhynchus mykiss*): implications for conservation. *Canadian Journal of Fisheries and Aquatic Sciences* 65 2592-2605.
- Kelley KM, Desai P, Roth JT, Haigwood JT, Arope SA, Flores RM, Schmidt KE, Perez M, Nicholson GS & Song WW 2000 Evolution of endocrine growth regulation: the insulin like growth factors (IGFs), their regulatory binding proteins (IGFBPs), and IGF receptors in fishes and other ectothermic vertebrates. In *Recent Advances in Marine Biotechnology*, pp 189-228. Eds M Fingerman, M-F Thompson & R Nagabhushanam. New Delhi: Oxford and IBH Publishing.
- Kendall NW, McMillan JR, Sloat MR, Buehrens TW, Quinn TP, Pess GR, Kuzishchin KV, McClure MM & Zabel RW 2015 Anadromy and residency in steelhead and rainbow trout (*Oncorhynchus mykiss*): a review of the processes and patterns. *Canadian Journal of Fisheries and Aquatic Sciences* 72 319-342.
- Kennedy J, Witthames PR, Nash RDM & Fox CJ 2008 Is fecundity in plaice (*Pleuronectes platessa* L.) down-regulated in response to reduced food intake during autumn? *Journal of Fish Biology* 72 78-92.
- Krogdahl A & Bakke-McKellep AM 2005 Fasting and refeeding cause rapid changes in intestinal tissue mass and digestive enzyme capacities of Atlantic salmon (*Salmo salar* L.). *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 141 450-460.
- Lahood, E. S., J. J. Miller, C. Apland, and M. J. Ford. 2008. A rapid, ethanol-free fish tissue collection method for molecular genetic analyses. *Transactions of the American Fisheries Society* 137(4):1104–1107.
- Larsen DA, Beckman BR & Dickhoff WW 2001 The effect of low temperature and fasting during the winter on growth and smoltification of coho salmon, *Oncorhynchus kisutch*. *North American Journal of Aquaculture* in press.

- Larsen DA, Beckman BR, Strom CR, Parkins PJ, Cooper KA, Fast DE & Dickhoff WW 2006 Growth modulation alters the incidence of early male maturation and physiological development of hatchery-reared Spring Chinook Salmon: A comparison with wild fish. *Transactions of the American Fisheries Society* 135 1017-1032.
- Le Bail PY, Sumpter JP, Carragher JF, Mourot B, Niu PD & Weil C 1991 Development and validation of a highly sensitive radioimmunoassay for chinook salmon (*Oncorhynchus tshawytscha*) growth hormone. *Gen Comp Endocrinol* 83 75-85.
- Leitritz, E., and R.C. Lewis. 1976. Trout and Salmon Culture (Hatchery Methods). State of California The Resources Agency Department of Fish and Game Fish Bulletin 164.
- Lindsey, C. C., T. G. Nortcote, and G. F. Hartman. 1959. Homing of rainbow trout to inlet and outlet spawning streams at Loon Lake, British Columbia. *Journal of the Fisheries Research Board of Canada* 16:695–719.
- Lotspeich, F.B., and F.H. Everest. 1981. A New Method for Reporting and Interpreting Textural Composition of Spawning Gravel. U.S. Department of Agriculture: Forest Service; Pacific Northwest Forest and Range Experiment Station; Research Note PNW-369.
- Lubzens E, Young G, Bobe J & Cerda J 2010 Oogenesis in teleosts: how eggs are formed. *General and Comparative Endocrinology* 165 367-389.
- Luckenbach JA, Dickey JT & Swanson P 2010 Regulation of pituitary GnRH receptor and gonadotropin subunits by IGF1 and GnRH in prepubertal male coho salmon. *General and Comparative Endocrinology* 167 387-396.
- Lupu F, Terwilliger JD, Lee K, Segre GV & Efstratiadis A 2001 Roles of growth hormone and insulin-like growth factor 1 in mouse postnatal growth. *Developmental Biology* 229 141-162.
- Marshall, T.C.J. Slate, L. Kruuk, and J.M. Pemberton. 1998. Statistical confidence for likelihood-based paternity inference in natural populations. *Molecular Ecology* 7:639-655.
- Matala, A. P., M. W. Ackerman, M. R. Campbell, S. R. Narum (2014) Relative contributions of neutral and non-neutral genetic differentiation to inform conservation of steelhead trout across highly variable landscapes. *Evolutionary Applications*, 7, 682–701.
- McBride RS, Somarakis S, Fitzhugh GR, Albert A, Yaragina NA, Wuenschel MJ, Alonso-Fernandez A & Basilone G. 2015. Energy acquisition and allocation to egg production in relation to fish reproductive strategies. *Fish and Fisheries* 16 23-57.
- McCleave, J. D. 1967. Homing and orientation of cutthroat trout (*Salmo clarki*) in Yellowstone Lake, with special reference to olfaction and vision. *Journal of the Fisheries Research Board of Canada* 24:2011–2044.
- McCormick SD 2012 Smolt Physiology and Endocrinology. In *Fish Physiology*, pp 200-253. Eds SD McCormick, AP Farrell & CJ Brauner.
- McGlaufflin, M. T., D. E. Schindler, L.W. Seeb, C. T. Smith, C. Habicht and J. E. Seeb. 2011. Spawning Habitat and Geography Influence Population Structure and Juvenile Migration Timing of Sockeye Salmon in the Wood River Lakes, Alaska, *Transactions of the American Fisheries Society*, 140:3, 763-782, DOI: 10.1080/00028487.2011.584495
- Meador JP, Sommers FC, Ylitalo GM & Sloan CA 2006 Altered growth and related physiological responses in juvenile Chinook salmon (*Oncorhynchus tshawytscha*) from dietary exposure to polycyclic aromatic hydrocarbons (PAHs). *Canadian Journal of Fisheries and Aquatic Sciences* 63 2364-2376.

- Medeiros LR, Elliott M & Nagler JJ. 2016. Stressor timing, not cortisol, is an important embryo viability determinant in female rainbow trout *Oncorhynchus mykiss*. *Journal of Fish Biology* 88 557-566.
- Medeiros LR, Galbreath PF, Knudsen CM, Stockton CA, Koch IJ, Bosch WJ, Narum SR, Nagler JJ & Pierce AL. 2018. Plasma 11-Ketotestosterone in Individual Age-1 Spring Chinook Salmon Males Accurately Predicts Age-2 Maturation Status. *Transactions of the American Fisheries Society* 147 1042-1051.
- Medeiros, L.R., J.J. Nagler, A.L. Pierce. 2020. Establishment of time-resolved fluoroimmunoassays for detection of growth hormone and insulin-like growth factor I in rainbow trout plasma. *Comparative Biochemistry and Physiology*.
- Moffett IJJ, Kennedy GJA & Crozier WW 1996 Freshwater reconditioning and ranching of Atlantic salmon, *Salmo salar* L., kelts: growth and reproductive performance. *Fisheries Management and Ecology* 3 35-44.
- Mommsen TP 1998 Growth and metabolism. In *Physiology of Fishes*, pp 65-97. Ed D Evans. Boca Raton, Florida: CRC Press.
- Mommsen TP, French CJ & Hochachka PW 1980 Sites and patterns of protein and amino acid utilization during the spawning migration of salmon. *Can J Zool* 58 1785-1799.
- Moore DS, Chaput G & Pickard R 1995 The effect of fisheries on the biological characteristics and survival of mature Atlantic salmon (*Salmo salar*) from the Miramichi River. In *Water, Science, and the Public: the Miramichi Ecosystem*, pp 229-247. Ed EMP Chadwick. Volume 123. Canadian Special Publication of Fisheries and Aquatic Sciences, National Research Council of Canada, Ottawa.
- Moore, J.W., J.D. Yeakel, D. Peard, J. Lough, and M. Beere. 2014. Life-history diversity and its importance to population stability and persistence of a migratory fish: steelhead in two large North American watersheds. *Journal of Animal Ecology* 83, 1035-1046.
- Moriyama S, Ayson FG & Kawauchi H 2000 Growth regulation by insulin-like growth factor-I in fish. *Bioscience Biotechnology and Biochemistry* 64 1553-1562.
- Moriyama S, Swanson P, Nishii M, Takahashi A, Kawauchi H, Dickhoff WW & Plisetskaya EM 1994 Development of a Homologous Radioimmunoassay for Coho Salmon Insulin-Like Growth-Factor-I. *General and Comparative Endocrinology* 96 149-161.
- Nagahama Y 1994 Endocrine regulation of gametogenesis in fish. *International Journal of Developmental Biology* 38 217-229.
- Nagler JJ, Cavileer TD, Verducci JS, Schultz IR, Hook SE & Hayton WL 2012 Estrogen receptor mRNA expression patterns in the liver and ovary of female rainbow trout over a complete reproductive cycle. *General and Comparative Endocrinology* 178 556-561.
- Narum, S. R., D. Hatch, A. J. Talbot, P. Moran and M. S. Powell. 2008. Iteroparity in complex mating systems of steelhead *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Biology* 72:1-16.
- Navarro I & Gutierrez J 1995 Fasting and starvation. In *Biochemistry and Molecular Biology of Fishes*, pp 393-434. Eds PW Hochachka & TP Mommsen. Amsterdam: Elsevier Science.
- NMFS (National Marine Fisheries Service). 1996. Status review of west coast steelhead from Washington, Idaho, Oregon, and California. Seattle, WA.
- NMFS (National Marine Fisheries Service). 2008. Endangered Species Act Section 7(a) (2) Consultation Biological Opinion and Magnuson-Stevens Fishery Conservation and

- Management Act Essential Fish Habitat Consultation on Remand for Operation of the Federal Columbia River Power System, 11 Bureau of Reclamation Projects in the Columbia Basin and ESA Section 10(a) (I) (A) Permit for Juvenile Fish Transportation Program (Revised and reissued pursuant to court order, NWF v. NMFS, Civ. No. CV 01-640-RE (D. Oregon))
- NMFS (National Marine Fisheries Service). 2000. Guidelines for Electrofishing Waters Containing Salmonids Listed Under the Endangered Species Act.
- NMFS (National Marine Fisheries Service). 2010. Endangered Species Act Section 7(a) (2) Consultation Supplemental Biological Opinion Supplemental Consultation on Remand for Operation of the Federal Columbia River Power System, 11 Bureau of Reclamation Projects in the Columbia Basin and ESA Section 10(a) (I) (A) Permit for Juvenile Fish Transportation Program.
- NMFS (National Marine Fisheries Service). 2014. Endangered Species Act Section 7(a) (2) Supplemental Biological Opinion Supplemental Consultation on Remand for Operation of the Federal Columbia River Power System.
- NMFS (National Marine Fisheries Service). 2019. Endangered Species Act Section 7(a) (2) Consultation Biological Opinion and Magnuson-Stevens Fishery Conservation and Management Act Essential Fish Habitat Response Continued Operation and Maintenance of the Columbia River System. NMFS Consultation Number: WCRO-2018-00152. March 29, 2019.
- NPCC (National Power and Conservation Council. 1980. Pacific Northwest Electric Power Planning and Conservation Act. 16 U.S. code Ch. 12H (1994 & Supp. I 1995). Act of Dec. 5, 1980, 94 Stat. 2697. P.L. No. 96-501, S. 885.
- NPCC (National Power and Conservation Council. 2017. 2017 Research Plan. Doc. 2017-4/June 2017.
- Norbeck LA, Kittilson JD & Sheridan MA 2007 Resolving the growth-promoting and metabolic effects of growth hormone: Differential regulation of GH-IGF-I system components. *General and Comparative Endocrinology* 151 332-341.
- Nielsen JL, Turner SM & Zimmerman CE 2011 Electronic tags and genetics explore variation in migrating steelhead kelts (*Oncorhynchus mykiss*), Ninilchik River, Alaska. *Canadian Journal of Fisheries and Aquatic Sciences* 68 1-16.
- Niemelä E, Erkinaro J, Julkunen M, Hassinen E, Lansman M & Brors S 2006a Temporal variation in abundance, return rate and life histories of previously spawned Atlantic salmon in a large subarctic river. *Journal of Fish Biology* 68 1222-1240.
- Niemelä E, Orell P, Erkinaro J, Dempson JB, Brors S, Svenning MA & Hassinen E 2006b Previously spawned Atlantic salmon ascend a large subarctic river earlier than their maiden counterparts. *Journal of Fish Biology* 69 1151-1163.
- Norberg B & Haux C 1985 Induction, isolation and a characterization of the lipid content of plasma vitellogenin from 2 salmo species - rainbow trout (*Salmo gairdneri*) and sea trout (*Salmo trutta*). *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* 81 869-876.
- Oguri M & Ooshima Y 1977 Early changes in plasma osmolality and ionic concentrations of rainbow trout and goldfish following direct transfer from fresh-water to sea water. *Bulletin of the Japanese Society of Scientific Fisheries* 43 1253-1257.

- Olsen, J.B., P. Bentzen, M.A. Banks, J.B. Shaklee, & S. Young. 2000. Microsatellites reveal population identity of individual pink salmon to allow supportive breeding of a population at risk of extinction. *Transactions of the American Fisheries Society* 129:232-242.
- Osborne, V. (Ed.). (2015, April 1). Anglers Help Catch Steelhead Brood Stock. Retrieved January 14, 2016, from <https://fishgame.idaho.gov/content/article/anglers-help-catch-steelhead-broodstock>.
- Palstra AP, Crespo D, van den Thillart GE & Planas JV 2010 Saving energy to fuel exercise: swimming suppresses oocyte development and downregulates ovarian transcriptomic response of rainbow trout *Oncorhynchus mykiss*. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* 299 R486-499.
- Pankhurst NW, King HR, Anderson K, Elizur A, Pankhurst PM & Ruff N 2011 Thermal impairment of reproduction is differentially expressed in maiden and repeat spawning Atlantic salmon. *Aquaculture* 316 77-87.
- Penney ZL & Moffitt CM 2014a Histological assessment of organs in sexually mature and post-spawning steelhead trout and insights into iteroparity. *Reviews in Fish Biology and Fisheries* 24 781-801.
- Penney ZL & Moffitt CM 2014b Proximate composition and energy density of stream-maturing adult steelhead during upstream migration, sexual maturity, and kelt emigration. *Transactions of the American Fisheries Society* 143 399-413.
- Penney ZL & Moffitt CM 2015 Fatty-acid profiles of white muscle and liver in stream-maturing steelhead trout *Oncorhynchus mykiss* from early migration to kelt emigration. *Journal of Fish Biology* 86 105-120.
- Perez-Sanchez J & Le Bail PY 1999 Growth hormone axis as marker of nutritional status and growth performance in fish. *Aquaculture* 177 117-128.
- Perez-Sanchez J, Marti-Palanca H & Kaushik SJ 1995 Ration size and protein intake affect circulating growth hormone concentration, hepatic growth hormone binding and plasma insulin-like growth factor-I immunoreactivity in a marine teleost, the gilthead sea bream (*Sparus aurata*). *J Nutr* 125 546-552.
- Perez-Sanchez J, Simo-Mirabet P, Naya-Catala F, Martos-Sitcha JA, Perera E, Bermejo-Nogales A, Benedito-Palos L & Caldach-Giner JA. 2018. Somatotrophic Axis Regulation Unravels the Differential Effects of Nutritional and Environmental Factors in Growth Performance of Marine Farmed Fishes. *Frontiers in Endocrinology* 9.
- Perez-Sanchez J, Weil C & Le Bail PY 1992 Effects of human insulin-like growth factor-I on release of growth hormone by rainbow trout (*Oncorhynchus mykiss*) pituitary cells. *J Exp Zool* 262 287-290.
- Picha ME, Turano MJ, Beckman BR & Borski RJ 2008a Endocrine biomarkers of growth and applications to aquaculture: A minireview of growth hormone, insulin-like growth factor (IGF)-I, and IGF-Binding proteins as potential growth indicators in fish. *North American Journal of Aquaculture* 70 196-211.
- Picha ME, Turano MJ, Tipsmark CK & Borski RJ 2008b Regulation of endocrine and paracrine sources of Igfs and Gh receptor during compensatory growth in hybrid striped bass (*Morone chrysops* X *Morone saxatilis*). *Journal of Endocrinology* 199 81-94.

- Pierce AL, Beckman BR, Shearer KD, Larsen DA & Dickhoff WW 2001 Effects of ration on somatotrophic hormones and growth in coho salmon. *Comp Biochem Physiol B Biochem Mol Biol* 128 255-264.
- Pierce AL, Blodgett JW, Cavileer TD, Medeiros LR, Boyce J, Caldwell LK, Bosch WJ, Branstetter R, Fast DE, Hatch DR, et al. 2017 Reproductive development in captive reconditioned female steelhead kelts: evidence for consecutive and skip spawning life histories. *Canadian Journal of Fisheries and Aquatic Sciences* 74 1049-1060.
- Pierce AL, Breves JP, Moriyama S, Hirano T & Grau EG 2011 Differential regulation of Igf1 and Igf2 mRNA levels in tilapia hepatocytes: effects of insulin and cortisol on GH sensitivity. *Journal of Endocrinology* 211 201-210.
- Pierce AL, Fukada H & Dickhoff WW 2005a Metabolic hormones modulate the effect of growth hormone (GH) on insulin-like growth factor-I (IGF-I) mRNA level in primary culture of salmon hepatocytes. *Journal of Endocrinology* 184 341-349.
- Pierce AL, Shimizu M, Beckman BR, Baker DM & Dickhoff WW 2005b Time course of the GH/IGF axis response to fasting and increased ration in chinook salmon (*Oncorhynchus tshawytscha*). *Gen Comp Endocrinol* 140 192-202.
- Piry, S., A. Alapetite, J. M. Cornuet, D. Paetkau, L. Baudouin and A. Estoup. 2004. GeneClass2: A Software for Genetic Assignment and First-Generation Migrant Detection. *Journal of Heredity* 95:536-539.
- Plummer, M. 2003. JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. *Proceedings of the 3rd International Workshop on Distributed Statistical Computing*. Vienna, Austria.
- Prat F, Sumpter JP & Tyler CR 1996 Validation of radioimmunoassays for two salmon gonadotropins (GTH I and GTH II) and their plasma concentrations throughout the reproductive cycle in male and female rainbow trout (*Oncorhynchus mykiss*). *Biology of Reproduction* 54 1375-1382.
- Quinn TP 2005 *The Behavior and Ecology of Pacific Salmon and Trout*. Seattle: University of Washington Press.
- Quinn TP, McGinnity P & Reed TE 2016 The paradox of "premature migration" by adult anadromous salmonid fishes: patterns and hypotheses. *Canadian Journal of Fisheries and Aquatic Sciences* 73 1015-1030.
- Quinn TP & Myers KW 2004 Anadromy and the marine migrations of Pacific salmon and trout: Rounsefell revisited. *Reviews in Fish Biology and Fisheries* 14 421-442.
- Quinn TP, Seamons TR, Vøllestad LA & Duffy E 2011 Effects of Growth and Reproductive History on the Egg Size-Fecundity Trade-off in Steelhead. *Transactions of the American Fisheries Society* 140 45-51.
- Rannala, B., and J. L. Mountain. 1997. Detecting immigration by using multilocus genotypes. *Proceedings of the National Academy of Sciences of the United States of America* 94(17):9197-9201.
- Reid JE & Chaput G 2012 Spawning history influence on fecundity, egg size, and egg survival of Atlantic salmon (*Salmo salar*) from the Miramichi River, New Brunswick, Canada. *ICES Journal of Marine Science* 69 1678-1685.
- Ricker, W.E. 1954. Stock and recruitment. *J. Fish. Res. Board Can.* 11: 559–623.

- Reindl KM & Sheridan MA 2012 Peripheral regulation of the growth hormone-insulin-like growth factor system in fish and other vertebrates. *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 163 231-245.
- Rideout RM, Rose GA & Burton MPM 2005 Skipped spawning in female iteroparous fishes. *Fish and Fisheries* 6 50-72.
- Rideout RM & Tomkiewicz J 2011 Skipped Spawning in Fishes: More Common than You Might Think. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* 3 176-189.
- Reiser, D.W. 1999. Sediment in gravel bed rivers: ecological and biological consideration. In *Gravel-bed rivers in the environment*, 199-225. Edited by P.C. Klingeman, R.L. Beschta, P.D. Komar, and J.B. Bradely. Highlands Ranch, Colorado: Water Resources Publications.
- Reiser, D. W., and R. G. White. Effects of two sediment size classes on survival of steelhead and Chinook salmon eggs. *N. Am. J. Fish. Management*, 8: 432–437 (1988).
- Roberts RJ, Johnson KA & Kasten MT 2004 Control of *Salmincola californiensis* (Copepoda: Lernaeapodidae) in rainbow trout, *Oncorhynchus mykiss* (Walbaum): a clinical and histopathological study. *Journal of Fish Diseases* 27 73-79.
- Rousseau K, Huang YS, Le Belle N, Vidal B, Marchelidon J, Epelbaum J & Dufour S 1998 Long-term inhibitory effects of somatostatin and insulin-like growth factor 1 on growth hormone release by serum-free primary culture of pituitary cells from European eel (*Anguilla anguilla*). *Neuroendocrinology* 67 301-309.
- Rowe DK & Thorpe JE 1990 Differences in growth between maturing and nonmaturing male Atlantic salmon, *Salmo salar* L., parr. *Journal of Fish Biology* 36 643-658.
- Sadoul B & Vijayan MM 2016 Stress and Growth. In *Fish Physiology*, pp 167-205.
- Satterthwaite WH, Beakes MP, Collins EM, Swank DR, Merz JE, Titus RG, Sogard SM & Mangel M 2009 Steelhead life history on California's Central Coast: Insights from a State - Dependent Model. *Transactions of the American Fisheries Society* 138 532-548.
- Seamons, TR, TP Quinn. 2010. Sex-specific patterns of lifetime reproductive success in single and repeat breeding steelhead trout (*Oncorhynchus mykiss*). *Behav. Ecol. Sociobiol.* 64:505–513. DOI 10.1007/s00265-009-0866-7.
- Schindler DE, Hilborn R, Chasco B, Boatright CP, Quinn TP, Rogers LA & Webster MS 2010 Population diversity and the portfolio effect in an exploited species. *Nature* 465 609-612.
- Shearer K, Parkins P, Gadberry B, Beckman B & Swanson P 2006 Effects of growth rate/body size and a low lipid diet on the incidence of early sexual maturation in juvenile male spring Chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture* 252 545-556.
- Shearer KD & Swanson P 2000 The effect of whole body lipid on early sexual maturation of 1+ age male Chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture* 190 343-367.
- Sheridan MA 1994 Regulation of lipid metabolism in poikilothermic vertebrates. *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* 107 495-508.
- Shimizu M, Swanson P, Fukada H, Hara A & Dickhoff WW 2000 Comparison of extraction methods and assay validation for salmon insulin-like growth factor-I using commercially available components. *General and Comparative Endocrinology* 119 26-36.
- Schindler, D. E., and coauthors. 2010. Population diversity and the portfolio effect in an exploited species. *Nature* 465(7298):609-612.

- Schreck CB, Contreras-Sanchez W & Fitzpatrick MS 2001 Effects of stress on fish reproduction, gamete quality, and progeny. *Aquaculture* 197 3-24.
- Schroder, S.L., C.M. Knudsen, T.N. Pearsons, T.W. Kassler, S.F. Young, C.A. Busack, & D.E. Fast. 2008. Breeding Success of Wild and First-Generation Hatchery Female Spring Chinook Salmon spawning in an Artificial Stream. *Transactions of the American Fisheries Society* 137:1475-1489.
- Schroder, S. L., C. M. Knudsen, T. N. Pearsons, T. W. Kassler, S. F. Young, E.P. Beall, & D. E. Fast. 2010. Behavior and Breeding Success of Wild and First-Generation Hatchery Male Spring Chinook Salmon Spawning in an Artificial Stream. *Transactions of the American Fisheries Society*, 139:989-1003.
- Shearer, K. D., and P. Swanson. 2000. The effect of whole body lipid on early sexual maturation of 1+age male chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture* 190(3-4):343-367.
- Silverstein JT, Shearer KD, Dickhoff WW & Plisetskaya EM 1998 Effects of growth and fatness on sexual development of chinook salmon (*Oncorhynchus tshawytscha*) parr. *Canadian Journal of Fisheries and Aquatic Sciences* 55 2376-2382.
- Simpkins DG, Hubert WA, Del Rio CM & Rule DC 2003 Physiological responses of juvenile rainbow trout to fasting and swimming activity: effects on body composition and condition indices. *Transactions of the American Fisheries Society* 132 576-589.
- Small BC & Peterson BC 2005 Establishment of a time-resolved fluoroimmunoassay for measuring plasma insulin-like growth factor I (IGF-I) in fish: effect of fasting on plasma concentrations and tissue mRNA expression of IGF-I and growth hormone (GH) in channel catfish (*Ictalurus punctatus*). *Domestic Animal Endocrinology* 28 202-215.
- Stead SM, Houlihan DF, McLay HA & Johnstone R 1999 Food consumption and growth in maturing Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences* 56 2019-2028.
- Stearns SC 1992 *The Evolution of Life Histories*. New York: Oxford University Press.
- Steele, C., M. Ackerman, J. McCane, M. Campbell, M. Hess and S. Narum. 2012. Parentage based tagging of Snake River hatchery-origin steelhead and Chinook salmon. 2011 Annual Report. Submitted to: U.S. Department of Energy, Bonneville Power Administration, Division of Fish and Wildlife, Project Number 2010-031-00; Contract Number 53238. Available at: <https://pisces.bpa.gov/release/documents/documentviewer.aspx?doc=P127156>
- Steele, C. A., E. C. Anderson, M. W. Ackerman, M. A. Hess, N. R. Campbell, S. R. Narum, and M. R. Campbell. 2013. A validation of parentage-based tagging using hatchery steelhead in the Snake River basin. *Canadian Journal of Fisheries and Aquatic Sciences* 70:1046–1054.
- Stephenson, J., M. Campbell, J., Hess et al. (2009). A centralized model for creating shared, standardized, microsatellite data that simplifies inter-laboratory collaboration. *Conservation Genetics*, 10, 1145–1149.
- Stoddard JW, Parsons JE & Nagler JJ 2005 Early onset of embryonic mortality in sub-fertile families of rainbow trout (*Oncorhynchus mykiss*). *Reproduction Fertility and Development* 17 785-790.

- Sumpter JP, Lebail PY, Pickering AD, Pottinger TG & Carragher JF 1991 The Effect of Starvation on Growth and Plasma Growth-Hormone Concentrations of Rainbow-Trout, *Oncorhynchus-Mykiss*. *General and Comparative Endocrinology* 83 94-102.
- Sutton SG, Bult TP & Haedrich RL 2000 Relationships among fat weight, body weight, water weight, and condition factors in wild Atlantic salmon parr. *Transactions of the American Fisheries Society* 129 527-538.
- Takei Y & McCormick SD 2012 Hormonal Control of Fish Euryhalinity. In *Fish Physiology*, pp 70-125. Eds SD McCormick, AP Farrell & CJ Brauner: Academic Press.
- Tannenbaum GS 1993 Genesis of Episodic Growth-Hormone Secretion. *Journal of Pediatric Endocrinology* 6 273-282.
- Tappel, P.D. and T.C. Bjornn. 1983. A New Method of Relating Size of Spawning Gravel to Salmonid Embryo Survival. Idaho Cooperative Fishery Research Unit. University of Idaho, Moscow, Idaho 83843.
- Taranger GL, Carrillo M, Schulz RW, Fontaine P, Zanuy S, Felip A, Weltzien FA, Dufour S, Karlsen O, Norberg B, et al. 2010 Control of puberty in farmed fish. *General and Comparative Endocrinology* 165 483-515.
- Taylor JF, Migaud H, Porter MJR & Bromage NR 2005 Photoperiod influences growth rate and plasma insulin-like growth factor-I levels in juvenile rainbow trout, *Oncorhynchus mykiss*. *General and Comparative Endocrinology* 142 169-185.
- Taylor JF, Porter MJR, Bromage NR & Migaud H 2008 Relationships between environmental changes, maturity, growth rate and plasma insulin-like growth factor-I (IGF-I) in female rainbow trout. *General and Comparative Endocrinology* 155 257-270.
- Thissen JP, Ketelslegers JM & Underwood LE 1994 Nutritional Regulation of the Insulin-Like Growth-Factors. *Endocrine Reviews* 15 80-101.
- Thissen JP, Underwood LE & Ketelslegers JM 1999 Regulation of insulin-like growth factor-I in starvation and injury. *Nutr Rev* 57 167-176.
- Thorpe JE 1994 Reproductive strategies in Atlantic salmon, *Salmo salar* L. *Aquaculture and Fisheries Management* 25 77-87.
- Thorpe JE 2007 Maturation responses of salmonids to changing developmental opportunities. *Marine Ecology Progress Series* 335 285-288.
- Thorpe JE, Mangel M, Metcalfe NB & Huntingford FA 1998 Modelling the proximate basis of salmonid life-history variation, with application to Atlantic salmon, *Salmo salar* L. *Evolutionary Ecology* 12 581-599.
- Trammell JIJ, Fast DE, Hatch DR, Bosch WJ, Branstetter R, Pierce AL, Blodgett JW & Frederiksen CR. 2016. Evaluating steelhead kelt treatments to increase iteroparous spawners in the Yakima River Basin. *North American Journal of Fisheries Management* 36 876-887.
- Tyler CR & Sumpter JP 1996 Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6 287-318.
- Tyler CR, Sumpter JP & Witthames PR 1990 The dynamics of oocyte growth during vitellogenesis in the rainbow trout (*Oncorhynchus mykiss*). *Biology of Reproduction* 43 202-209.
- Uchida K, Kajimura S, Riley LG, Hirano T, Aida K & Grau EG 2003 Effects of fasting on growth hormone/insulin-like growth factor I axis in the tilapia, *Oreochromis mossambicus*. *Comp Biochem Physiol A Mol Integr Physiol* 134 429-439.

- Whitehead C, Bromage NR & Breton B 1983 Changes in serum levels of gonadotropin, estradiol 17-beta and vitellogenin during the 1st and subsequent reproductive cycles of female rainbow trout. *Aquaculture* 34 317-326.
- Wilkinson RJ, Longland R, Woolcott H & Porter MJR 2010 Effect of elevated winter-spring water temperature on sexual maturation in photoperiod manipulated stocks of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 309 236-244.
- Wilkinson RJ, Porter M, Woolcott H, Longland R & Carragher JF 2006 Effects of aquaculture related stressors and nutritional restriction on circulating growth factors (GH, IGF-I and IGF-II) in Atlantic salmon and rainbow trout. *Comparative Biochemistry and Physiology* 145 214-224.
- Winans, G. A., M. M. Paquin, D. M. Van Doornik, B. M. Baker, P. Thornton, D. Rawding, A. Marshall, P. Moran, and S. Kalinowski. 2004. Genetic stock identification of steelhead in the Columbia River Basin: an evaluation of different molecular markers. *North American Journal of Fisheries Management* 24:672–685. doi:10.1577/M03-052.1.
- Withler, I. L. 1966. Variability in life history characteristics of steelhead trout (*Salmo gairdneri*) along the Pacific coast of North America. *Journal of the Fisheries Research Board of Canada* 23, 365–393.
- Wood AW, Duan C & Bern HA 2005a Insulin-like growth factor signaling in fish. *International Review of Cytology* 243 215-285.
- Wood AW, Duan C & Bern HA 2005b Insulin-like growth factor signaling in fish. *Int Rev Cytol* 243 215-285.
- Wootton RJ & Smith C 2015 *Reproductive Biology of Teleost Fishes*. West Sussex, UK: John Wiley & Sons, Ltd.
- Xu Y, Wang B, Liu X, Shi B & Zang K 2017 Evidences for involvement of growth hormone and insulin-like growth factor in ovarian development of starry flounder (*Platichthys stellatus*). *Fish Physiol Biochem* 43 527-537.
- Yamamoto Y, Adam Luckenbach J, Goetz FW, Young G & Swanson P 2011 Disruption of the salmon reproductive endocrine axis through prolonged nutritional stress: changes in circulating hormone levels and transcripts for ovarian genes involved in steroidogenesis and apoptosis. *General and Comparative Endocrinology* 172 331-343.
- Yamashita S & Melmed S 1986 Insulin regulation of rat growth hormone gene transcription. *J Clin Invest* 78 1008-1014.
- Zaldua N & Naya DE 2014 Digestive flexibility during fasting in fish: a review. *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 169 7-14.

Appendices

A: Master Kelt Tracking Table

										Treatment benefit relative to Hockersmith 1.66	
Strategy	Year	Location	# Collected	# Released		# @ Bonneville	Return Rate to Bonneville (%)	# @ Lower Granite Dam (or Prosser)	Return Rate to Lower Granite Dam (or Prosser) (%)	Transportation Benefit relative to in-river	Transportation Benefit relative to Bonneville natural
In-river	2005	Prosser	67	67		3	4.5			1.54	25.61
In-river	2006	Prosser	51	51		1	2.0			0.67	3.16
In-river	2007	Prosser	53	53		3	5.7			1.95	9.28
In-river	2008	Prosser	88	88		4	4.5			1.56	6.64
In-river	2009	Prosser	58	58		3	5.2			1.78	11.54
In-river	2010	Prosser	155	154		2	1.3			0.44	3.74
In-river	2011	Prosser	85	85		3	3.5			1.21	7.01
In-river	2012	Prosser	59	59		2	3.4			1.17	3.15
In-river	2013	Prosser	52	52		0	0.0			0.00	0.00
In-river	2014	Prosser	45	45		3	6.7			2.29	11.52
In-river	2015	Prosser	121	121		0	0.0	0	0.0	0.00	0.00
In-river	2016	Prosser	56	56		2	3.6	2	3.6	1.23	57.50
In-river	2017	Prosser	5	5		0	0.0	0	0.0	0.00	0.00
Disc. In 2018 Total and weighted mean	13		895	894		26	2.91	2	0.22	1.07	5.64
In-river	2009	Lower Granite	178	176		2	1.1	2	1.1	3.55	1.96
In-river	2010	Lower Granite	1410	1399		5	0.4	4	0.3	1.12	0.42
In-river	2011	Lower Granite	1633	1613		3	0.2	3	0.2	0.58	0.10
In-river	2012	Lower Granite	2098	2098		4	0.2	3	0.1	0.60	0.10
In-river	2013	Lower Granite	840	827		3	0.4	2	0.2	1.13	0.37
In-river	2014	Lower Granite	2584	2571		11	0.4	9	0.4	1.34	0.50
In-river	2015	Lower Granite	1195	1193		0	0.0	0	0.0	0.00	0.00
In-river	2016	Lower Granite	1841	1837		4	0.2	2	0.1	0.69	2.11
In-river	2017	Lower Granite	824	821		0	0.0	0	0.0	0.00	0.00
In-river	2018	Lower Granite	868	863		3	0.3	1	0.1	1.09	0.31
In-river	2019	Lower Granite	1062	1034		0	0	0	0	0.00	0.00
In-river	2020	Lower Granite	296	284		12	4.1	12	1.2	12.79	7.46
In-river	2021	Lower Granite	250	248		TBD 2022	TBD 2022	TBD 2022	TBD 2022	TBD 2022	TBD 2022
Total and weighted mean	12		15079	14964		47	0.3	38	0.2	0.92	0.37
In-river	2020	Little Goose	22	22		7	31.8	7	31.8	100.39	52.39
In-river	2021	Little Goose	14	14		TBD 2022	TBD 2022	TBD 2022	TBD 2022	TBD 2022	TBD 2022
Total and weighted mean	2		36	36							

Strategy	Year	Location	# Collected	# Released									
Transported Immature (Fall Release @ Hamilton Is.)	2019	Prosser	103	103									
Transported Immature (Fall Release @ Hamilton Is.)	2020	Prosser	52	52									
Transported Immature (Fall Release @ Hamilton Is.)	2021	Prosser	53	53									
total years	3		208	208									

											Treatment benefit relative to Hockersmith 1.66	Transportation Benefit relative to Bonneville natural
Strategy	Year	Location	# Collected	# Released	S @ release (%)	# Remature	Retained	skip remature				
Long-term	2000	Prosser	512	91	17.77							
Long-term	2001	Prosser	551	197	35.75							
Long-term	2002	Prosser	420	140	33.33							
Long-term	2003	Prosser	482	298	61.83							
Long-term	2004	Prosser	662	253	38.22							
Long-term	2005	Prosser	386	86	22.28							
Long-term	2006	Prosser	279	85	30.47							
Long-term	2007	Prosser	422	221	52.37							
Long-term	2008	Prosser	472	269	56.99							
Long-term	2009	Prosser	510	140	27.45	91						
Long-term	2010	Prosser	1157	404	34.92	101						
Long-term	2011	Prosser	680	223	32.79	120						
Long-term	2012	Prosser	550	340	61.82	275						
Long-term	2013	Prosser	546	266	48.72	166	41	8				
Long-term	2014	Prosser	481	292	60.71	149	96	22				
Long-term	2015	Prosser	1098	396	36.07	382	74	37				
Long-term	2016	Prosser	471	360	76.43	236	74	15				
Long-term	2017	Prosser	118	55	46.61	55	29	4				
Long-term	2018	Prosser	227	103	45.37	103	49	5				
Long-term	2019	Prosser	371	248	66.85	145	NA	NA				
Long-term	2020	Prosser	463	320	69.11	268	NA	NA				
Long-term	2021	Prosser	360	219	60.83	149	NA	NA				
Total and weighted mean	21		11218	5006	44.1	2240	363	91				

										Treatment benefit relative to Hockersmith 1.66		
Strategy	Year	Location	# Collected	# Survived	S @ release (%)	Released	# cons. remature	Retained	skip remature		Transportation Benefit relative to in-river	Transportation Benefit relative to Bonneville natural
Long-term	2011	Lower Granite	111	2	1.80	2	2	-	-		9.80	3.58
Long-term	2012	Lower Granite	124	10	8.06	10	10	0	-		42.30	7.49
Long-term	2013	Lower Granite/S.F. Clearwater	134	69	51.49	69	69	0	-		144.18	47.01
Long-term	2014	Lower Granite/Fish Cr.	122	37	30.33	35	35	2	2		71.24	52.41
Long-term	2015	Lower Granite/S.F. Clearwater/Fish Cr.	140	43	30.71	24	22	21	18		96.91	59.58
Long-term	2016	Lower Granite/S.F. Clearwater	227	120	52.86	37	19	101	77		243.30	851.10
Long-term	2017	Lower Granite	269	59	21.93	98	21	58	29		69.20	42.55
Long-term	2018	Lower Granite	259	177	68.34	79	50	99	1		197.73	102.05
Long-term	2019	Lower Granite	288	121	42.01	40	39	85	58		132.56	114.84
Long-term	2020	Lower Granite/Little Goose	137	71	51.82	66	8	63	61		163.51	100.53
Long-term	2021	Lower Granite/Little Goose	66	22	33.33	65	4	22	TBD 2022		105.17	64.66
Total and weighted mean	10		1877	731	39.1	525	279	451	185		123.52	75.94
Long-term	2020	Little Goose	49	18	36.7	1	1	17	3			
Long-term	2021	Little Goose	12	1	8.3	0	0	1	TBD 2022			
Total and weighted mean			61	19	31.15	1	1	18	3			

Strategy	Year	Location	# Collected		# @ Bonneville	Return Rate to Bonneville (%)
Natural repeat	2004	Bonneville Dam	1125		4	0.36
Natural repeat	2005	Bonneville Dam	572		1	0.17
Natural repeat	2006	Bonneville Dam	1452		9	0.62
Natural repeat	2007	Bonneville Dam	1967		12	0.61
Natural repeat	2008	Bonneville Dam	2630		18	0.68
Natural repeat	2009	Bonneville Dam	2454		11	0.45
Natural repeat	2010	Bonneville Dam	1740		6	0.34
Natural repeat	2011	Bonneville Dam	1391		7	0.50
Natural repeat	2012	Bonneville Dam	1486		16	1.08
Natural repeat	2013	Bonneville Dam	1278		14	1.10
Natural repeat	2014	Bonneville Dam	1728		10	0.58
Natural repeat	2015	Bonneville Dam	904		0	0.00
Natural repeat	2016	Bonneville Dam	1610		1	0.06
Natural repeat	2017	Bonneville Dam	837		0	0.00
Natural repeat	2018	Bonneville Dam	896		6	0.67
Natural repeat	2019	Bonneville Dam	820		3	0.37
Natural repeat	2020	Bonneville Dam	1222		4	0.33
Natural repeat	2021	Bonneville Dam	TBD 2022		TBD 2022	TBD 2022
Total	18		24112		122	0.52

B: Publications

Kelt steelhead have been largely understudied. Our initial interest came from the availability of kelt steelhead at the Chandler Juvenile Collection Facility on the Yakima River coupled with Atlantic salmon kelt reconditioning literature (Gray et al. 1987; Crim et al. 1992; Johnston et al. 1992). Below is a brief chronological bibliography of kelt steelhead-related literature with contributions associated with this project noted with an *.

Everman (1895) surveyed fish populations in the Snake River and noted local knowledge that steelhead don't die after spawning and must return to the sea.

Whitt (1954) estimated that approximately 2% of adult Clearwater River steelhead were repeat spawners. Unfortunately, Whitt's estimates were conducted after the construction of two hydropower facilities on main-stem river sites; the estimates also relied on scale analysis, which may have resulted in an underestimation of kelt abundance due to scale reabsorption.

[Busby et al.](#) (1996) reported iteroparity rates in excess of 17% for winter run steelhead in the Kalama River (lower Columbia River).

*[Evans](#) (2003) estimated that 17% of the Snake River steelhead spawning population was observed as kelts in the Lower Granite Dam juvenile by-pass facility during a 10-week monitoring period in 2000.

[Wertheimer and Evans](#) (2005) noted high kelt abundance and poor emigration survival of steelhead from upper Columbia River tributaries to the ocean as the underlying limiting factor inhibiting iteroparity.

[Evans et al.](#) (2008) identified transportation of downstream-migrating kelts around hydropower dams as a potential method to improve kelt survival.

[Keefer et al.](#) (2008) sampled steelhead kelts at main-stem hydropower dams from 2001 to 2004 and estimated that between 0.5% and 1.2% of those fish from the Snake River and tributaries and 2.9% to 9.0% of fish from the Lower Columbia exhibited an iteroparous life history.

*[Narum et al.](#) (2008) used microsatellite genotypes to identify kelt individuals to upstream populations of origin. Repeat spawning was negatively correlated with body size and iteroparity may be a valuable source of genetic variability and a conservation priority, especially in years with poor recruitment.

[Scott and Gill](#) (2008) anthropogenic modifications to the environment may select against repeat spawning.

[Seamons and Quinn](#) (2010) reported that repeat spawning adults have life-time reproductive success more than twice that of one-time spawners, and the average number of offspring produced by both male and female repeat spawners is much higher (1.9-time higher females and 2.7 times higher for males).

[Nielsen et al.](#) (2011) analyzed molecular genetic data across various life-history types (single versus repeat-spawning groups) of steelhead inhabiting the Ninilchik River in Alaska. They concluded that steelhead inhabiting this river formed a single panmictic spawning population, suggesting that iteroparity is a ubiquitous trait, potentially governed solely by environmental conditions that are more or less conducive to the survival of post-spawned adults. This is consistent with the fact that expression of iteroparity persists among interior Columbia River steelhead despite decades of passage-related selection against this life history type. Nonetheless, we are not aware of any study that has quantitatively established the presence or absence of specific genetic factors that contribute to iteroparity. Given the lack of conclusive evidence either supporting or negating a genetic component for iteroparity it is reasonable to conclude that enhancing the potential for iteroparity is a risk-averse strategy for steelhead.

*[Caldwell et al.](#) (2013) found that a feed restricted regime arrested reproductive development and affected factors associated with energy balance purported to play a role in initiating reproductive development within 2-4 months after spawning.

#[Courter et al.](#) (2013) found evidence for variable maternal resident contribution rates to anadromous returns based on otolith microchemistry relative to water chemistry in the Yakima River.

*[Hatch et al.](#) 2013 established the sex ratio of kelts being skewed toward female 92% over a collection 9,738 fish. Further reporting described the reconditioning process, survival rates, condition of fish, and release strategies.

[Null et al.](#) (2013) reconditioned post-spawned hatchery-origin steelhead for a short period of time at Coleman Hatchery on the Sacramento River and achieved return rates of 26%. Radio tracking showed that some of the reconditioned kelts did not migrate to the estuary and instead resided in the lower Sacramento River until the next spawning run.

*[Penney and Moffitt](#) (2013) findings indicate that the physiological processes causing rapid senescence and death in semelparous salmon are not evident in steelhead and recovery begins in freshwater.

*[Caldwell et al.](#) (2014) found that ration restriction after spawning arrested ovarian development by 10 weeks after spawning.

*[Buelow and Moffitt](#) (2014) found kelts in good external condition had plasma profiles similar to downstream-migrating smolts. In addition, they found more than 80% of kelts ranked in good external condition had smolt-like body silvering. Migrating kelts had Na⁺, K⁺ ATPase activity that was significantly elevated over hatchery fish at the time of spawning.

#[Hernandez, K., Copeland, T., Wright, K.](#) 2014. Quantitative Assessment of Scale Resorption in Migrating and Spawning Steelhead of the Snake River Basin. Transactions of the American Fisheries Society 143:1562-1568.

*[Penney and Moffitt](#) (2014a) found no evidence of significant interannual variation in lipid, protein, and energy density of sexually mature steelhead and steelhead kelts. Postspawning survival of steelhead is likely limited by the low energy density of kelts.

*[Penney and Moffitt](#) (2014b) investigated energy change in kelt steelhead and found liver fatty acids were depleted more rapidly than muscle.

[Moore et al.](#) (2014) iteroparity provides extra population resilience for anadromous salmonids. "Life history strategies can buffer individuals and populations from environmental variability. For instance, it is possible that asynchronous dynamics among different life histories can stabilize populations through portfolio effects."

[Keefer et al.](#) (2016) assessed sea-to-sea survival of 1,212 adult steelhead from the Snake and Columbia rivers. They found survival from Snake River tributary entry to postspawn kelt was 14% to 17% and kelt outmigration survival through the lower Snake River dams was 31% to 39% and 13% to 20% through all eight main-stem dams.

*[Matala et al.](#) (2016) identified kelt steelhead collected at Lower Granite Dam to A-run (82%) and B-run (18%) components.

*[Penney, Z.L., C.M. Moffitt, B. Jones, B. Marston.](#) 2016. Physiological comparisons of steelhead kelts emigrating from the Situk River, AK and Clearwater River, ID. Environmental Fish Biology Vol. 99 No. 4.

*[Trammell et al.](#) (2016) compared kelt reconditioning management strategies and found that long-term reconditioning resulted in approximately 5 times more repeat spawners than the next best strategy or control group.

*[Bosch et al.](#) (2017) provided an outreach article to The Osprey describing the benefits of steelhead kelt reconditioning to wild steelhead populations.

[Copeland et al.](#) (2017) reports on the life history diversity across management areas in the Snake River and finds broad overlap among populations in several areas forming a gradient in life history characteristics rather than a dichotomous break.

*[Pierce, A.L., J.W. Blodgett, T.D. Cavileer, L.R. Medeiros, J. Boyce, L.K. Caldwell, W.J. Bosch, R. Branstetter, D.E. Fast, D.R. Hatch, and J.J. Nagler.](#) 2017. Reproductive development in captive

reconditioned female steelhead kelts: evidence for consecutive and skip spawning life histories. Canadian Journal of Fisheries and Aquatic Sciences 74(7): 1049-1060.

*[Jenkins LE, Pierce AL, Graham N, Branstetter R, Hatch DR, and Nagler JJ](#). 2018. Reproductive performance and energy balance in consecutive and skip repeat spawning reconditioned female steelhead trout *Oncorhynchus mykiss*. Transactions of the American Fisheries Society.

[Copeland et al.](#) (2019) investigated patterns of iteroparity in wild Snake River steelhead and reported a greater percentage of skip spawners relative to consecutive spawners. Additionally, survival to a second spawning was greater for smaller fish.

*[Jenkins LE, Pierce AL, Graham ND, Medeiros LR, Hatch DR, and Nagler JJ](#). 2019. Elevated plasma triglycerides and growth are early indicators of reproductive status in post-spawning female steelhead trout *Oncorhynchus mykiss*. Conservation Physiology.

*[Jenkins et al.](#) (2020) found no support for the idea of a trade-off between current and future reproduction instead reproductive effort and its allocation to egg size and fecundity depended on energetic status during early oogenesis.

*[Medeiros et al.](#) (2020) developed a cost-comparable, non-radioisotope method for quantifying salmonid plasma growth hormone and IGF-I using commercially available reagents.

*Indicates articles directly associated with BPA project 2007-401-00.

#Indicates articles where material contributions were made from BPA project 2007-401-00.

Presentations 2021

Bosch B. Yakima Basin steelhead kelt reconditioning: Program overview. Yakima Chapter of Trout Unlimited, Zoom Presentation, December 15, 2021.

Hatch D. Kelt Reconditioning and Reproductive Success Evaluation Research. Independent Science Review Panel: Anadromous Fish Habitat and Hatchery Review Presentations, Go-to-meeting, July 15, 2021.

C: List of Metrics and Indicators

Data Collection Methods

Air Spawning: <https://www.monitoringresources.org/Document/Method/Details/5343>

Coloration Rating: <https://www.monitoringresources.org/Document/Method/Details/5302>

Determining Adult Anadromous Salmonid Gender:

<https://www.monitoringresources.org/Document/Method/Details/1429>

Determining Sex of Adult Steelhead:

<https://www.monitoringresources.org/Document/Method/Details/5334>

DNA Extraction Kit Protocol:

<https://www.monitoringresources.org/Document/Method/Details/1353>

Downloading Data from PTAGIS:

<https://www.monitoringresources.org/Document/Method/Details/4095>

Electrofishing- Determine Electrofisher Settings:

<https://www.monitoringresources.org/Document/Method/Details/115>

Electrofishing - Fish Processing and Recovery:

<https://www.monitoringresources.org/Document/Method/Details/117>

Estimating Lipid Content in Muscle Tissue of Adult Salmonids:

<https://www.monitoringresources.org/Document/Method/Details/4215>

Extracting Fish Plasma to Measure Reproductive Development:

<https://www.monitoringresources.org/Document/Method/Details/4239>

Fish Wet Weight: <https://www.monitoringresources.org/Document/Method/Details/1734>

Genetic Sampling and Storage Using Chromatography Filter:

<https://www.monitoringresources.org/Document/Method/Details/4087>

Identifying Marks/Tags on Fish:

<https://www.monitoringresources.org/Document/Method/Details/342>

Identifying Steelhead Kelts:

<https://www.monitoringresources.org/Document/Method/Details/5310>

Measuring Fish Length- Fork length:

<https://www.monitoringresources.org/Document/Method/Details/4041>

Measuring Fish Length- Mid-orbital hypural length:

<https://www.monitoringresources.org/Document/Method/Details/1549>

Modified Assessment of Fish Condition:

<https://www.monitoringresources.org/Document/CustomizedMethod/Details/22915>

Modified Off Ladder Adult Trapping Procedures:

<https://www.monitoringresources.org/Document/CustomizedMethod/Details/31061>

Modified PIT Tag Marking Procedures:

<https://www.monitoringresources.org/Document/CustomizedMethod/Details/22818>

O. mykiss and O. tshawytscha SNP Marker Sets for PBT and GSI Use in the Columbia River Basin:

<https://www.monitoringresources.org/Document/Method/Details/1356>

SNP Genotyping on Fluidigm Platform:

<https://www.monitoringresources.org/Document/Method/Details/1332>

Tissue Sampling for Parentage Based Tagging:

<https://www.monitoringresources.org/Document/Method/Details/1432>

Tissue Sampling of Salmonids in Nature for Genetic Analysis:

<https://www.monitoringresources.org/Document/Method/Details/933>

Weights: Green Egg Mass:

<https://www.monitoringresources.org/Document/Method/Details/1457>

Weights: Sampling for Green Egg Fecundity Estimate:

<https://www.monitoringresources.org/Document/Method/Details/1458>

Data Analysis/Interpretation Methods

Assessing Genetic Population Structure Using Bayesian Clustering Methods:

<https://www.monitoringresources.org/Document/Method/Details/1351>

Calculating a Mean, Variance and Standard Deviation:

<https://www.monitoringresources.org/Document/Method/Details/4245>

Estimating Relative Reproductive Success (RRS):

<https://www.monitoringresources.org/Document/Method/Details/696>

Estradiol Assay of Fish Plasma Samples:

<https://www.monitoringresources.org/Document/Method/Details/5320>

Fulton's Fish Condition Factor:

<https://www.monitoringresources.org/Document/Method/Details/952>

Genetic Assignment Tests Using GeneClass2:

<https://www.monitoringresources.org/Document/Method/Details/487>

Genetic Stock Mixture Analysis Using the Software Program BAYES:

<https://www.monitoringresources.org/Document/Method/Details/488>

Modified Analysis of Variance Models:

<https://www.monitoringresources.org/Document/CustomizedMethod/Details/22904>

Modified Binary Logistic Regression of Multi-Year Monitoring Data:

<https://www.monitoringresources.org/Document/CustomizedMethod/Details/31060>

Modified Calculating the smolt to adult return rate (SAR):

<https://www.monitoringresources.org/Document/CustomizedMethod/Details/22932>

Modified Growth Rate for Individual Fish:

<https://www.monitoringresources.org/Document/CustomizedMethod/Details/31057>

Modified Integrated Status and Effectiveness Monitoring Program Salmonid Life Cycle Modeling:

<https://www.monitoringresources.org/Document/CustomizedMethod/Details/31063>

Monitor Survival and Mortality Rates of Fish:

<https://www.monitoringresources.org/Document/Method/Details/3992>

Parentage Analysis Using Cervus:

<https://www.monitoringresources.org/Document/Method/Details/1430>

Predicting the Accuracy of Genetic Stock Identification:

<https://www.monitoringresources.org/Document/Method/Details/1346>

Solvent Extraction of Plasma Samples:

<https://www.monitoringresources.org/Document/Method/Details/5319>

Tests of Significance: T-Test:

<https://www.monitoringresources.org/Document/Method/Details/5291>

Vitellogenin Assay of Fish Plasma Samples:

<https://www.monitoringresources.org/Document/Method/Details/5335>

Metrics

Title	Category	Subcategory	Subcategory Focus 1	Subcategory Focus 2
"Stock composition"	Fish	Composition: Fish Species Assemblage (ID: 56)	Fish Life Stage: Adult - Outmigrant	Fish Origin: Natural
"Kelt condition"	Fish	Condition of Life Stage: Fish (ID: 57)	Fish Life Stage: Adult - Outmigrant	NA
"Maturation rate"	Fish	Condition of Life Stage: Fish (ID: 57)	Fish Life Stage: Adult - Returner	NA
"Reconditioned kelt condition"	Fish	Condition of Life Stage: Fish (ID: 57)	Fish Life Stage: Adult Fish	NA
"Kelt homing"	Fish	Distribution of Fish Species (ID: 62)	Fish Life Stage: RANGE: Adult to Adult	NA
"Fecundity"	Fish	Fecundity: Fish (ID: 68)	NA	NA
"Growth rate"	Fish	Growth Rate: Fish (ID: 73)	Fish Life Stage: RANGE: Juvenile to Adult	NA
"Kelt length"	Fish	Length: Fish Species (ID: 75)	Fish Life Stage: Adult - Outmigrant	NA
"Reconditioned kelt length"	Fish	Length: Fish Species (ID: 75)	Fish Life Stage: Adult Fish	NA
"Lipid content"	Fish	Lipid Content (ID: 200)	NA	NA
"Parentage analysis"	Fish	Relative Reproductive Success (RRS) (ID: 88)	Fish Origin: Both	NA
"Reproductive success"	Fish	Reproductive Success (Nb/N) (ID: 89)	Fish Origin: Natural	NA
"Reconditioned kelt survival rate"	Fish	Survival Rate: Fish (ID: 99)	Fish Life Stage: Adult - Outmigrant	Fish Origin: Both
"Natural kelt survival rate"	Fish	Survival Rate: Fish (ID: 99)	Fish Life Stage: Adult - Outmigrant	Fish Origin: Both
"Kelt weight"	Fish	Weight: Fish (ID: 206)	Fish Life Stage: Adult - Outmigrant	Fish Origin: Both

"Reconditioned kelt weight"	Fish	Weight: Fish (ID: 206)	Fish Life Stage: Adult Fish	Fish Origin: Both
-----------------------------	----------------------	--	---	-----------------------------------

Indicators

Title	Category	Subcategory	Subcategory Focus 1
"Relative reproductive success of artificially reconditioned kelt steelhead"	Fish	Relative Reproductive Success (RRS) (ID: 88)	Fish Origin: Natural